

**GESAMP**

Joint Group of Experts on the
Scientific Aspects of Marine
Environmental Protection

SOURCES, FATE AND EFFECTS OF MICROPLASTICS IN THE MARINE ENVIRONMENT: A GLOBAL ASSESSMENT



IMO



FAO



UNESCO



IOC



WMO



UNIDO



IAEA



UN



UNEP



UNDP



SOURCES, FATE AND EFFECTS OF MICROPLASTICS IN THE MARINE ENVIRONMENT: A GLOBAL ASSESSMENT

Published by the
INTERNATIONAL MARITIME ORGANIZATION
4 Albert Embankment, London SE1 7SR
www.imo.org

Printed by Polestar Wheatons (UK) Ltd, Exeter, EX2 8RP

ISSN: 1020-4873

Cover photo: Microplastic fragments from the western North Atlantic, collected using a towed plankton net.
Copyright Giora Proskurowski, SEA

Notes:

GESAMP is an advisory body consisting of specialized experts nominated by the Sponsoring Agencies (IMO, FAO, UNESCO-IOC, UNIDO, WMO, IAEA, UN, UNEP, UNDP). Its principal task is to provide scientific advice concerning the prevention, reduction and control of the degradation of the marine environment to the Sponsoring Agencies.

The report contains views expressed or endorsed by members of GESAMP who act in their individual capacities; their views may not necessarily correspond with those of the Sponsoring Agencies.

Permission may be granted by any of the Sponsoring Agencies for the report to be wholly or partially reproduced in publication by any individual who is not a staff member of a Sponsoring Agency of GESAMP, provided that the source of the extract and the condition mentioned above are indicated.

Information about GESAMP and its reports and studies can be found at: <http://gesamp.org>

ISSN 1020-4873 (GESAMP Reports & Studies Series)

Copyright © IMO, FAO, UNESCO-IOC, UNIDO, WMO, IAEA, UN, UNEP, UNDP 2015

For bibliographic purposes this document should be cited as:

GESAMP (2015). "Sources, fate and effects of microplastics in the marine environment: a global assessment" (Kershaw, P. J., ed.). (IMO/FAO/UNESCO-IOC/UNIDO/WMO/IAEA/UN/UNEP/UNDP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection). Rep. Stud. GESAMP No. 90, 96 p.

Report editor: Peter Kershaw

Contributors to the report:

Alison Anderson, Anthony Andrady, Courtney Arthur, Joel Baker, Henk Bouwman, Sarah Gall, Valeria Hidalgo-Ruz, Angela Köhler, Kara Lavender Law, Heather Leslie, Peter Kershaw, Sabine Pahl, Jim Potemra, Peter Ryan, Won Joon Shim, Richard Thompson, Hideshige Takada, Alexander Turra, Dick Vethaak and Kayleigh Wyles.

Contents

Page

EXECUTIVE SUMMARY	5
ACKNOWLEDGMENTS.....	7
1 BACKGROUND TO GESAMP ASSESSMENT.....	9
1.1 Microplastics in the ocean – an emerging issue of international concern	9
1.2 GESAMP response	11
1.2.1 Scoping activities	11
1.2.2 Working Group 40	11
2 ASSESSMENT FRAMEWORK.....	12
2.1 DPSIR Conceptual model	12
2.2 Scope of assessment	12
3 SOURCES AND FATE OF MICROPLASTICS IN THE MARINE ENVIRONMENT	14
3.1 Introduction	14
3.1.1 Defining ‘plastic’.....	14
3.1.2 Defining ‘microplastics’	14
3.1.3 Origin and types of plastic.....	14
3.2 ‘Primary’ and ‘Secondary’ microplastics	18
3.2.1 Generation of microplastics.....	18
3.2.2 Weathering degradation of plastics in the ocean	18
3.2.3 Fragmentation of plastics	19
3.2.4 Biodegradation and Mineralization	19
3.3 Sampling methods for microplastics in the marine environment	19
3.3.1 Introduction	19
3.3.2 Sampling seawater.....	20
3.3.3 Sampling sediments	21
3.3.4 Sampling biota	21
3.4 Determining the composition of microplastics	21
3.5 Distribution of microplastics in the marine environment	22
3.5.1 Influence of the source.....	22
3.5.2 Distribution of microplastics based on direct observations	23
3.5.3 Transport pathways	27
3.5.4 Modelling the transport and distribution of microplastics	28
3.6 Recommendations for further research	29
4 EFFECTS OF MICROPLASTICS ON MARINE BIOTA.....	30
4.1 Introduction	30
4.2 Exposure	30
4.2.1 Exposure through the gills	30
4.2.2 Ingestion.....	30
4.2.3 Uptake and transition into tissues, cells and organelles.....	43
4.2.4 Excretion	44
4.2.5 Transfer of microplastics in the food web	45
4.3 Microplastics as a vector of chemical transport into marine organisms	45
4.3.1 Introduction	45

4.3.2	Contaminant transfer from μm -size plastics to lower-trophic-level organisms	45
4.3.3	Field evidence of contaminant transfer from mm-size plastics to higher-trophic-level organisms	48
4.4	Biological impacts	48
4.4.1	Physical effects	49
4.4.2	Comparison with observed effects in mammalian systems	49
4.4.3	Chemical effects	51
4.4.4	Potential effects on populations, communities and ecosystems	51
4.4.5	Potential effects on humans	52
4.5	Recommendations for further research	53
5	SOCIAL ASPECTS OF MICROPLASTICS IN THE MARINE ENVIRONMENT	53
5.1	Introduction	53
5.2	Perceptions of marine litter and microplastics	54
5.2.1	Perceptions of marine litter in general	54
5.2.2	Perceptions about microplastics	55
5.2.3	Public perceptions and coverage in the printed and digital media	56
5.2.4	Perceived Risks	58
5.3	Social and socio-economical impacts	60
5.4	The role of individual, group and regional differences	61
5.5	Overcoming barriers and towards solutions	62
5.5.1	Education and Public Engagement	63
5.6	Recommendations for further research	63
6	KEY OBSERVATIONS AND CONCLUSIONS	64
6.1	Sources, distribution and fate of microplastics	64
6.2	Effects	64
6.3	Social aspects	65
7	KEY POLICY-RELATED RECOMMENDATIONS	65
7.1	Rationale	65
7.2	Action-orientated recommendations addressing marine microplastics	66
7.2.1	Challenge 1	66
7.2.2	Challenge 2	66
7.2.3	Challenge 3	67
7.3	Recommendations to improve a future assessment	67
7.3.1	Challenge 4	67
7.3.2	Challenge 5	68
7.3.3	Challenge 6	68
	REFERENCES	69
	ANNEX I – MEMBERSHIP OF THE WORKING GROUP	92
	ANNEX II – LIST OF GESAMP REPORTS AND STUDIES	93

EXECUTIVE SUMMARY

Society has used the ocean as a convenient place to dispose of unwanted materials and waste products for many centuries, either directly or indirectly via rivers. The volume of material increased with a growing population and an increasingly industrialized society. The demand for manufactured goods and packaging, to contain or protect food and goods, increased throughout the twentieth century. Large-scale production of plastics began in the 1950s and plastics have become widespread, used in a bewildering variety of applications. The many favourable properties of plastics, including durability and low cost, make plastics the obvious choice in many situations. Unfortunately, society has been slow to anticipate the need for dealing adequately with end-of-life plastics, to prevent plastics entering the marine environment. As a result there has been a substantial volume of debris added to the ocean over the past 60 years, covering a very wide range of sizes (metres to nanometres in diameter). This is a phenomenon that has occurred wherever humans live or travel. As a result there are multiple routes of entry of plastics into the ocean, and ocean currents have transported plastics to the most remote regions. It is truly a global problem.

The GESAMP assessment focuses on a category of plastic debris termed 'microplastics'. These small pieces of plastic may enter the ocean as such, or may result from the fragmentation of larger items through the influence of UV radiation. Section 1 provides an introduction to the problem of microplastics in the marine environment, and the rationale for the assessment. The principal purpose of the assessment is to provide an improved evidence base, to support policy and management decisions on measures that might be adopted to reduce the input of microplastics to the oceans. The GESAMP assessment can be considered as contributing to a more formal Assessment Framework, such as the Driver-Pressure-State-Impact-Response (DPSIR) Assessment Framework, which is introduced in Section 2.

The nature of man-made polymers, different types and properties of common plastics and their behaviour in the marine environment are introduced in Section 3. There is no internationally agreed definition of the size below which a small piece of plastic should be called a microplastic. Many researchers have used a definition of <5 mm, but this encompasses a very wide range of sizes, down to nano-scales. Some microplastics are purposefully made to carry out certain functions, such as abrasives in personal care products (e.g. toothpaste and skin cleaners) or for industrial purposes such as shot-blasting surfaces. These are often termed 'primary' microplastics. There is an additional category of primary particle known as a 'pellet'. These are usually spherical or cylindrical, approximately 5 mm in diameter, and represent the common form in which newly produced plastic is transported between plastic producers and industries which convert the simple pellet into a myriad of different types of product.

The potential physical and chemical impacts of microplastics, and associated contaminants, are discussed in detail in Section 4. The physical impacts of larger litter items, such as plastic bags and fishing nets, have been demonstrated, but it is much more difficult to attribute physical impacts of microplastics from field observations. For this reason researchers have used laboratory-based experimental facilities to investigate particle uptake, retention and effects. Chemical effects are even more difficult to quantify. This is partly because seawater, sediment particles and biota are already contaminated by many of the chemical substances also associated with plastics. Organic contaminants that accumulate in fat (lipids) in marine organisms are absorbed by plastics to a similar extent. Thus the presence of a contaminant in plastic fragments in the gut of an animal and the measurement of the same contaminant in tissue samples does not imply a causal relationship. The contaminant may be there due to the normal diet. In a very small number of cases, contaminants present in high concentrations in plastic fragments with a distinctive chemical 'signature' (a type of flame retardant) can be separated from related contaminants present in prey items and have been shown to transfer across the gut. What is still unknown is the extent to which this might have an ecotoxicological impact on the individual.

It is recognized that people's attitudes and behaviour contribute significantly to many routes of entry of plastics into the ocean. Any solutions to reducing these sources must take account of this social dimension, as attempts to impose regulation without public understanding and approval are unlikely to be effective. Section 5 provides an opportunity to explore issues around public perceptions towards the ocean, marine litter, microplastics and the extent to which society should be concerned. Research specifically on litter is rather limited, but useful analogies can be made with other environmental issues of concern, such as radioactivity or climate change.

Section 6 summarizes some of the main observations and conclusions, divided into three sections: i) sources, distribution and fate; ii) effects; and, iii) social aspects. Statements are given a mark of high, medium or low confidence. A common theme is the high degree of confidence in what we do not know.

The assessment report concludes (Section 7) with a set of six Challenges and related Recommendations. Suggestions for how to carry out the recommendations are provided, together with a briefing on the likely consequences of not taking action. These are divided into three Action-orientated recommendations and three recommendations designed to improve a future assessment:

Action-orientated recommendations:

- Identify the main sources and categories of plastics and microplastics entering the ocean.
- Utilize end-of-plastic as a valuable resource rather than a waste product.

- Promote greater awareness of the impact of plastics and microplastics in the marine environment.

Recommendations for improving a future assessment:

- Include particles in the nano-size range.
- Evaluate the potential significance of plastics and microplastics as a vector for organisms.
- Address the chemical risk posed by ingested microplastics in greater detail.

ACKNOWLEDGMENTS

The report was subject to rigorous review prior to publication, including the established internal review process by Members of GESAMP. In addition, we wish to gratefully acknowledge the invaluable critical review, insights and suggestions made by an independent group of acknowledged experts in their fields: Francois Galgani, K. Irvine, Branden Johnson, Chelsea Rochman, Peter Ross and Martin Thiel. The review process greatly improved the content and presentation of the final report. However, any factual errors or misrepresentation of information remains the responsibility of the principal editor.

The following organizations provided in-kind or financial support to the working group: IOC of UNESCO, IMO, UNIDO, UNEP, NOAA. In addition, the American Chemistry Council (ACC) and Plastics Europe (PE) provided financial support, without which the Working Group could not have functioned. Luis Valdes (IOC), Edward Kleverlaan, Fredrik Haag and Jennifer Rate (IMO) provided invaluable encouragement and in-kind support.

Ashley Carson (ACC), Keith Christman (ACC), Roberto Gomez (PE) and Ralph Schneider (PE) provided encouragement and technical advice on the plastics industry and related matters.

1 BACKGROUND TO GESAMP ASSESSMENT

1.1 Microplastics in the ocean – an emerging issue of international concern

Marine debris from natural sources, such as floating vegetation or volcanic ash deposits (tuff), is commonplace in the ocean. Unfortunately, man-made debris has increased substantially, particularly in the past hundred years. Marine debris, or litter, from non-natural sources is usually defined as ‘any persistent, manufactured or processed solid material discarded, disposed of or abandoned in the marine and coastal environment’ (Galgani et al. 2010). This includes items that have been made or used by people and deliberately discarded or unintentionally lost directly into the sea, or on beaches, and materials transported into the marine environment from land by rivers, drainage or sewage systems or by wind transport. Such items may consist of metal, glass, paper, fabric or plastic. Of these, plastic is considered to be the most persistent and problematic.

Larger plastic objects are readily visible and many types of negative social, economic and ecological impact have been demonstrated, ranging from the entanglement of wildlife in fishing gear to the blockage of cooling water intakes on boats, requiring intervention by the rescue services. Smaller plastic objects, by definition, are less visible to the casual observer. Potential negative impacts are less obvious. However, reports of floating plastic micro-debris in the North Atlantic were first published in the scientific literature in the early 1970s. These publications raised concerns about the likelihood of ingestion of plastic particles by organisms and the potential for adverse physical and chemical impacts. A number of further publications, some in the ‘grey’ literature, in the 1970s and 1980s confirmed the occurrence of plastic particles in the North Pacific, Bering Sea and Japan Sea. But, the topic was largely ignored by the wider scientific and non-scientific community for many years.

Up until relatively recently, there has been a widespread tendency to treat the ocean as a convenient place to dispose of all sorts of unwanted material, either deliberately or unwittingly. Of course, the ocean is of finite capacity and persistent substances will tend to be transported globally and accumulate in certain locations, under the influence of oceanic and atmospheric transport. Plastics began to enter the ocean in increasing quantities from the 1950s, from a wide variety of land- and sea-based sources. There are no reliable estimates of inputs at a regional or global scale, but it is reasonable to assume that the total quantities have increased, even if the rate of increase is unknown.

National and international concern started to increase

in the mid 1980s. For example, the National Oceanic and Atmospheric Administration (NOAA) of the USA initiated the first of a series of international conferences of marine debris, held at regular intervals with the most recent (the fifth, 5IMDC) taking place in 2011. Increasingly this has included plastic particles as an important category of marine litter. The Honolulu Strategy was launched at 5IMDC.¹ NOAA has been at the forefront of developing assessment guidelines and funding initiatives designed to reduce the impact of marine litter (Arthur & Baker 2009; 2012).² At a regional level the European Union has adopted the Marine Strategy Framework Programme, with marine litter as one of 11 descriptors of environmental state. A Technical Support Group was set up to guide the selection of sampling methods and assessment strategies, including microplastics.

At an international level, marine litter was one of the categories incorporated in the 1995 Washington Declaration concerning a Global Programme of Action (GPA) for the protection of the marine environment from land-based sources (UNEP 1995). It was listed as being of concern in the GESAMP 71 report on land-based activities (GESAMP 2001). More recently, the problem of marine debris, and the need for increased national and international control, was dealt with at the 60th session of UNGA within the Oceans and the Law of the Sea session (UNGA 2005); paragraphs 65-70).

A more definitive assessment was provided by the analytical overview of marine litter, initiated by UNEP with input from IOC, IMO and FAO (UNEP 2005). This provided a useful overview of the issue, including type, source and distribution of litter, and measures to combat the problem. FAO has expressed concern over lost, abandoned or otherwise discarded fishing gear and has addressed this issue through a correspondence group with IMO and in a joint study with UNEP (Macfadyen et al. 2009). UNEP pursued this issue within the Regional Sea Programme and published a review of their global initiative on marine litter in 2009 (UNEP 2009) together with a series of Regional Sea status reports with regards to marine litter. Subsequently, marine debris was one of three topics selected for inclusion in the 2011 UNEP Year Book, with specific emphasis on microplastics as an emerging issue of environmental concern (UNEP 2011). The MARPOL Convention (International Convention for the Prevention of Pollution from Ships; MARPOL 1973), governed by IMO, covers the disposal of solid wastes under Annex V. Annex V specifically prohibits the disposal of any plastic waste anywhere in the world ocean.

¹ <http://5imdc.wordpress.com/about/honolulustrategy/>

² <http://marinedebris.noaa.gov>

Marine litter was raised as an issue of concern at the UN Conference on Sustainable Development in 2012 (Rio+20). This resulted in a specific reference to marine litter in the outcome document (paragraph 163, A/RES/66/288), 'The future we want':

163. We note with concern that the health of oceans and marine biodiversity are negatively affected by marine pollution, including marine debris, especially plastic, persistent organic pollutants, heavy metals and nitrogen-based compounds, from a number of marine and land-based sources, including shipping and land run-off. We commit to take action to reduce the incidence and impacts of such pollution on marine ecosystems, including through the effective implementation of relevant conventions adopted in the framework of the International Maritime Organization (IMO), and the follow-up of the relevant initiatives such as the Global Programme of Action for the Protection of the Marine Environment from Land-based Activities, as well as the adoption of coordinated strategies to this end. We further commit to take action to, by 2025, based on collected scientific data, achieve significant reductions in marine debris to prevent harm to the coastal and marine environment.

The Global Partnership on Marine Litter³ was launched during Rio+20. This UNEP-led initiative is designed to encourage all sectors of governance, business, commerce and society to work together to bring about a reduction in the input of marine litter, especially plastics, into the ocean. The problem of marine litter and microplastics was raised at the First UN Environment Assembly, which took place in Nairobi in June 2014. This resulted in agreement on a Resolution on 'Marine plastic debris and microplastics' (UNEP 2014). The resolution referred specifically to the GESAMP assessment on microplastics (the subject of this report):

'13. Welcomes the initiative by the Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection to produce an assessment report on microplastics, which is scheduled to be launched in November 2014;

15. Requests the Executive Director, in consultation with other relevant institutions and stakeholders, to undertake a study on marine plastic debris and marine microplastics, building on existing work and taking into account the most up-to-date studies and data, focusing on:

- (a) Identification of the key sources for marine plastic debris and microplastics;*
- (b) Identification of possible measures and best available techniques and environmental practices to prevent the accumulation and minimize the level of microplastics in the marine environment;*
- (c) Recommendations for the most urgent actions;*

³ <http://gpa.unep.org/index.php/global-partnership-on-marine-litter>

(d) Specification of areas especially in need of more research, including key impacts on the environment and on human health;

(e) Any other relevant priority areas identified in the GESAMP assessment of the Joint Group of Experts described in paragraph [13].'

In a related development, the Global Partnership for Oceans was also launched at Rio+20.⁴ The World Bank acts as the Secretariat, with over 140 partners representing governments, NGOs, international organizations and the private sector. Marine litter is one of three (waste water and excess nutrients) priority topics under the pollution theme.

An assessment of floating plastic marine debris, and contaminants contained in plastic resin pellets, forms part of the Transboundary Waters Assessment Programme (TWAP), a full-size project (2013–2014) co-financed by the Global Environmental Facility.⁵ There has been coordination between the work on marine plastics and microplastics carried out under the TWAP and the work of WG40. The TWAP report is due for publication in early 2015.

Concern at an institutional level has been matched by a surge in interest amongst the academic community, with publications increasing almost exponentially over the past 5 years (Figure 1.1). The term 'microplastics' entered the popular lexicon at the start of this period of increased activity. Several regional seas organizations (e.g. NOWPAP, UNEP-MAP, OSPAR, HELCOM) have produced guidelines for assessing marine litter, including microplastics, in recent years, and some have organized regional workshops to encourage capacity building and the spread of good practice. The plastics production industry has developed a Joint 'Declaration of the Global Plastics Associations for Solutions on Marine Litter',⁶ launched at the 5IMDC in 2011. This includes a commitment to support a number of litter assessment and litter reduction programmes.⁷ Several Non-Governmental Organizations (NGOs), also referred to as Not-for-Profit organizations, have developed programmes both to raise awareness and help to quantify the extent of microplastic contamination and effects, at a national, regional and international scale. This has provided valuable additional information, often in a very cost-effective manner.

⁴ <http://www.globalpartnershipforoceans.org/about>

⁵ <http://geftwap.org>

⁶ <http://www.marinelittersolutions.com/who-we-are/joint-declaration.aspx>

⁷ Plastics Europe and the American Chemistry Council have jointly supported the GESAMP Working Group 40 on microplastics.




Figure 1.1 Publications by year, 1970 – July 2014, using the search terms ‘plastic pellets’ and ‘microplastics’ – compiled by Sarah Gall, Univ. Plymouth, UK.

1.2 GESAMP response

1.2.1 Scoping activities

The topic of microplastics in the marine environment was raised as part of GESAMP’s Emerging Issues programme in 2008. It was agreed to produce a scoping paper, published as an internal document in 2009. This was followed by a scoping workshop in June 2010, hosted by UNESCO-IOC in Paris, entitled: ‘International Workshop on Microplastic particles as a vector in transporting persistent, bioaccumulating and toxic substances in the ocean’. The proceedings were published in the GESAMP Reports & Studies Series (GESAMP 2010). One of the recommendations of the workshop was GESAMP should initiate a Working Group to conduct an assessment of microplastics in the coastal and open ocean, resulting in the setting up of WG40.

1.2.2 Working Group 40

Institutional support

Lead Agency: UNESCO-IOC

Other supporting UN Agencies: IMO, UNIDO

Other supporting organizations: NOAA, American Chemicals Council, Plastics Europe

Terms of reference

(as agreed at the 39th Session of GESAMP, April 2012, UNDP, New York)

1. Assess inputs of microplastic particles (e.g. resin pellets, abrasives, personal care products) and macroplastics (including main polymer types) into the ocean; to include pathways, developing methodology, using monitoring data, identifying proxies (e.g. population centres, shipping routes, tourism revenues);
2. Assess modelling of surface transport, distribution & areas of accumulation of plastics and microplastics, over a range of space- and time-scales;
3. Assess processes (physical, chemical & biological) controlling the rate of fragmentation and degradation, including estimating long-term behaviour and estimate rate of production of ‘secondary’ microplastic fragments;
4. Assess long-term modelling including fragmentation, seabed and water column distribution, informed by the results of ToR 3;
5. Assess uptake of particles and their contaminant/additive load by biota, as well as their physical and biological impacts at a population level;
6. Assess the socio-economic aspects, including public awareness.

Details of the working group members are provided in Annex I.

2 ASSESSMENT FRAMEWORK

2.1 DPSIR Conceptual model

The Driver-Pressure-State-Impact-Response (DPSIR) model has been used quite widely as a conceptual framework on which to base assessments of human impacts on both the terrestrial and aquatic environments (Ness et al. 2010). Drivers refer to high level demands society places on the environment, for example: energy supply, food security, transport, housing and recreation. In turn this may result in Pressures, or stresses, on the environment such as fisheries,

shipping, coastal tourism and waste generation. The State of the environment may change, for example by increasing levels of noise or introducing litter. This may lead to an Impact, for a Response may be implemented. Any response has to consider the cost-benefit trade-offs between the overriding Drivers and the desired reduction of the Impact. An example of the DPSIR model applied to the generation and impacts on marine litter, including microplastics, is provided in Figure 2.1.




Figure 2.1 DPSIR/DPSWR model as applied to the generation and potential impacts of marine litter.

There has been considerable debate, largely between natural scientists and social scientists and economists, about what constitutes an Impact. In natural sciences the injury or death on an organism might be construed as an Impact, whereas in socio-economics it is usually argued that Impacts imply a loss of an ecosystem service, i.e. a welfare Impact that society is concerned about. Attempts have been made to clarify this by replacing 'Impact' with 'Welfare impact' (i.e. DPSWR) (Cooper 2013), but this has not been universally accepted. The Response may include several formal and informal measures to mitigate the Impact, acting on one or more of the Driver, Pressure, State or Impact (Figure 2.2). The example used to illustrate this involves the impact of larger plastic items on turtles, as this provides an easily understood set of relationships. However, the model can also be applied to microplastics, although the Response pathways are likely to be much more complex.

2.2 Scope of assessment

The assessment was constrained by the Terms of Reference, to consider the sources, fate and effects of microplastics in the marine environment. It was designed to provide an improved evidence base for the intended audience, summarized as follows:

- UNESCO-IOC, IMO, UNIDO – main WG40 supporting agencies
- Other international agencies: e.g. UNEP, UNDP, FAO, World Bank, GEF, WHO, IWC
- Regional Seas Organizations, e.g. NOWPAP
- National and local governance bodies
- Other funding/development bodies
- Industry/commercial sectors, e.g. tourism, aquaculture, fisheries, retail, plastic producers, plastic recyclers
- General public, NGOs, school children etc.
- Social and natural scientists, economists




Figure 2.2 DPSIR/DPSWR model showing an example of potential Responses to reduce the Impact of marine litter on turtles.

The scope did not include providing recommendations for implementing specific measures to reduce the input of plastic and microplastics into the ocean. But, it is anticipated that the improved evidence base will inform such processes. One critical aspect of designing measures is to consider the potential economic loss associated with the loss of an ecosystem service, to allow a reliable cost-benefit analysis of any trade-offs that will

be necessary. This was outside the scope of the present assessment.

The working group considered three main topics, reflected in the structure of this report: Section 3 – sources and fate of microplastics; Section 4 – physical and chemical effects of microplastics; and, Section 5 – social aspects. These have been mapped onto the DPSIR model in Figure 2.3.




Figure 2.3 Coverage of the DPSIR/DPSWR model by the three main sections of the report: Section 3 – sources and fate of microplastics; Section 4 – physical and chemical effects of microplastics; and, Section 5 – social aspects.

3 SOURCES AND FATE OF MICROPLASTICS IN THE MARINE ENVIRONMENT

3.1 Introduction

3.1.1 Defining 'plastic'

Plastic is a term used in many fields, to describe the physical properties and behaviour of materials (e.g. soils, geological formations) as well as the name of a class of materials. The term 'plastic' is used here to define a sub-category of the larger class of materials called polymers. Polymers are very large molecules that have characteristically long chain-like molecular architecture and therefore very high average molecular weights. They may consist of repeating identical units (homopolymers) or different sub-units in various possible sequences (copolymers). Those polymers that soften on heating, and can be moulded, are generally referred to as 'plastic' materials. These include both virgin plastic resin⁸ pellets (easily transported prior to manufacture of plastic objects) as well as the resins mixed (or blended) with numerous additives to enhance the performance of the material. Additives may typically include fillers, plasticizers, colorants, stabilizers and processing aids. In addition to the thermoplastics, marine debris also includes some thermoset materials such as polyurethane foams, epoxy resins and some coating films. Thermosets are cross-linked materials that cannot be re-moulded on heating. However, these too are generally counted within the category of 'plastics' in marine debris.

3.1.2 Defining 'microplastics'

Small pieces of floating plastics in the surface ocean were first reported in the scientific literature in the early 1970s (Carpenter and Smith 1972; Carpenter et al. 1972), and later publications described studies identifying plastic fragments in birds in the 1960s (Harper and Fowler 1987). It is unclear when the term 'microplastic' was first used in relation to marine debris. It was mentioned by Ryan and Moloney (1990) in describing the results of surveys of South African beaches, and in cruise reports of the Sea Education Association in the 1990s and by Thompson et al. (2004) describing the distribution of plastic fragments in seawater. No formal size definition was proposed at the time but generally the term implied material that could only be readily identified with the aid of a microscope. It has since become widely used to describe small pieces of plastic in the millimetre to sub-millimetre size range, although it has not been formally recognized.

Particles in the size range 1 nm to <5 mm were considered microplastics for the purposes of this assessment

A more scientifically rigorous definition of plastic pieces might refer to nano-, micro-, meso-, macro- and mega-size ranges, although this has not yet been formally proposed for adoption by the international research community (Figure 3.1). At present, the lack of an agreed nomenclature, together with practical difficulties of sampling and measuring different size ranges in the field, has encouraged the widespread adoption of microplastics as a generic term for 'small' pieces of plastic.

The size definition of microplastics was discussed at the first international research workshop on the occurrence, effects and fate of microplastic marine debris in 2008, hosted by NOAA (Arthur et al. 2009). The participants adopted a pragmatic definition, suggesting an upper size limit of 5mm. This was based on the premise that it would include a wide range of small particles that could readily be ingested by biota, and such particles that might be expected to present different kinds of threat than larger plastic items (such as entanglement).

It was also recognized that the size ranges reported in field studies are constrained by the sampling techniques employed. For example, many studies have reported concentrations and size ranges of microplastics based on sampling with a plankton net, typically with a mesh size of 330 microns. Material <330 microns will be under-sampled. In this report we consider all available evidence; for example, including that relating to impacts of nano-particles, even though such particles cannot be detected at present in the environment on a routine basis.

3.1.3 Origin and types of plastic

Many different types of plastic are produced globally, but the market is dominated by 6 classes of plastics: polyethylene (PE, high and low density), polypropylene (PP), polyvinyl chloride (PVC), polystyrene (PS, including expanded EPS), polyurethane (PUR) and polyethylene terephthalate (PET). Plastics are usually synthesized from fossil fuels, but biomass can also be used as feedstock. The production chain for the most common artificial and natural polymers is illustrated in Figure 3.2. The figure includes some examples of common applications, including the manufacture of microplastics for particular industrial or commercial applications.

⁸ Resin – a semi-viscous liquid capable of hardening permanently http://en.wikipedia.org/wiki/Synthetic_resin




Figure 3.1 Size range of plastic objects observed in the marine environment and some comparisons with living material. These size distinctions could form the basis of a more rigorous description.

The availability of bio-based raw materials (feedstock) is expected to increase in the near future, providing alternative feedstock to fossil fuel raw materials. Being bio-based, however, does not necessarily make the plastic biodegradable; in fact, bio-based resins such as bio-PE or bio-PET are developed to mirror the properties of their conventional counterparts to allow same lifetime, applications and recycling capabilities. They are drop-in substitutes for the conventional plastic resin with the same structure. For instance a very small fraction of bio-based PET resin or bio-PET is presently used in soda bottles.

The bulk of common thermoplastics manufactured (PE, PP) are used in packaging products that have a relatively short useful lifetime that end up in the waste and litter streams rapidly. Plastics used in building construction (e.g. PVC) constitute about a third of the production but have much longer service lives. Figure 3.3 provides a summary of the major application areas for the commodity thermoplastics used in high volume.

Table 3.1 Frequency of occurrence of different polymer types in 42 studies of microplastic debris sampled at sea or in marine sediments (from Hidalgo-Ruz et al. 2012)

Polymer type	% studies (n)
Polyethylene (PE)	79 (33)
Polypropylene (PP)	64 (27)
Polystyrene (PS)	40 (17)
Polyamide (nylon) (PA)	17 (7)
Polyester (PES)	10 (4)
Acrylic (AC)	10 (4)
Polyoximethylene (POM)	10 (4)
Polyvinyl alcohol (PVA)	7 (3)
Polyvinyl chloride (PVC)	5 (2)
Poly methylacrylate (PMA)	5 (2)
Polyethylene terephthalate (PET)	2 (1)
Alkyd (AKD)	2 (1)
Polyurethane (PU)	2 (1)




Figure 3.2. Production of the most common artificial (plastic) and natural polymers, including some typical applications. Microplastics are manufactured for particular applications, such as industrial scrubbers or in personal cleaning products such as toothpaste. All plastics can be subject to fragmentation on environmental exposure and degradation into (secondary) microplastics. The proportion of plastic reaching the ocean to become plastic litter depends on the effectiveness of the re-use, recycle and waste management chain.

Table 3.2 Selection of reported polymer compositions in a variety of media (see Table 3.1 for abbreviations)

Matrix	Size	Polymer composition	Reference
sediment/ shoreline	<1 mm	PES (56%), AC (23%), PP (7%), PE (6%), PA (3%)	Browne et al. (2011)
sediment/ sewage disposal site	<1 mm	PES (78%), AC (22%)	Browne et al. (2011)
sediment/ beach	<1 mm	PES (35%), PVC (26%), PA (18%), AC, PP, PE, EPS	Browne et al. (2008)
Sediment/ Inter- and sub-tidal	0.03–0.5 mm	PE (48.4%), PP (34.1%), PP+PE (5.2%), PES (3.6%), PAN ^a (2.6%), PS (3.5%), AKD (1.4%), PVC (0.5%), PVA ^b (0.4%), PA (0.3%)	Vianello et al. (2013)
sediment/ beach	1–5 mm (pellet)	PE (54, 87, 90, 78%), PP (32, 13, 10, 22%)	Karapanagioti et al. (2011)
water/ coastal surface microlayer	<1 mm	AKD (75%), PSA ^c (20%), PP+PE (2%), PE, PET, EPS	Song et al. (2014)
water/ sewage effluent	<1 mm	PES (67%), AC (17%), PA (16%)	Browne et al. (2011)
fish	0.13–14.3 mm	PA (35.6%), PES (5.1%), PS (0.9%), LDPE (0.3%) AC (0.3%), rayon ^d (57.8%)	Lusher et al. (2013)
bird	-	PE (50.5%), PP (22.8%), PC and ABS ^e (3.4%), PS (0.6%), not-identified (22.8%)	Yamashita et al. (2011)

^a PAN: polyacrylonitrile, ^b PVA: polyvinyl alcohol, ^c PSA: poly(styrene:acrylate), ^d rayon – a semi-synthetic compound produced from pure cellulose, ^e ABS: acrylonitrile butadiene styrene




Figure 3.3. European plastics demand (EU27 + Norway & Switzerland) by resin type and industrial sector in 2012. Nylons (mainly Nylon 6 and Nylon 66) in fishing gear applications and polystyrene, polyurethane foams used in vessel insulation and floats, are employed extensively in the marine environment. Figure courtesy of PlasticsEurope (PEMRG)/Consultic/ECEBD.

Table 3.3 Densities and common applications of plastics found in the marine environment (adapted from Andrady 2011)

Resin type	Common applications	Specific gravity
Polyethylene	Plastic bags, storage containers	0.91–0.95
Polypropylene	Rope, bottle caps, gear, strapping	0.90–0.92
Polystyrene (expanded)	Cool boxes, floats, cups	0.01–1.05
Polystyrene	Utensils, containers	1.04–1.09
Polyvinyl chloride	Film, pipe, containers	1.16–1.30
Polyamide or Nylon	Fishing nets, rope	1.13–1.15
Poly(ethylene terephthalate)	Bottles, strapping	1.34–1.39
Polyester resin + glass fibre	Textiles, boats	>1.35
Cellulose Acetate	Cigarette filters	1.22–1.24

It would be mistaken to assume that the volume or production trend of particular polymer types or applications inevitably corresponds with the pattern of plastic litter observed. The variety of resin types produced is reflected in the composition of plastic debris recovered from the marine environment (Tables 3.1, 3.2), but there are many societal, economic, technical and environmental factors at play in determining the distribution and composition of plastic litter.

One key property that influences the behaviour of plastics in the marine environment is the density with respect to the density of seawater. Objects containing a void, such as a bottle, will tend to float initially but once objects lose their integrity it is the density of the plastic that will determine whether objects float and sink. The rate at which this occurs will influence the distance the object will be transported from its source. The densities of common plastic resins are shown in Table 3.3. In addition, the development of biofilms on the surface of the particle may alter the density sufficiently to cause it to float, even if the 'clean' polymer is less dense than seawater.

3.2 'Primary' and 'Secondary' microplastics

The distinction between primary and secondary microplastics is based on whether the particles were originally manufactured to be that size (primary) or whether they have resulted from the breakdown of larger items (secondary). It is a useful distinction because it can help to indicate potential sources and identify mitigation measures to reduce their input to the environment. Primary microplastics include industrial 'scrubbers' used to blast clean surfaces, plastic powders used in moulding, micro-beads in cosmetic formulation, and plastic nanoparticles used in a variety of industrial processes. In addition, spherical or cylindrical virgin resin pellets, typically around 5 mm in diameter, are widely used during plastics manufacture and transport of the basic resin 'feedstock' prior to production of plastic products. Secondary microplastics result from the fragmentation and weathering of larger plastic items. This can happen during the use phase of products such as textiles, paint and tyres, or once the items have been released into the environment. The rate of fragmentation is controlled by a number of factors (Section 3.2.2).

In addition to synthetic microplastics, naturally occurring biopolymers (Figure 3.2) may be present in the oceans. However, regardless of their particle size these are less of a concern because they are generally biodegradable and are less hydrophobic compared to synthetic plastics. Most natural polymers readily biodegrade into CO₂ and H₂O in the oceans (Poulicek et al. 1991). Biopolymers have been always present in the oceans unlike the microplastics whose origins are recent.

Although microplastics greatly outnumber large plastic items in marine systems, they still make up only a small proportion of the total mass of plastics in the ocean (Browne et al. 2010). This means that even if we were able to stop the discharge of macroplastic litter into the sea today, on-going degradation of the larger litter items already at sea and on beaches would likely result in a sustained increase in microplastics for many years to come.

3.2.1 Generation of microplastics

As plastic marine debris on beaches and floating in water is exposed to solar UV radiation it undergoes weathering degradation and gradually loses its mechanical integrity (Pegram and Andrady 1989). With extensive weathering plastics generally develop surface cracks (Cooper and Corcoran 2010) and fragment into progressively smaller particles (Qayyum and White 1993; Yakimets et al. 2004). This is the most likely process for the generation of secondary microplastics in the marine environment. Weathering degradation would occur particularly rapidly on beaches and at relatively low rates in floating debris. In the aphotic and low-oxygen environments in mid-water or sediment, the degradation and fragmentation are particularly slow. Knowledge of the fragmentation rates and mechanisms for common plastics in debris are needed to reliably infer the rates of microplastics generation, their particle size distribution (PSD) and their impact on

different biota. Such crucial information, especially the fragmentation mechanics, are not known reliably even for common plastics materials.

3.2.2 Weathering degradation of plastics in the ocean

The dominant cause of degradation of plastics outdoors is solar UV radiation, which facilitates oxidative degradation of polymers (Andrady et al. 1996). Photodegradation of common plastics such as polyethylenes, polypropylenes and polystyrene are free-radical mediated oxidation reactions (Celina 2013). The basic mechanism of this autocatalytic oxidation of common plastics is well established (Cruz-Pinto et al. 1994). During advanced stages of degradation, the plastic debris typically discolours, develops surface features, becoming weak and brittle (embrittle) in consequence over time. Any mechanical force (e.g. wind, wave, animal bite and human activity) can break the highly degraded, embrittled plastics into fragments.

Unlike virgin resin pellets, plastic products can incorporate a range of additives selected to modify the properties of the resin to meet the intended product application. These additives typically change the rate of oxidative degradation (and therefore weathering rates) of plastics. For example, UV and heat stabilizers and antioxidants used as additives often markedly retard light-induced degradation of the plastic material.

While the weathering of common plastics and their various formulations has been extensively studied in different environments, these studies have historically focused on the early stages of degradation that impact the useful lifetime of the product. Therefore, only very limited information is available on extensive oxidation and fragmentation of highly weathered plastics in the environment. Furthermore, there is virtually no information on weathering of plastics stranded on shorelines, floating in seawater (Andrady 2011; Muthukumar et al. 2011) and especially submerged in seawater or sediment. The effects of variables such as mechanical impact, salinity, temperature, hydrostatic pressure, presence of pollutants such as oil in seawater and bio-fouling (reducing UV exposure) on the rates of weathering according to various types of plastic items are virtually unknown.

Presently, there are no reliable methodologies to determine the age (or duration of outdoor exposure) of microplastics collected from the field, making it very difficult to investigate the degradation dynamics of microplastics in marine systems from collected field samples. While it is possible to quantify their extent of weathering by chemical analysis (FTIR⁹ or Raman spectroscopy) the duration of their exposure in the marine environment cannot be deduced from such information. The highly variable rates of weathering of large plastic items due to differences in polymer type, additive composition and environmental factors complicate the assessment of age. Without an accurate assessment of age, developing three-dimensional models of microplastic abundance is somewhat unconstrained. For example, a particular plastic can move great distances either because it is in a fast current or because it simply has been in the ocean for a long time.

⁹ Fourier transform infrared spectroscopy

3.2.3 Fragmentation of plastics

As plastic debris photo-degrades, especially on land, objects begin to show characteristic surface cracks and pits (Cooper and Corcoran 2010). Localization of cracks to the surface layer (of a couple of 100 microns) is a consequence of the rapid attenuation of the damaging solar UV radiation as it transmits in the bulk of the plastic. The microplastics enter the environment from this embrittled weak surface layer of oxidizing plastic debris. These reactions continue on in the microplastic particles generated (Gregory and Andrady 2003), possibly progressing to yield particles at the nano-scale. However, the existence of nano-scale plastic in the ocean has not been reported as yet.

Both weathering and fragmentation rates are relatively rapid on beaches but generally several orders of magnitude slower, decreasing in the following order; plastics floating in water, in the mid-water column or in the marine sediments. The degradation on beaches may be enhanced by the higher UV radiation, higher sample temperatures and mechanical abrasion attained by the beach litter, although some interaction between those factors is expected. The relative rates of degradation of plastic in different compartments of the marine environments have not been quantified but in any event depends on the plastic. However, the degradation of floating plastic is well known to be impeded by low water temperatures. Biofouling of floating plastics in the ocean is ubiquitous, and often leads to a rich growth of surface fauna. This shields the plastic from solar UV and invariably increase its density and hence sinking out of the photic zone, preventing further UV-mediated degradation (Gregory and Andrady 2003; Ye and Andrady 1991). In the aphotic (dark) and cold sediment environment no appreciable degradation is expected. In cases where the surface foulants are foraged or otherwise removed, the plastic may subsequently resurface into the floating debris. However, virtually no information is available on the fate of plastics in aphotic marine sediment.

Formation of microplastics in the ocean is influenced by a combination of environmental factors and the properties of the polymer

The general lack of research information on weathering and fragmentation of plastics in the marine environment is a very significant gap in relevant scientific knowledge. Lack of data on how the combined effects of photo-oxidation, fragmentation, mechanical abrasion and additive chemicals affect the formation of microplastics is a significant barrier to the production of reliable quantitative models to describe the behaviour of plastics and microplastics in the ocean.

3.2.4 Biodegradation and Mineralization

An understanding of terminology is important when describing the fate of plastics in the ocean. Complete degradation refers to the destruction of the polymer chain and its complete conversion into small molecules such as carbon dioxide or methane (also called mineralization). This is distinct from degradation which refers to an alteration in the plastic's properties (e.g. embrittlement, discolouring; Section 3.2.2) or its chemistry.

Few plastics undergo complete degradation or mineralization in the marine environment. Plastics such as aliphatic polyesters, bacterial biopolymers and some bio-derived polymers are readily biodegradable in the environment. But often, these are more expensive to produce than commodity plastics. Ideally, biodegradability is desirable only after the useful service life when the product is in litter or marine debris. But, for most applications it is the durability of plastics that is the most sought after property; it is not clear if the existing biodegradable plastics deliver the requisite mechanical integrity and durability needed for most applications during their useful life.

3.3 Sampling methods for microplastics in the marine environment

3.3.1 Introduction

Sampling and analysis form our link to gaining knowledge about the natural world and are necessary for studying and assessing the environmental impacts of microplastics. Choices made in the design and selection of sampling and analytical methods determine what types of microplastics are sampled and what types of microplastics are detected; methods have limitations in particle size ranges and types of plastic materials targeted for measurement. Methods all have a given degree of specificity in what is targeted which depends on how the microplastics are extracted from the environmental matrix, such as seawater, sediment and biota.

Sampling and analysis is the first step in recording the presence or absence of microplastics in the ocean. Astute researchers in the early 1970s first realized the existence of microplastics in the open ocean (Carpenter et al. 1972) and there followed a steady trickle of confirmatory records until an explosion of interest and publication from about 2005 onwards (Figure 1.1). The ecological risk posed by microplastics (see Section 4) is calculated by combining the potential hazard with the probability of encountering that hazard. In order to assess probability it is important to gain an adequate understanding of the distribution of microplastics in the various environmental compartments.

Whether sampling at the sea surface, on the seabed or in the intertidal it is important to recognize that a variety of methods are available (Nuelle et al. 2014; Hidalgo Ruz et al. 2012). Currently there are no universally accepted methods for any of these matrices and the methods available all have potential biases. For example, the mesh size of the net or sieve used to extract microplastics from the bulk medium will influence the shape and size of particle captured. Separation of microplastics from a bulk medium becomes increasingly more challenging as the particle size decreases. For particles less than 100 μm visual separation is typically followed by forensic analysis, for example by spectroscopy. However, whether or not a particle is selected for subsequent forensic determination will be influenced by the degree to which it stands out against the background of other particulates in the sample. For example brightly coloured plastic fibres are conspicu-

ous compared to more spherical natural sediment and are likely to be more completely sampled than plastic particles which have similar size and shape to that of natural sediment. Plastic particles of this latter type are hence likely to be under-sampled (Woodall et al. 2014). In addition the distribution of microplastics within a particular matrix is variable in time and space; for example particles can become redistributed vertically in the water column as a consequence of surface turbulence (Kukulka et al. 2012). Similarly particles can become buried or exposed on sandy beaches according to the prevailing sediment depositional regime (Turra et al. 2014).

Methods development for sampling and analysis of microplastics is an important, emerging area of research and development in marine litter science (e.g. Galgani et al. 2011). Many decisions are made in the process of designing and implementing sampling plans that can affect the end result of a study, and reliability and representativeness of the results. Ancillary parameters such as the sample volume, collection technique and sea state (Kukulka et al. 2012) are important to consider. This section provides a brief overview of sampling approaches.

3.3.2 Sampling seawater

Sampling for microplastics in the water column requires decisions about the size ranges to be targeted for sampling and analysis, then selecting the appropriate equipment. Concentrations of microplastics are usually too low to allow sampling by standard water samples, used for routine chemical analysis (e.g. 1–100 l). Some form of filtration is usually required to allow much larger volumes of water (many m³) to be analysed. Some studies have employed bulk water sampling followed by vacuum filtration to capture microplastic particles (Ng and Obbard 2006; Desforges et al. 2014). More commonly, towed nets are utilized to filter large volumes of water in situ. A variety of surface-towing nets have been employed over time, designed primarily for sampling biological specimens (e.g. manta net, neuston net, ring net), with variable net mesh, net aperture and net length dimensions (Figure 3.4).

Towing protocols also vary in terms of the depth of water sampled (ranging from microns to tens of cm) and length of tow. Below the sea surface, obliquely towed nets have been employed to sample from the sea surface to a particular depth (e.g. Doyle et al. 2011), while multiple opening–closing nets sample discrete depth layers (Kukulka et al. 2012). Continuous plankton recorders (CPR)¹⁰ collect plankton at a depth of 10 m and CPR samples have been analysed for microplastics (Thompson et al. 2004). Archived samples, both from towed nets and CPR tows, can provide a valuable resource for microplastic research.

When nets, sieves or filtration systems are used, the mesh size selected determines the minimum size that is targeted for sampling, although smaller particles will be captured (Goldstein et al. 2013). The majority of nets used currently have a nominal mesh size of 333 μm , with the CPR membrane being 285 μm . A number of factors such as net clogging influence the actual

volume of water filtered and the sampling efficiency. This problem tends to increase in inverse proportion to the mesh size, and finer mesh nets are more liable to damage.



Figure 3.4 Manta net being used to collect microplastics from surface waters in the Pacific, operated from S/V Kaisei, 2009. Image courtesy of Doug Woodring, Project Kaisei; <http://projectkaisei.wordpress.com/the-science-team/>.

Results may be reported in terms of the abundance of microplastics (number km⁻² or number m⁻³ seawater) or the mass of microplastics (total mass km⁻² or mass m⁻³ seawater). Both metrics can be useful. However, surface nets can only be used to quantify abundance in a two dimensional area of sea surface; this is because surface nets are only partially submerged and can move vertically relative to the sea surface and so, unlike sub-surface nets, it is not possible to obtain reliable estimates of volume since the amount of water passing through the net will vary according to its vertical position in relation to the sea surface. Macroplastics, such as fishing debris, are far fewer in number but have a much greater mass, whereas microplastics are far more numerous but have a correspondingly lower mass. It is helpful if both mass and abundance are reported (Browne et al. 2010; Morét-Ferguson et al. 2012; Eriksen et al. 2013b). Mass is useful from an overall waste management perspective whereas number is likely to be of greater ecological significance. The mass of plastics is reported on a dry weight basis, as is the mass of natural inorganic particles. If comparisons are made with the abundance of living biota, such as zooplankton, it is essential to report the latter on the basis of wet weight, to avoid misleading interpretations on the relative quantities in the environment.

Improved techniques for bulk sampling and quantification of sub-millimetre sized particles in seawater are under development. Sampling of nanoparticles using towed nets is impractical (it would have to be a nano-membrane technology). The volumes required for detection of particles in the nano-size range is unknown, but likely to be in the same order as the volume needed for sampling microplastics that are >333 μm . Man-made nanoparticles in the nanotechnology size range (approximately 10–100 nm) are expected to be present in the environment as aggregates due to surface interactions (Quik et al. 2012),

¹⁰ <http://www.sahfos.ac.uk>

and may not be homogeneously distributed throughout the water column. Man-made nanoparticles are also likely to aggregate with any naturally occurring particles including those in the nano-size range. Particles in the 100 nm to 1 µm size range may not aggregate as strongly because surface interactions diminish with increasing particle size.

3.3.3 Sampling sediments

Sampling shoreline sediments is generally more straightforward than sampling seawater. However, sampling seabed sediments can require significantly more effort and resources. Sampling shoreline sediments for microplastics will vary according to the purpose of the study. Aspects to consider include the sampling depth (surface or vertical profile), the sample size, and the location in relation to the strandline (an accumulation zone). Sediment samples may undergo some form of separation in the field, such as a hand-held sieve (Figure 3.5), before further processing in the laboratory. Sampling finer-grained sediments will usually require more elaborate, laboratory-based separation techniques. Protocols are being developed for sampling shoreline sediments for marine debris, including some with particular regard to microplastics (e.g. Galgani et al. 2011).

Significant heterogeneity has been observed in the distribution of microplastics on beaches, both from surface transects and vertical profiles, emphasizing the need to include these considerations when designing an effective sampling strategy (Browne et al. 2010; Turra et al. 2014).



Figure 3.5 Collecting microplastics from surface sediments by dry sieving, prior to carrying out compositional analysis; Busan, Republic of Korea.

3.3.4 Sampling biota

Microplastics have been observed in several species of fish, bivalves, crustaceans and birds, with greatest focus on stomach content analysis (see Section 4). The stomach content of birds may contain plastic objects covering a wide size spectrum, whereas tissues tend to contain much smaller size ranges because smaller particles are more likely to be translocated from the

GI tract (see Section 4). Sampling of birds relies on the recovery of dead specimens, usually from shorelines or coastal nesting sites, for example from mid-ocean islands. Objects observed may be several cm in diameter (e.g. cigarette lighter) but much smaller objects have also been found (Figure 3.6). The longest running monitoring programme operates in the North Sea, based on the analysis of recovered carcasses of the northern fulmar (*Fulmarus glacialis*) (van Franeker et al. 2011). This has been incorporated into a wider programme for environmental management, providing the basis for establishing an Ecological Quality Objective, based on the average mass of plastic in the stomach.




Figure 3.6 Fragments of plastic, including microplastics, in the stomach contents of a northern fulmar (*Fulmarus glacialis*) from the North Sea. Image courtesy of OSPAR/KIMO, photograph Jan van Franeker.

3.4 Determining the composition of microplastics

The analysis of environmental samples is a multi-step process that may include: sample preparation (such as sample homogenization or pre-concentration steps); extraction of microplastics; further purification ('clean-up') of the microplastic extract to remove additional non-microplastic matrix; and, detection and quantification of particles and identification of polymer types.

The specific composition of the microplastics in the sea is potentially very broad, reflecting the innovations in materials science over the past century that have brought a multitude of different polymers, copolymers and blends, combined with a variety of fillers and additives, giving each new material and each new product its characteristic physical-chemical properties. These innovations and the diversity of plastic material composition that results, has created challenges especially for laboratory technicians carrying out analytical work for targeted, quantitative analyses of microplastics in environmental matrices.

Besides targeting the plastic materials, the other major challenge is analysing the broad range of particle sizes of microplastics, which is important for understanding microplastic distribution, dispersal and dynamics but also the size-dependent biological effects (Section 4). Our ability to identify polymers of microplastics varies with particle size. Many studies to date have focused

on large microplastics (1–5 mm), including pre-production resin pellets, which are visible to the naked eye and can be picked out of nets, beach sand or biota samples with tweezers. However, when smaller particles are targeted for analysis, they are harder to identify with the techniques presented below.

Separation

Microplastics need to be separated from other organic and inorganic particles materials in the sample prior to being counted, weighed and the polymer type identified. Initial separation can be achieved by density separation using solutions of varying density (e.g. ZnCl_2 , NaI and sodium polytungstate). Co-extracted small organic particles, which may interfere with microscopic as well as spectroscopic identification, can be removed using hydrogen peroxide and sulphuric acid. It becomes increasingly difficult to distinguish plastic from non-plastic particles with decreasing size, using microscopic examination alone (Figure 3.7). Spectroscopy can confirm the presence of plastics and provide the polymer composition. In addition there have been recent advances in separation of microplastic from natural organic material. Such separation has been achieved in surface water samples using an enzymatic digest to remove planktonic organisms (Cole et al. 2014).




Figure 3.7 Microscope image of a plastic pellet (left) and squid eye (right), both are approximately 1 mm in diameter.

Raman and FTIR spectroscopy

Spectroscopy is required to confirm the identification of plastics, and their synthetic polymer for particles <1 mm in size. Microscopic FTIR (Fourier transformed infra-red) and Raman spectroscopy are the most promising methodologies in this regard. Current methods of analysis of field-collected microplastics make it possible to identify polymer types in particles as small as approximately 10 μm . These particles are visible under the light microscope and are still large enough to be identified as plastic using FTIR or Raman microscopy. These techniques rely on light transmission and wavelengths of light, and if the particle is smaller than these wavelengths, no reliable polymer IR spectrum can be achieved. This results in a major challenge to the identification of plastic particles in the 20 nm to 10 μm (10,000 nm) size range.

Fluorescently labelled nano- and micro-sized plastic particles can be utilized in laboratory experiments examining microplastic behaviour, allowing measurement through fluorescence detection. This approach is suitable to laboratory studies but does not solve the problem of measuring unknown and unlabelled plastic particles in environmental matrices. Development of new analytical techniques is required to extract, isolate and identify micro- and nano-sized plastic particles in the marine environment but to date no validated methods yet exist for field samples.

Method validation

The pioneering microplastic field surveys to date have taken place in the research and development (R&D) sphere and have been largely semi-quantitative. When a new chemical contaminant begins to be measured in environmental matrices, much effort is required to evaluate and improve existing methods and develop new products and initiatives, such as reference materials, proficiency testing schemes, ring tests, inter-calibration exercises and standard operating protocols (SOPs). This helps to ensure that the quality of the data produced meets predefined performance criteria, which may lead to some form of accreditation. The field of microplastics research is at a less mature stage by comparison. In addition, the extremely varied nature of microplastics in the field, in terms of composition and distribution, makes it unlikely that such stringent procedures are either possible to achieve in a meaningful way or strictly necessary. The resulting data can still be of use for determining the relative state of the environment, and informing decisions on possible management measure.

3.5 Distribution of microplastics in the marine environment

3.5.1 Influence of the source

The distribution of microplastics in the ocean is influenced by the nature and location of the source of entry, as well as the subsequent complex interaction of physical, chemical and biological processes. There is growing information about all these aspects, but there remain major uncertainties about the spatial and temporal distribution of microplastics, and likely trends. Identification of the sources is important to gain an accurate assessment of the quantities of plastics and microplastics entering the ocean, to provide an indication of regional or local ‘hot spots’ of occurrence, and to determine the feasibility of introducing management measures to reduce these inputs.

Primary microplastics are manufactured particles designed for particular applications (Section 3.2.1). A proportion of these particles is released from discrete point sources such as factories and sewage discharges. In addition, there is evidence of virgin resin pellets being lost during transport at sea¹¹ or during transshipment. There is also evidence of point source inputs

¹¹ Loss of PP pellets from the container vessel Yong Xin Jie near Hong Kong during Typhoon Vicente, August 2012; <http://e-info.org.tw/node/79464>

near to plastic processing plants where the abundance of plastic pellets or powders can be considerable (e.g. Norén and Ekendahl 2009). However, there is a lack of quantitative data on diffuse inputs via small, but regular and persistent, losses from multiple sources such as loss of plastics pellets from processing plants or of micro-beads discharged through domestic wastewater systems. One study estimated that 200 tons of micro-beads were used in the USA annually, in personal care products, and that approximately 50% of these would pass through sewage treatment to the ocean (Gouin et al. 2011).

Estimating the distribution of microplastic based on secondary inputs is particularly difficult since it relies on accurate assessment of the distribution of macroplastics and the degradation process (which is also not well known). There is a lack of data comparing the abundance of macroplastics and microplastics at local scales. However, it is unlikely that the abundance of microplastic and macroplastics will be closely correlated as large and small objects will be influenced by environmental processes to differing degrees. For example, larger floating objects will be more prone to transport by winds than microplastics (Browne et al. 2010), and this is reflected in circulation models used to simulate the transport of micro- and macro-debris (Eriksen et al. 2014; Lebreton et al. 2012).

In some cases it is possible to link the presence of microplastics to particular industrial sectors. For example, data from shorelines in southern Korea indicate that the great majority of microplastic fragments are composed of EPS (Heo et al. 2013). The prevalence in this region is explained by the wide scale use of EPS in buoys for aquaculture installations. Similar high occurrences of EPS have been reported from Japan and Chile, also linked to coastal aquaculture. In most other regions the composition of microplastics will be much more varied and less dominated by a single component, presenting great difficulties in attributing occurrence to a particular source.

3.5.2 *Distribution of microplastics based on direct observations*

Understanding how microplastics are distributed both horizontally and vertically in the oceans is a prerequisite to assessing potential impacts. In short, there can be no direct impacts where there are no, or very few, microplastics. However, there is likely to be a greater potential for impacts if microplastics accumulate in specific locations. Before we can properly assess risks it is therefore essential to understand how microplastics are distributed in space, for example between broad geographic regions (temperate, tropical, polar); between open and relatively enclosed seas (e.g. the Mediterranean versus the Pacific Ocean); between compartments including sea surface, water column and benthic sediments; and between coastal habitats (e.g. salt marsh, mangrove, coral reef, mussel bed). It is also important to quantify the distribution across international boundaries as differences in production, consumption, usage and waste management practices have the potential to influence inputs to the ocean and are essential when considering measures aimed at reducing such inputs.

Surface ocean

The geographical coverage of microplastics sampling is growing each year. The bulk of microplastics surveys have so far been conducted in the northern hemisphere, and the majority of surveys are of the sea surface using plankton nets and on shorelines. Microplastics at the sea surface have been reported in the coastal ocean (Carpenter et al. 1972; Doyle et al. 2011; Reisser et al. 2013), the open ocean (Carpenter and Smith 1972; Law et al. 2010; Goldstein et al. 2012; Eriksen et al. 2013b), and in enclosed or semi-enclosed seas such as the Mediterranean Sea (Collignon et al. 2012; Law et al. 2014), North Sea (Dubai and Liebezeit 2013) and South China Sea (Zhou et al. 2011). Three studies have reported microplastics in the near-surface water column (typically upper tens of metres; Lattin et al. 2004; Doyle et al. 2011; Kukulka et al. 2012) but not in the deeper water column beyond ~200 m depth, possibly because there have been no dedicated sampling efforts in this regime. The North Atlantic and North Pacific Oceans and Mediterranean Sea are the best-sampled regions of the ocean for floating microplastics (Figure 3.8). Microplastics are widespread in the oceans across temperate and tropical waters, and have been reported near to population centres and also in considerable concentrations in more remote locations, as a result of long-distance transport in the surface ocean. While there is considerable spatial variability in reported abundances, studies have reported at least the presence of microplastics at all of the locations they examined. A key challenge is to scale up from regional seas and ocean systems in order to assess the quantities of plastic and microplastic contamination at a global scale. Such work is essential in order to better estimate the scale of the problem and to facilitate monitoring efforts that will be essential to evaluate the success of measures to reduce inputs of debris to the ocean. Considerable progress has been made recently in estimating quantities of surface plastic on a global scale (Cózar et al. 2014; Eriksen et al. 2014). There is also evidence of substantial accumulations of plastic and microplastics on the sea bed including areas of the deep sea down to 5766 m in the Kuril-Kamchatka Trench in the NW Pacific (Fischer et al. 2015), but estimating total quantities of plastic debris in the deep sea will be much more challenging than at the sea surface (Pham et al. 2014; Woodall et al. 2014).

Shorelines and seabed

The majority of studies on beaches have focused on larger items of debris. However, microplastics have also been reported on beaches at numerous locations worldwide (e.g. Gregory 1978; Wilber 1987; Browne et al. 2011; do Sul et al. 2009; Hidalgo-Ruz et al. 2012). Industrial resin pellets have frequently been reported as they are easily identifiable by eye and have been targeted for study of absorbed contaminants by the International Pellet Watch programme (Takada 2006). Most studies survey only the surface sediments, although some have collected samples below the surface (e.g. Claessens et al. 2011). Turra et al. (2014) found plastic pellets were distributed in beach sediments in Brazil to depths of at least 2 metres, with greater concentrations consistently being observed in sub-surface layers relative to the current beach surface.

There are several studies indicating that the abundance of microplastics in subtidal sediments can be greater than on shorelines (Thompson et al. 2004; Browne et al. 2011; Claessens et al. 2011). There have been very few reports of microplastics in deep sea sediments, but one study recorded their occurrence on the seabed at a water depth of 5000 m (Van Cauwenberghe et al. 2013c).

Spatial and temporal trends

Only a small number of data sets have sufficient sampling to begin to tease apart spatial and temporal trends in microplastic concentration. This is exacerbated by inherent space–time variability in environmental drivers, together with microplastic sources and transport between the various compartments (surface ocean, intertidal, seabed). Thompson et al. (2004) published the first assessment of microplastic abundance over time, and found that while the amount of microplastics measured between the British Isles and Iceland increased from the 1960s and 1970s to the 1980s and 1990s, no significant increase was observed between the later decades. Similarly, Law et al. (2010) found no significant increase in annual mean concentrations of floating microplastics in the western North

Atlantic subtropical gyre between 1986 and 2008, or in the eastern North Pacific subtropical gyre between 2001 and 2012 (Law et al. 2014). Goldstein et al. (2012) reported a significant two-order-of-magnitude increase in floating microplastic in the eastern North Pacific from 41 measurements collected from 1972 to 1987 and 1999 to 2010; however, data in the later time period were deliberately collected in the predicted region of highest plastic concentration, while earlier microplastic data were incidentally collected in more broadly dispersed samples. While it is possible, and even likely, that concentrations of floating microplastic in the ocean are increasing over time (based on increased plastic production, use and disposal since the 1950s), it is a challenging task to isolate such a trend given the large variability in the data and the size of the ocean that would need to be regularly and randomly sampled to properly account for this variability.

The concentration of microplastics recorded is directly influenced by the sampling approach used, which can vary significantly between studies. There is a need for greater harmonization to facilitate comparability (Galgani et al. 2011). Comparing concentrations between matrices is fraught with difficulty because the methods used are quite different, for example to sample sediment and bulk seawater.




Figure 3.8 (a) Distribution of microplastics in the western North Atlantic, 1986-2008. Sea Education Centre, Woods Hole, MA (downloaded from: http://onesharedocean.org/open_ocean/pollution/floating_plastics)




Figure 3.8 (b) Distribution of microplastics in the western North Pacific, 2001-2012. Sea Education Centre, Woods Hole, MA (downloaded from: http://onesharedocean.org/open_ocean/pollution/floating_plastics)

Table 3.4. Observations of microplastics in seawater

Location	Sampling method	Reported concentration	Reference
Seawater measurements			
North Sea - open water - sea surface	Continuous plankton recorder; 280 μm mesh silkscreen	Maximum concentration 0.04 – 0.05 fibres/ m^3	Thompson et al. 2004
Western North Atlantic - coastal - sea surface	Plankton net, 333 μm mesh	0.01 – 14.1 particles/ m^3	Carpenter et al. 1972
Western North Atlantic - open ocean - sea surface	Plankton net, 0.33 mm mesh	47 – 12,080 particles/ km^2	Carpenter and Smith 1972
Western North Atlantic - coastal and open ocean - sea surface	Plankton net, 0.947 mm mesh	60.6 – 5465.7 particles/ km^2 (mean values)	Colton et al. 1974
Western North Atlantic - open ocean - sea surface	Plankton net, 335 μm mesh	20,328 particles/ km^2 (mean near 30°N)	Law et al. 2010
Eastern North Atlantic - coastal - sea surface	Plankton nets, 180 μm and 280 μm mesh; continuous plankton recorder, 335 μm	0.01 – 0.32 cm^3/m^3	Frias et al. 2014
Mediterranean Sea - sea surface	Plankton net, 333 μm mesh	Mean concentrations: 0.116 particles/ m^2 , 0.202 mg/m^2	Collignon et al. 2012

Table 3.4. Observations of microplastics in seawater (cont.)

Location	Sampling method	Reported concentration	Reference
Mediterranean Sea - sea surface	Plankton net, 200 µm mesh	Mean concentrations: 0.94 particles/m ³ (Ligurian Sea) 0.13 particles/m ³ (Sardinian Sea)	Fossi et al. 2012c
North Pacific - open ocean - surface	Plankton net, 150 µm mesh	Maximum concentrations: 34,000 pieces/km ² , 3.5 mg/m ²	Wong et al. 1974
North Pacific - open ocean - sea surface	Plankton net, 333 µm mesh	Mean concentrations: 80 particles/km ² (Bering Sea) 3370 particles/km ² (subarctic) 96,100 particles/km ² (subtropical)	Day and Shaw 1987
Eastern North Pacific - open ocean - sea surface	Plankton net, 330 µm mesh	31,982 – 969,777 pieces/km ² 64 – 30,169 g/km ²	Moore et al. 2001
Eastern North Pacific - open ocean - sea surface	Plankton net, 335 µm mesh	In accumulation zone: 156,800 pieces/km ² (mean) 33,909 pieces/km ² (median)	Law et al. 2014
Western North Pacific - Kuroshio Current - sea surface	Plankton net, 330 µm mesh	Mean concentrations: 174,000 pieces/km ² , 3600 g/km ²	Yamashita and Tanimura 2007
Australia - coastal and open ocean - sea surface	Plankton nets, 333 and 335 µm mesh	4256 pieces/km ² (mean) 1932 pieces/km ² (median)	Reisser et al. 2014

Table 3.5. Observations of microplastics in seawater and sea ice

Location	Sampling method	Reported concentration	Reference
Seawater measurements			
UK estuarine beach - surface to 3 cm deep	0.25 m ² quadrats, NaCl flotation	1 – 8 particles per 50 ml sediment	Browne et al. 2011
Belgian marine sediments - beach and subtidal	Van Veen grab for subtidal sampling; sediment cores; flotation in saline solution, 38 µm mesh sieve	Mean concentrations: 166.7 particles/kg dry sediment (harbours) 97.2 particles/kg dry sediment (sublittoral) 92.8 particles/kg dry sediment (beaches)	Claessens et al. 2011
Portugal beaches - surface to 2 cm deep	0.5 m ² and 2 m ² quadrats, flotation in NaCl solution, 1 µm filter	Mean concentrations: 185.1 items/m ² 36.4 g/m ²	Martins and Sobral 2011
Lagoon of Venice - surface to 5 cm deep	Box corer, flotation in NaCl solution, 32 µm mesh sieve	672 – 2175 particles/kg dry weight	Vianello et al. 2013
Equatorial western Atlantic island beaches - surface to 2 cm deep	900 cm ² quadrats, 1 mm mesh sieve	4.6 x 10 ⁻³ g marine debris per gram of sand (mean)	Ivar do Sul et al. 2009
Brazil beaches - surface to 2 m deep	1 m ² trenches, sampling plastic pellets at 0.1 m intervals from 0.2 to 2 m depth, flotation in seawater, 1 mm mesh sieve	0.10 – 163.31 pellets/m ³	Turra et al. 2014
Singapore mangroves - surface to 4 cm deep	1.5 m ² quadrats, flotation in saline solution, 1.6 µm filter	3.0 – 15.7 particles per 250 g dry sediment	Nor and Obbard 2014
Kuril-Kamchatka Trench (NW Pacific) - surface to 20 cm deep	Box corer (0.25 m ² sampling area), 300 – 1000 µm mesh sieves	60 – 2020 particles/m ²	Fischer et al. 2015
Sea Ice			
Arctic Ocean sea ice	Sea ice cores, melted and filtered with 0.22 µm filters	38 – 234 particles per m ³ of ice	Obbard et al. 2014

3.5.3 Transport pathways

Surface transport

The most extensive spatial pattern in sea surface microplastics is their accumulation in large-scale subtropical ocean gyres, where convergent ocean surface currents concentrate and retain debris over long time periods (e.g. Wilber, 1987; Law et al. 2010; Goldstein et al. 2012; Law et al. 2014), and in enclosed seas such as the Mediterranean (Collignon et al. 2012) where surface water is retained for long periods of time because of limited exchange with the North Atlantic. These retention mechanisms are further supported by results of numerical models that predict where floating debris will accumulate in the surface ocean based on ocean physics (Section 3.4.4). While the highest concentrations of floating debris have been observed in subtropical gyres and in the Mediterranean Sea, there is significant small-scale variability within these regions, with order of magnitude variations in concentration observed on scales of 10s of km (Law et al. 2014).

Floating microplastic acts as a passive tracer and is therefore transported in the direction of net flow, which results from a combination of large-scale (1000s km) wind-driven currents as in the subtropical gyre down to centimetre-scale turbulent motion, and including all scales in between from eddies and surface and internal waves, for example. In addition, while the gyres are relatively steady circulation features in time, smaller scale water movements are highly time-dependent because they are locally driven in response to often quickly changing wind and sea conditions. Thus, while it is easy to predict that floating microplastic will be encountered near the centre of subtropical gyres at any given time, one cannot predict the concentration of debris at a particular time and location within the gyre. Similarly, one is unlikely to encounter large amounts of floating microplastic in boundary currents at the edges of gyres, or at equatorial or sub-polar latitudes; however, microplastic might still be found in these places at any particular time, especially if located near a debris source.

There are additional considerations, for example estimates of the rates of transfer between reservoirs in marine habitats, organisms or compartments. For example if, in what is probably an unrealistically simplistic scenario, the ultimate fate for most microplastics was the sea bed then understanding the amount of time spent at the sea surface or in the water column, while en route to the sea bed, would be critical to our understanding of rates of accumulation, relative abundance in each of these compartments and how long it might take for any changes in the quantity of microplastics entering the ocean to be apparent. Therefore, in undertaking these assessments it is important to know not only the abundance of microplastics, but also the size distribution, shape and polymer type; all of these have the potential to influence the type and magnitude of any associated impacts.

Vertical transport in seawater

It is important to note that oceanic currents are reasonably well known, but vary with depth. To further complicate matters, if particles change buoyancy, their pathways will depend on time in a specific vertical layer in the ocean. Because of their likely slow sink rates microplastic dispersal might be influenced by varying circulations at different depths. A study by McDonnell and Buesseler (2010) found sinking rates of marine particles to vary between 10 to 150 metres per day. In the deep ocean this means particles would take about one month to a year to reach the sea floor from the surface. Typical horizontal currents are on the order 1 metre per second, but decay quite rapidly to just a few centimetres a second at 1000 metres. Thus, a sinking particle may transit from 1 kilometre (fast sinking rate) to 35 kilometres (slow sinking rate) horizontally from its point of origin. This assumes the particle is continuously sinking, however, it is also possible particles spend a long time in the surface layer before sinking, in which case the horizontal displacements would be much greater.

The role of biofouling on density changes in microplastics is poorly known. Some larger items of microplastics such as pre-production pellets support colonization by macrobiotic fouling organisms (Gregory 2009) and items are likely to develop a microscopic biofilm (Lobelle and Cunliffe 2011; Zettler et al. 2013). Experiments indicate that biofouling of large plastic items can cause sinking through increased density (Ye and Andrady 1991).

Retention in biota and sediment

Microplastics can be taken up and retained for varying periods by marine organisms, which can potentially transport them significant distances. In the case of seabirds and seals, microplastics can even be carried back onto land. Microplastics may be trapped in sediments for long periods, although wave action and beach erosion can release particles from at least shallow water sediments. Microplastics might be more readily re-suspended from bottom sediments than larger plastic items simply because they are small and of low density compared to natural sediment and hence more easily disturbed by wave action, currents or bioturbation. Hence it is important to consider temporary and permanent sinks of microplastic.

3.5.4 Modelling the transport and distribution of microplastics

Given the challenges and limitations in assessing microplastic distributions by direct observations, various techniques have been employed to *estimate* where microplastics might be found and in what numbers. These include both theoretical and numerical modelling. Both approaches involve deriving an estimate based on known factors that include sources, sinks and forcing. Ideally two of these are known and the model is used to estimate the third. In the specific case of marine microplastic either the source or sink may be known, and the transport mechanism is presumed to be ocean currents. Thus the model would utilize known input/output and transport to estimate output/input. The transport mechanisms include ocean (and in the case of aerosols, atmospheric) currents and translocation by marine organisms. The former can be estimated with reasonable accuracy using numerical ocean modelling; the latter is much more challenging. An example of these approaches comes from modelling of the track taken by macroscopic items of tsunami debris across the Pacific Ocean from Japan to the western seaboard of the USA (Lebreton et al. 2012). There are several examples of the application of numerical models to study a subset of these, including tracking individual large items (debris, boats, etc.), subsurface items (Wilcox et al. 2013), dispersal of smaller particles (e.g. oil spills, planktonic larvae), formation and distribution of water masses, etc.

Using numerical models to track marine debris has thus far focused mainly on surface floating debris (Lebreton et al. 2012; Maximenko et al. 2012; van Sebille et al. 2012). Due to the nature of this particular problem, models start with an initial condition, for example a point source of known amount. Relevant forcing is then primarily wind and wind-driven near-surface currents. The model integration then gives an estimate of the pathways and landing points for the particles.

Extending this to examining pathways of marine microplastic is not straightforward for many reasons. Firstly, the initial conditions (sources) are not well known. Lebreton et al. (2012) experimented using different source points for debris, including the area of urbanized watershed, coastal population density, and shipping density, and their study also included estimates of landing locations (deposition on shore). Maximenko et al. (2012), in contrast, assumed a continuous distribution of particles. In both studies the particles aggregated in discrete regions of the open ocean, but in the latter case the simulation failed to reproduce the observed higher concentrations in the Mediterranean. The output from the Lebreton model was used to estimate the relative abundance of microplastics in Large Marine Ecosystems (Figure 3.9), as a contribution to the GEF Transboundary Waters Assessment Programme (<http://geftwap.org>). The approach cannot provide an accurate estimate of actual abundances, but can provide useful insights to help inform management decisions on future funding decisions.




Figure 3.9. Estimated relative distribution of microplastic abundance in Large Marine Ecosystems, based on Lebreton et al. 2012. Screen shot from www.oneshared.ocean.org

Secondly, particles may change density over time adding an additional (vertical) dimension. Most studies have assumed a constant density, assuming that surface currents are the sole drivers. Thirdly, the ultimate fate of these particles is not well known, but it likely includes vastly different geophysical regimes, including open-ocean, benthic, near-shore and on-shore. Finally, these particles become integrated into marine ecosystems by biological uptakes and discharges (redistribution). Thus, particle pathways cannot be treated as “passive” tracers in ocean currents. Nonetheless, numerical modelling may provide important insights into estimating pathways and thus locations of marine microplastic. One way to do this would be an extension of modelling work used to study floating debris, and to add a vertical component based on known rates of density changes.

Regional modelling studies have attempted to identify local accumulation areas at-sea (Pichel et al. 2007; Martinez et al. 2009; Eriksen et al. 2013b) and on-shore (Kako et al. 2007). Pichel et al. (2007) investigated the accumulation of debris in the North Pacific Subtropical Convergence Zone (STCZ) and were able to relate various forcing (e.g. winds) and indicators of convergence (e.g. sea surface temperature gradients) to derive a Debris Estimated Likelihood Index (DELI).

In summary, numerical models represent a tool for estimating pathways, sources and sinks of marine microplastics. In order to be effective however, the models need to have accurate descriptions of the initial conditions (sources for forward trajectories or sinks for backward trajectories) and changes in particle density. Finally, it should be pointed out that the particular issue of microplastic pathways in the ocean, and the application of numerical models of circulation, is in some sense a matter of statistics. Ocean models could therefore be used to estimate or predict areas

of accumulation in a broad sense (percentages, likelihood, etc.) but it would be extremely difficult to apply these techniques to individual particles.

3.6 Recommendations for further research

- Generate data on weathering-induced fragmentation of at least the PE, PP and EPS plastics in the marine environment.
- Examine the influence of weathering on particle sorption characteristics.
- Establish improved and validated methods for sampling at the sea surface in sediments and in biota.
- Organize inter-calibration exercises and harmonize reporting units to make future data comparable around the world.
- Design sampling strategies to establish time trends and spatial trends in selected marine areas.
- Conduct additional sampling of sub-tidal and in particular deep sea sediment.
- Investigate nano-sized plastic particles in marine organisms as a critical input for future risk assessments.
- Develop more realistic transport models, to incorporate variable particle properties, 3D circulation and sources of plastics and microplastics.

4 EFFECTS OF MICROPLASTICS ON MARINE BIOTA

4.1 Introduction

The potential ecological and human health risks of microplastics are relatively new areas of research, and there is currently a large degree of uncertainty surrounding this issue.

Risk is a function of hazard and exposure (dose), and evaluating the risks from microplastics requires knowledge of hazard (i.e. the potential of microplastics to cause adverse effects through plausible mechanisms), exposure levels (i.e. the quantities of microplastics detected in the environment, including in living organisms) and their effects (identification of dose-response relationships and threshold levels). The risk assessment of microplastics in the marine environment is still in the hazard characterization phase due to limited information on exposure levels and established effect levels. Rational policy measures are difficult to develop given the current incomplete and uncertain risk analysis, and it is important that priority should be given to systematically improving assessment of the risk of microplastics in the world's oceans.

The definitions of hazard, exposure and risk used in this document follow the International Union of Pure and Applied Chemistry (<http://www.iupac.org>). Hazard is a set of inherent properties of a substance, mixture of substances or a process involving substances that, under production, usage or disposal conditions, make it capable of causing adverse effects to organisms or the environment, depending on the degree of exposure; in other words, it is a source of danger. Consider the hazards posed by plastics in the environment that differ based on size of the plastic debris and size of the organism. Exposure is the concentration, amount or intensity of a particular physical or chemical agent or environmental agent that reaches the target population, organism, organ, tissue or cell, usually expressed in numerical terms of concentration, duration and frequency (for chemical agents and microorganisms) or intensity (for physical agents). Risk expresses either the probability of adverse effects caused under specified circumstances by an agent in an organism, a population or an ecological system; or the expected frequency of occurrence of a harmful event arising from such an exposure.

Particles have their effect as a consequence of several potential factors, relating either to physical or chemical effects. For physical effects particle size and shape will be important. For chemical effects two key factors act together to determine their potential to cause harm: their large surface area and reactivity, and the intrinsic toxicity of the polymer and absorbed contaminants. Smaller particles have more surface area per unit mass and therefore will likely exhibit more intrinsic toxicity. From this perspective, smaller microplastic particles less than 100 micrometres may be considered to be more likely to cause chemical effects in marine organisms, but this hypothesis has not been robustly tested. Shapes of microplastic range from fibres to spheres with varying surface roughness and sizes include fine particles (~200 nm) down to ultra-fine particles (<200 nm) and it is likely that far smaller

sizes in the size range of nanoparticles occur in the environment as well. It is important to note that the toxicities of engineered nanoparticles (ENP) are themselves diverse, and the toxicity of a given ENP may not be directly extrapolated to secondary nanoplastics (Andrady 2011).

4.2 Exposure

4.2.1 *Exposure through the gills*

External exposure results when microplastics contact the outer surfaces of the organism, including gills. Subsequently they may be translocated from the outside into the organism. The magnitude of external exposure depends on the concentration and size distribution of the microplastic particles and upon the specific nature of the organism. For most or all organisms that actively feed, external exposure of microplastics is small relative to exposure through ingestion (see below). The possible exception to this is very small (less than 40 μm) particles which may pass across gills. In non-filter feeding marine organisms such as the shore crab (*Carcinus maenas*), Watts et al. (2014) tested uptake of fluorescently labelled polystyrene microspheres (8–10 μm) through inspiration across the gills as well as ingestion of pre-exposed food. Ingested microspheres were retained within the body tissues of the crabs for up to 14 days following ingestion and up to 21 days following inspiration across the gill, with uptake significantly higher into the posterior versus anterior gills. These results identify ventilation as a possible route of uptake of microplastics into a common marine non-filter feeding species. Results were used to construct a simple conceptual model of particle flow for the gills and the gut.

4.2.2 *Ingestion*

Field studies have demonstrated that microplastics are ingested by a large variety of marine taxa representing various trophic levels, including fish-eating birds, marine mammals, fish and invertebrates, e.g. lugworms, amphipods and barnacles, mussels, sea cucumbers, zooplankton. The occurrence of plastic particle ingestion is reported from all oceanic regions (Table 4.1 (a–g)) in numerous species, for example in pelagic planktivorous fish from the North Pacific Central Gyre (Boerger et al. 2010), pelagic piscivorous fish from the North Pacific Ocean (Jantz et al. 2013) and pelagic and benthic (bottom dwelling) fish from the English channel (Lusher et al. 2013) and the North Sea (Foekema et al. 2013), in nephrops in Clyde sea (Murray and Cowie 2011), marine mussels from Belgian breakwaters (Van Cauwenberghe et al. 2012a), stranded whales (de Stephanis 2013), harbour seals from the North Sea (Rebolledo 2013), Franciscana dolphins from the coast of Argentina (Denuncio et al. 2011), Northern Fulmars from the North Sea (van Franeker et al. 2011), wedge-tailed shearwaters from the Great Barrier Reef, Australia (Verlis et al. 2013) and Magellanic penguins from the Brazilian coast (Brandao et al. 2011).

Invertebrates that ingest microplastics include deposit feeding lugworms *Arenicola marina* (Thompson et al., 2004) and sea cucumbers (Graham and Thompson, 2009), filter feeding salps *Thetys vagina* (sponges, polychaetes, echinoderms, bryozoans, bivalves, barnacles *Semibalanus balanoides* (Thompson et al. 2004; Ward and Shumway, 2004; Van Cauwenberghe et al. 2013a, and detritivores, such as amphipods *Orchestia gammarellus* (Thompson et al. 2004). There is also evidence of planktonic organisms other than salps consuming microplastics, namely arrow worms and larval fish (Carpenter et al. 1972 cited in Fendall et al. 2009), copepods in laboratory feeding trials (Wilson 1973 cited in Fendall et al. 2009), invertebrate larvae such as trochophores (Bolton and Havenhand, 1998 cited in Fendall et al. 2009), the echinoderms echinoplutei, ophioplutei, bipinnaria and auricularia (Hart 1991 cited in Fendall et al. 2009) and freshwater zooplankton (Bern 1990). Results are from both laboratory and field studies.

A number of studies have focused on the exposure of marine microplastics under controlled conditions (Browne et al. 2013; Rochman et al. 2013a; Wright et al. 2013a, Cole et al. 2014; Chua et al. 2014). In addition, there is a body of literature on the particle size selection or preferences by many animals, using artificial fluorescent plastic beads or industrial primary plastic powder. A selection of the work is presented in Table 4.2. Two review articles that appeared in 2011 (Andrady 2011; Cole et al. 2011), and one in 2013 (Wright et al. 2013b) listed some of the laboratory studies that have been conducted.

Table 4.1 (a) Field observations of the occurrence of plastic particles in biota in the North Sea

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
North Sea						
Mytilus edulis, Blue mussel	Belgium	March 2013	Comparison of consumption mussels and wild type mussels	Microscopic fibres ranging from 200 – 1500 µm identified in mussel tissues	Not investigated	De Witte et al. 2014
Mytilus galloprovincialis, Mediterranean mussel	Belgium	March 2013	Comparison of consumption mussels and wild type mussels	Microscopic fibres ranging from 200 – 1500 µm identified in mussel tissues	Not investigated	De Witte et al. 2014
Mytilus edulis/galloprovincialis hybrid	Belgium	March 2013	Comparison of consumption mussels and wild type mussels	Microscopic fibres ranging from 200 – 1500 µm identified in mussel tissues	Not investigated	De Witte et al. 2014
Mytilus edulis, Blue mussel	Germany	2013	Quantification of microplastic level in bivalve species reared for human consumption	Average of 0.36 ± 0.07 particles per gram tissue (ww) identified	Not investigated	Van Cauwenberghe & Janssen 2014
Clupea harengus, Herring	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	8/566 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013
Gadus morhua, Cod	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	10/80 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013
Merlangius merlangus, Whiting	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	6/105 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013
Melanogrammus aeglefinus, Haddock	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	6/97 individuals contained plastic particles	Condition factor significantly lower in individuals that contained plastic than those without plastic. Data deemed insufficient to confirm the hypothesis	Foekema et al. 2013
Trachurus trachurus, Horse mackerel	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	1/100 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
Eutrigla gurnardus, Grey gurnard	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	1/171 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013
Scomber scombrus, Atlantic mackerel	North Sea	July – October 2010	Quantify the occurrence, number and size of plastic particles and test the hypothesis that ingested plastic adversely affects the condition of fish	1/84 individuals contained plastic particles	No significant difference in condition factor between individuals that contained plastic and those without	Foekema et al. 2013
Fulmarus glacialis, Northern fulmar	North Sea	2003–2007	Disseminate concept of the Fulmar-Plastic-EcoQO. Long term beached bird study.	95% of sampled birds had plastic in their stomachs. Critical level of 0.1 g of plastic exceeded in 58% of individuals and 60% of Dutch birds	Not investigated	van Franeker et al. 2011
Prionace glauca, Blue shark	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	4.7% contained plastic in gut contents	Not investigated	Pinnegar 2014
Gadus morhua, Cod	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	28.57% contained plastic in gut contents	Not investigated	Pinnegar 2014
Eutrigla gurnardus, Grey gurnard	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	14.29% contained plastic in gut contents	Not investigated	Pinnegar 2014
Scyllorhinus canicula, Lesser spotted dogfish	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	4.76% contained plastic in gut contents	Not investigated	Pinnegar 2014
Pollachius virens, Saithe	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	23.81% contained plastic in gut contents	Not investigated	Pinnegar 2014
Merlangius merlangus, Whiting	North Sea, Channel, Irish Sea, Spitzbergen	1955–2011	CEFAS fish stomach contents records	23.81% contained plastic in gut contents	Not investigated	Pinnegar 2014

Table 4.1 (b) Field observations of the occurrence of plastic particles in biota in the Mediterranean Sea

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
Mediterranean Sea						
Calonectris diomedea, Cory's shearwater	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	47/49 individuals contained plastic particles. Average of 14.6 ± 24.0 particles per bird and 22.4 ± 48.8 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Puffinus mauretanicus, Balearic shearwater	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	32/46 individuals contained plastic particles. Average of 2.5 ± 2.9 particles per bird and 3.8 ± 8.4 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Puffinus yelkouan, Yelkouan shearwater	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	22/71 individuals contained plastic particles. Average of 4.9 ± 7.3 particles per bird and 29.8 ± 86.6 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Ichthyophaga audouinii, Audouin's gull	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	2/15 individuals contained plastic particles. Average of 9.8 ± 35.7 particles per bird and 22.7 ± 67.6 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Larus michahellis, Yellow-legged gull	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	4/12 individuals contained plastic particles. Average of 0.9 ± 1.98 particles per bird and 1.4 ± 2.3 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Ichthyophaga melanocephalus, Mediterranean gull	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	1/4 individuals contained plastic particles. Average of 3.7 ± 7.5 particles per bird and 12.6 ± 25.3 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Rissa tridactyla, Blacked-legged kittiwake	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	2/4 individuals contained plastic particles. Average of 1.2 ± 1.9 particles per bird and 4.9 ± 6.7 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Morus bassanus, Gannet	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	1/8 individuals contained plastic particles. Average of 0.1 ± 0.3 particles per bird and 0.3 ± 0.8 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013
Catharacta skua, Great skua	Catalan Coast, Spain	2003–2010	Opportunistic study to quantify and characterize plastics in seabirds caught as bycatch by longliner fishing vessels	1/2 individuals contained plastic particles. Average of 0.5 ± 0.7 particles per bird and 0.4 ± 0.5 mg in weight	No significant correlation identified between mean body mass and the number or mass of ingested plastic	Codina-Garcia et al. 2013

Table 4.1 (c) Field observations of the occurrence of plastic particles in biota in the North Atlantic Ocean

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
North Atlantic Ocean						
Nephrops norvegicus, Norway lobster	Clyde Sea, UK	May – June 2009	To determine whether N. norvegicus ingests small plastic fragments in the Clyde Sea	100/120 animals contained plastic in their stomach. Predominantly monofilament	Not investigated	Murray & Cowie 2011
Crassostrea gigas, Pacific oyster	Brittany, France	2013	Quantification of microplastic level in bivalve species reared for human consumption	Average of 0.47 ± 0.16 particles per gram tissue (ww) identified	Not investigated	Van Cauwenberghe & Janssen 2014
Myoxocephalus aenus, Grubby	Long Island Sound, USA	February – May 1972	Opportunistic sampling of fish caught for analysis of power plant impacts	White, opaque polystyrene spherules ingested and found in the gut	Not investigated	Carpenter et al. 1972
Pseudopleuronectes americanus, Winter flounder	Long Island Sound, USA	February – May 1972	Opportunistic sampling of fish caught for analysis of power plant impacts	White, opaque polystyrene spherules ingested and found in the gut	Not investigated	Carpenter et al. 1972
Roccus americanus, White perch	Long Island Sound, USA	February – May 1972	Opportunistic sampling of fish caught for analysis of power plant impacts	White, opaque polystyrene spherules ingested and found in the gut	Not investigated	Carpenter et al. 1972
Menidia menidia, Silversides	Long Island Sound, USA	February – May 1972	Opportunistic sampling of fish caught for analysis of power plant impacts	White, opaque polystyrene spherules ingested and found in the gut	Not investigated	Carpenter et al. 1972
Merlangius merlangus, Whiting	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	16/50 contained plastic particles	Not investigated	Lusher at al. 2013
Micromesistius poutassou, Blue whiting	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	14/27 contained plastic particles	Not investigated	Lusher at al. 2013
Trachurus trachurus, Atlantic horse mackerel	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	16/56 contained plastic particles	Not investigated	Lusher at al. 2013
Trisopterus minutus, Poor cod	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	20/50 contained plastic particles	Not investigated	Lusher at al. 2013
Zeus faber, John dory	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	20/42 contained plastic particles	Not investigated	Lusher at al. 2013
Aspitrigla cuculus, Red gurnard	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	34/66 contained plastic particles	Not investigated	Lusher at al. 2013

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
Callionymus lyra, Dragonet	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	19/50 contained plastic particles	Not investigated	Lusher at al. 2013
Cepola macrophthalma, Red band fish	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	20/62 contained plastic particles	Not investigated	Lusher at al. 2013
Buglossisium luteum, Solenette	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	13/50 contained plastic particles	Not investigated	Lusher at al. 2013
Microchirus variegates, Thickback sole	Plymouth, UK	June 2010 – July 2011	Part of routine long term trawl sampling. 70–75 mm cod end mesh size	12/51 contained plastic particles	Not investigated	Lusher at al. 2013
Uria aalge, Common murre	Newfoundland & Labrador, northwest Atlantic	1980s, 1990s, 2010s	Evaluate change over time in the ingestion of plastic by murres	4/374 contained plastic particles	Not investigated	Bond et al. 2013
Uria lomvia, Thick-billed murre	Newfoundland & Labrador, northwest Atlantic	1980s, 1990s, 2010s	Evaluate change over time in the ingestion of plastic by murres	114/1662 contained plastic particles	Not investigated	Bond et al. 2013

Table 4.1 (d) Field observations of the occurrence of plastic particles in biota in the South Atlantic Ocean

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
South Atlantic Ocean						
<i>Pachyptila vittata</i> , Broad-billed prion	Gough Island, Central South Atlantic Ocean	October & November 1979	Opportunistic sampling after observing plastic pellets in regurgitated skua pellets	Thirty-three plastic pellets observed with prion bones and feathers in regurgitated predatory Great skua <i>Catharacta skua</i> pellets	Not investigated	Bourne & Imber 1980
<i>Puffinus gravis</i> , Great shearwater	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	11/13 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Pelagodroma marina</i> , White faced storm petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	16/19 birds contained plastic particles in their gizzards	Statistically weak correlation identified between mass of ingested plastic and body mass	Furness 1985a
<i>Pachyptila vittata</i> , Broad-billed prion	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	12/31 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Fregetta grallaria</i> , White-bellied storm petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	5/13 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Garrodia nereis</i> , Grey-backed storm petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	3/11 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Stercorarius antarcticus</i> , Tristan skua	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	1/11 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Puffinus assimilis</i> , Little shearwater	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	1/13 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Lugensa brevirostris</i> , Kerguelen petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	1/13 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Pterodroma incerta</i> , Atlantic petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	1/13 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a
<i>Pterodroma mollis</i> , Soft-plumaged petrel	Gough Island, Central South Atlantic Ocean	September – October 1983	Dead birds collected for analysis and others harvested as necessary for sampling	1/18 birds contained plastic particles in their gizzards	No negative impacts identified	Furness 1985a

Table 4.1 (e) Field observations of the occurrence of plastic particles in biota in the North Pacific Ocean

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
North Pacific						
<i>Fulmarus glacialis</i> , Northern fulmar	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	16/19 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Puffinus tenuirostris</i> , Short-tailed shearwater	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	4/5 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Oceanodroma furcata</i> , Fork-tailed storm petrel	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	18/19 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Oceanodroma leucorhoa</i> , Leach's storm petrel	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	31/64 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Phalacrocorax pelagicus</i> , Pelagic cormorant	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	2/10 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Larus canus</i> , Mew gull	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	1/4 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Rissa tridactyla</i> , Black-legged kittiwake	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	20/256 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Rissa brevirostris</i> , Red-legged kittiwake	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	4/15 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Uria aalge</i> , Common guillemot	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	1/134 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Ptychoramphus aleuticus</i> , Cassin's auklet	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	4/35 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Aethia psittacula</i> , Parakeet auklet	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	195/208 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Aethia cristatella</i> , Crested auklet	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	1/40 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Cephus columba</i> , Pigeon guillemot	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	1/43 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Fratercula corniculata</i> , Horned puffin	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	44/120 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995
<i>Fratercula cirrhata</i> , Tufted puffin	Alaska	1988–1990	Targetted sampling of seabirds for feeding ecology studies	120/489 birds had ingested plastic particles (mainly pellets)	Not investigated	Robards et al. 1995

Table 4.1 (f) Field observations of the occurrence of plastic particles in biota in the Central Pacific Ocean

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
Central Pacific						
Phalaropus fulicarius, Red phalarope	California	May 1980	Opportunistic sampling of 7 dead migratory birds	6 birds contained plastic particles in their stomachs. Most were 'nibs' and some was styrofoam	Not investigated	Connors & Smith 1982
Pterodroma rostrata, Tahiti petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/121 birds had ingested plastic	Not investigated	Spear et al. 1995
Puffinus nativitatus, Christmas shearwater	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	2/5 birds had ingested plastic	Not investigated	Spear et al. 1995
Puffinus pacificus, Wedge-tailed shearwater	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	17/85 birds had ingested plastic	Negative relationship identified between plastic ingestion and physical condition (body weight)	Spear et al. 1995
Fregatta grallaria, White-bellied storm petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/18 birds had ingested plastic	Not investigated	Spear et al. 1995
Oceanodroma tethys, Wedge-rumped storm petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/296 birds had ingested plastic	Not investigated	Spear et al. 1995
Sterna fuscata, Sooty tern	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/64 birds had ingested plastic	Not investigated	Spear et al. 1995
Gygis alba, White tern	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/8 birds had ingested plastic	Not investigated	Spear et al. 1995
Pterodroma externa, Juan Fernandez petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/183 birds had ingested plastic	Not investigated	Spear et al. 1995
Pterodroma cervicalis, White-necked petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/12 birds had ingested plastic	Not investigated	Spear et al. 1995
Pterodroma pycrofti, Pycroft's petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	2/5 birds had ingested plastic	Not investigated	Spear et al. 1995
Pterodroma leucop-tera, White-winged petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	13/110 birds had ingested plastic	Negative relationship identified between plastic ingestion and physical condition (body weight)	Spear et al. 1995
Pterodroma brevipes, Collared petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	2/3 birds had ingested plastic	Not investigated	Spear et al. 1995

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
<i>Pterodroma nigripennis</i> , Black-winged petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	3/66 birds had ingested plastic	Not investigated	Spear et al. 1995
<i>Pelagodroma marina</i> , White-faced storm petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	11/15 birds had ingested plastic	Not investigated	Spear et al. 1995
<i>Oceanodroma leucorhoa</i> , Leach's storm petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	70/354 birds had ingested plastic	Negative relationship identified between plastic ingestion and physical condition (body weight)	Spear et al. 1995
<i>Stercorarius longicaudus</i> , Long-tailed jaeger	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/2 birds had ingested plastic	Not investigated	Spear et al. 1995
<i>Pterodroma ultima</i> , Murphy's petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	1/7 birds had ingested plastic	Not investigated	Spear et al. 1995
<i>Pterodroma longirostris</i> , Stejneger's petrel	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	34/46 birds had ingested plastic	Negative relationship identified between plastic ingestion and physical condition (body weight)	Spear et al. 1995
<i>Puffinus griseus</i> , Sooty shearwater	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	27/36 birds had ingested plastic	Negative relationship identified between plastic ingestion and physical condition (body weight)	Spear et al. 1995
<i>Puffinus bulleri</i> , Buller's shearwater	Tropical Pacific	1989–1991	Targetted sampling of seabirds to quantify incidence of plastic ingestion	3/3 birds had ingested plastic	Not investigated	Spear et al. 1995
<i>Lepas anatifera</i> , a gooseneck barnacle	North Pacific Subtropical Gyre	2009–2012	Opportunistic sampling of floating debris items with barnacles attached to assess microplastic ingestion	90/243 barnacles had ingested plastic	No evidence found of acute harm but negative long term effects could not be ruled out	Goldstein & Goodwin 2013
<i>Lepas pacifica</i> , a gooseneck barnacle	North Pacific Subtropical Gyre	2009–2012	Opportunistic sampling of floating debris items with barnacles attached to assess microplastic ingestion	34/85 barnacles had ingested plastic	No evidence found of acute harm but negative long term effects could not be ruled out	Goldstein & Goodwin 2013
<i>Astronesthes indopacifica</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 1.0 plastic particles and 0.03 mg plastic per fish gut	Not investigated	Boerger et al. 2010

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
<i>Cololabis saira</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 3.2 plastic particles and 1.97 mg plastic per fish gut	Not investigated	Boerger et al. 2010
<i>Hygophum reinhardtii</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 1.3 plastic particles and 1.82 mg plastic per fish gut	Not investigated	Boerger et al. 2010
<i>Loweina interrupta</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 1.0 plastic particles and 0.64 mg plastic per fish gut	Not investigated	Boerger et al. 2010
<i>Myctophum auro lanternatum</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 6.0 plastic particles and 4.66 mg plastic per fish gut	Not investigated	Boerger et al. 2010
<i>Symbolophorus calliforniensis</i>	North Pacific Central Gyre	February 2008	To determine if small mesopelagic planktivorous fish were ingesting plastic fragments	Mean of 7.2 plastic particles and 5.21 mg plastic per fish gut	Not investigated	Boerger et al. 2010

Table 4.1 (g) Field observations of the occurrence of plastic particles in biota in the South Pacific and Indian Oceans

Marine species	Location	Date of study	Purpose of study	Details of ingestion	Reported impact of ingestion	Reference
South Pacific Ocean						
<i>Ardenna tenuirostris</i> ¹² , Short-tailed shear-water	Phillip Island, Victoria, Australia	April – May 2010	Opportunistic sampling of 67 dead, beach cast chicks analysed for stomach content and body condition	Mean mass of plastic per bird was 113 ±8.2 mg (not certain that all was microplastic). No relationship identified between plastic mass or number of particles found and fat scores	None identified	Carey 2011
<i>Arctocephalus</i> spp., Antarctic fur seals	Macquarie Island, Australia		Opportunistic sampling of scats for plastic particles	All 145 scats contained at least 1 plastic particle. Majority 3–5 mm length. Hypothesized that they were ingested via predation of <i>Electrona subaspera</i> by the fur seals	Not investigated	Eriksson & Burton 2003
Indian Ocean						
<i>Stolephorus commersoni</i> , Commerson's anchovy	Kerala, India	August 2013	Targetted sampling as part of a larger study into mud banks	6/16 individuals had microplastic particles in their gut	Not investigated	Kripa et al. 2014

¹²*Ardenna tenuirostris* – formerly known as *Puffinus tenuirostris*

Table 4.2. Selected laboratory studies of microplastics exposure in marine organisms

Marine species	Plastic particle exposure and effect	Reference
Blue mussel <i>M. edulis</i> to crab <i>C. maenas</i>	Transition 0.5 µm PS from mussel to crab by trophic transfer	Farrell & Nelson 2013
Blue mussel <i>M. edulis</i>	Translocation to hemolymph of mussel after ingestion; 3–9.6 µm, into blood cells including macrophages 0–80 µm. Transfer into cells	Browne et al. 2008; von Moos et al. 2012; Höher et al. 2012; von Moos et al. 2012
Blue mussel <i>M. edulis</i>	Exposure to 10, 30 90 µm MPs Indications for selective uptake of 10 µm MPs Reduced clearance rate	Van Cauwenberghe et al. 2012b, 2013a
Zooplankton & mysid shrimp	Neomysis integer Ingestion, trophic transfer of fluorescent 10 µm PS from zooplankton to mesozooplankton Trophic transfer	Setälä et al. 2014
Shore crab <i>Carcinus maenas</i>	Uptake via gills of 8–10 µm PS and ingestion. Retention in foregut Watts et al. 2014 Echinoderm larvae Active capture and ingestion, 20 µm	Hart 1991
Lancet fish	Polystyrene beads 0.05–25 µm Unrestricted intake of polystyrene beads, max 100 µm	Ruppert et al. 2000
copepod		
Centropages typicus	Microbeads 10-70 µm, selective ingestion of 59–65 µm	Wilson 1973

4.2.3 Uptake and transition into tissues, cells and organelles

Up until now only a very few studies have examined the presence of microplastics in tissues or body fluid of field-collected organisms. Evidence for such internal microplastic exposure relates mainly to filter feeding mussels and sediment deposit feeding polychaetes (see Table 4.2) and is described below.

In experimental studies, marine mussels – a species also used for human consumption – were exposed to seawater containing microplastics accumulated plastic particles in the hemolymph and once the particles were ingested they were able to move from the gut to the circulatory system and be retained in the tissues (Browne et al. 2008). Van Cauwenberghe et al. (2012b, 2013a) exposed *Mytilus edulis* (filter feeder) and *Arenicola marina* (deposit feeder) to different sizes of microplastics at a total concentration of 110 particles per mL. After exposure, lugworms had on average 19.9 ± 4.1 particles in their tissue and coelomic fluid, while mussels had on average 4.5 ± 0.9 particles in their tissue and 5.1 ± 1.1 per 100 µL of extracted hemolymph (Van Cauwenberghe et al. 2012b, 2013a). Experimental studies by Browne et al. (2008) have shown that microscopic polystyrene particles 3 and 9.6 microm of size are ingested and accumulated in the gut of the mussel *Mytilus edulis* and translocate to the circulatory system within 3 days and were taken up by hemocytes. Studies with HDPE powder in a size range of >0 to 80 µm

by von Moos et al. (2012) demonstrated intracellular uptake of microplastic particles into the cells of digestive tubules and transition into cell organelles of the lysosomal system.

Accumulation of plastic inside of lysosomes coincides with breakdown of the lysosomal membrane and release of degrading enzymes into the cytoplasm causing cell death (von Moos et al. 2012). A strong immune response towards those HDPE particles expelled from the digestive tubules into the surrounding storage tissue and fibrous encapsulation of the plastic engulfing macrophages. In the same exposure study Höher et al. (2012) evidenced uptake into hemolymph from the digestive system, with specific uptake of HDPE into proliferating granulocytes and basophilic hemocytes.

Ecotoxicological experiments on sea urchin embryos showed that plastic pellets that have not entered the marine environment have a stronger effect on embryo development than beach collected ones (Nobre et al. 2015), suggesting that leaching of additives would have a higher toxicity than organic pollutants absorbed into the stranded pellets. Toxicity of beach collected pellets depended on the ecotoxicological method and varied among trials, suggesting variability in toxicity among samples of pellets at the beach.

Potential Endocytotic Pathways for Nanoparticle Entry into Cells




Figure 4.1. Pathways for endocytosis in the cell which could be exploited by manufactured and defragmented microplastic. Endocytosis via clathrin-coated pits (receptor mediated) or uncoated pits (fluid phase) transfers materials to the lysosomal degradative compartment, while caveolar endocytosis can result in translocation to the endoplasmic reticulum (ER), Golgi or through the cell by transcytosis (Shin and Abraham 2001; van der Goot and Gruenberg 2002) modified by Moore 2006.

Uptake routes into (eucaryotic) cells are evolutionary conserved and particles traverse by diffusion selectively by the size of pores in cell membranes, of which only 2% are large sized pores (400 nm pore radius); 30% of medium size (40 nm); and 68% of small size (1.3 nm). Specialized coated vesicles transport particles smaller than 200 nm. In contrast, larger particles up to 40 micrometres are taken up by engulfment through endocytosis and phagocytosis.

In general, in vivo exposure to small sized microplastic indicates that, due to the physiological mechanisms involved in the feeding process, nanoparticle agglomerates taken up by the gills are directed to the digestive gland where intra-cellular uptake of nano-sized materials induces lysosomal perturbations (Canesi et al. 2012) (Figure 4.1).

4.2.4 Excretion

In the field, the occurrence of microplastics in organisms represents recent exposures to these materials. Our current knowledge of the excretion of microplastics from marine organisms is based solely on laboratory studies. After microplastics are assimilated into the

organism (hemolymph or tissues) they either accumulate or are excreted depending on the size, shape and composition of the particles used in the studies. If accumulated, the chemical and/or physical effects are expected to develop and to be maintained through time. If excreted, these effects are expected to be reversed in a healing and regeneration process.

Microplastic concentrations in the hemolymph peak at a certain time after a single discrete exposure (which depends on species, plastic type and exposure time) and then reduce in abundance (Browne et al. 2008; Farrell & Nelson 2013;). It is not known how much is eliminated or transferred to other organ compartments or tissues. Recent studies in mussels after acute exposure to HDPE (0–80 µm size range) for 12 h followed by regeneration in plastic-free seawater indicate elimination of microplastic from the digestive tubules after a period of 12 to 48 hours, and a shift of HDPE particles into newly formed connective tissue (fibrosis) around the tubules, indicating a repair mechanism of injured tissue. Similar experiments with PVC microplastics revealed particle retention in the gut for up to 12 days and that small size particles had a higher retention time than larger ones.

4.2.5 *Transfer of microplastics in the food web*

Microplastic particles may be passed through the food web as predators consume prey. Farrell and Nelson (2013) fed mussels (*Mytilus edulus*) which had been exposed to 0.5 µm polystyrene microspheres to the crab *Carcinus maenas*. Microplastics were found in the stomach, hepatopancreas, ovary and gills of the crabs, and the maximum amount of microplastics were detected 24 hours post feeding. Nearly all of the ingested microplastics were cleared from the crabs after 21 days. In a laboratory feeding study, all ten zooplankton taxa tested from the Baltic Sea ingested 10 µm polystyrene microspheres (Setälä et al. 2014). Microspheres contained within copepods were transferred after mysid shrimps ingested them, again demonstrating trophic transfer of microplastics. Lusher et al. (2013) documented microplastic particles in the gastrointestinal tracts of 36% of 504 individual fish collected from the English Channel, representing 10 species of teleost fish, confirming ingestion of microplastics in prey items in the field. Similarly, Murray and Cowie (2013) found microplastics (mostly plastic strands) in the gut contents of 62% of Norway lobsters (*Nephrops norvegicus*) collected from the Clyde Sea, and confirmed in companion laboratory studies that ingested plastic fibres were effectively retained with the GI tract.

4.3 Microplastics as a vector of chemical transport into marine organisms

4.3.1 *Introduction*

Microplastics in marine environments carry chemicals that can be considered as contaminants from an ecotoxicological risk perspective, from two principal sources (Teuten et al. 2009). The first includes the additives, monomers and by-products contained in plastic particles. The size, condition and residence time of microplastic particles and the hydrophobicity of the chemicals controls how much of these chemicals are retained by microplastics in marine environments. The second type of contaminants are hydrophobic compounds and metals sorbed from surrounding seawater. This includes most POPs and PBTs, including polycyclic aromatic hydrocarbons (PAHs) and the other petroleum hydrocarbons. Sorption will tend towards equilibrium between the plastic and seawater, phases with the direction of absorption/desorption depending on the absorption kinetics and the relative concentrations in the two media. The size of microplastics, polymer type and hydrophobicity of the contaminant will all exert an influence.

This characteristic has been employed in sampling methods to isolate and concentrate dissolved POPs from natural waters (e.g. Adams et al. 2007; Cornelissen et al. 2008), and to assess POP/PBT bioavailability in water and sediments (e.g. Leslie et al. 2013). Indeed, the International Pellet Watch Programme (IPW) deliberately uses marine microplastic particles as passive samplers of organic contaminants in waters throughout the world (Ogata et al. 2009; IPW website: <http://www.pelletwatch.org/>). The magnitude of the POP-plastic

interaction depends on the nature of the plastic as well as the hydrophobicity of the chemical, as quantified by several studies (Teuten et al. 2009; et al. 2012; Endo et al. 2005).

The rate of uptake and release of POPs from plastics largely depend on size of plastics. With increase in the size (thickness) of plastics, sorption and desorption becomes slower. To thin polyethylene film with thickness of 50 µm PCBs (CB52) sorbed in 50 days to reach equilibrium (Adams et al. 2007), whereas sorption of PCBs to PE pellets with diameter of 3 mm is slower (Mato et al. 2001) and takes approximately one year to reach equilibrium (Rochman et al. 2012, 2013b, 2014a). The slow sorption and desorption was explained by slow diffusion in aqueous boundary layer (Endo et al. 2013) and/or in intra-particle (matrix) diffusion (Karapanagioti and Klontza 2008). In addition, the slow desorption may transport POPs to remote ecosystems. Plastic pieces that contain higher concentrations of POPs than the other plastics are often found on the beaches of remote islands (e.g. Hirai et al. 2011; Heskett et al. 2012).

However, recent laboratory experiments demonstrated that stomach oil acts as organic solvent, facilitates the elution of PBDEs from the plastic matrix and enhances bioavailability of the additive chemicals (Tanaka et al. 2013). Because stomach oil is common among many species of seabirds and plastic ingestion is observed for various species of seabirds, wider species of seabirds in various regions should be investigated in future efforts. However, only limited species of seabirds and the other animals have been examined. Magnitude and expansion of the chemical transfer by microplastics in the biosphere should be studied.

4.3.2 *Contaminant transfer from µm-size plastics to lower-trophic-level organisms*

Well-established pharmacokinetic-based models have established that dietary uptake is often the majority mechanism of POPs exposure for fish and shellfish (e.g. Thomann et al. 1992; Arnot and Gobas 2004). While diffusive uptake of dissolved POPs across the surfaces of skin, gills and other respiratory surfaces should not be ignored, it is the ingestion of contaminated prey items and subsequent transfer of POPs from food to organism that supplies most of the contaminant burden. As reviewed in this chapter, a growing number of studies demonstrate that, under the right conditions, many species of marine organisms will ingest microplastic particles. As organisms consume a mixed diet consisting of a variety of particles (including perhaps microplastics), the total dietary exposure equals the weighted average POPs concentration of the mixture weighted for selective assimilation. Assimilation efficiency is defined as the fraction of the POP concentration in a prey item or food particle that is transferred into the organism, and is a function of the diffusional gradient between food and organism and the gut residence time. Species-specific features, including especially digestive enzymes and active transport mechanisms, also influence the POP assimilation efficiency. It is important to note that net POPs transfer from food to organism is thought to only occur when the POPs concentration in the partially digested

food exceeds that in the organism, driving the diffusional gradient across the digestive walls (Kelly et al. 2004). Therefore, the magnitude of dietary exposure to POPs depends explicitly on over- or under-enrichment of POPs in the food relative to the organism, regardless of how enriched the food is relative to the surrounding water or sediment.

Partitioning, bioavailability and assimilation

Several studies document that marine microplastics are covered with biofilm communities (Zettler et al. 2013 and references therein). This organic layer likely acts as a reservoir for POPs, although studies demonstrating persistent differences in POPs affinities among plastic types deployed in marine waters suggest that the biofilms modify rather than control POPs associations with aged marine microparticles. When these particles are ingested, it is likely that the nutritionally rich biofilm organic matter is stripped from the particles, with coincident release of the co-adsorbed POPs. Emerging studies have suggested that contaminant transfer does take place (e.g. Chua et al. 2014; Rochman et al. 2014b; Browne et al. 2013; Gaylor et al. 2011). However, whether biofilms facilitate the transfer of contaminants into the organism has not been investigated.

As described earlier, many organisms are able to translocate assimilated microplastic particles within their tissues, storing the foreign body within intracellular structures. It is possible that this defence mechanism would deliver microplastic-associated POPs and additive chemicals to different tissue types and locations than those resulting from uptake from food and water. Given the long residence time of such sequestered particles relative to the lifetime of the organism, even slow chemical release may cause low but chronic delivery within the animal. This mechanism has not been studied in marine organism and is not included in current pharmacokinetic models. The relatively low frequency of occurrence and density of tissue-encapsulated microplastics in field collected marine specimens suggests this mechanism is likely not an important vector in overall chemical delivery to marine species. However, it is possible that this unstudied mechanism may provide a unique process to deliver chemicals to specific organs, especially for very small plastic particles that can cross membranes. Further quantitative investigations, especially heuristic application of existing models to explore the potential importance, are required to confirm this. Considerable literature exists examining the release of chemicals from polymers and other macromolecules in human systems, where these solid materials are used as drug delivery vehicles.

As noted above, transfer of POPs and additive chemicals from food to organism depends on the relative concentration of chemicals in the food compared to what already exists in the animal. Due to the influence of growth dilution (i.e. the increase in organism weight due to growth reduces the POPs concentration in the tissue even though the total POPs mass in the organism remains the same), the diffusional gradient in the gut is often positive (i.e. the food is enriched in POPs relative to the organism). However, the opposite situation may occur, where relatively contaminated organisms ingest cleaner food. In this case, partitioning of POPs from the organism back into the food during digestion,

with subsequent excretion, is possible. This process is analogous to those used to depurate commercial shellfish prior to harvest and limit the effects of ingested poisons by administration of activated carbon.

Model approaches to assess the role of μm -size plastics on contaminant accumulation in lower-trophic-level organisms

In a recently published series of papers, the issue of the net effect of microplastics on the transfer of PCBs to the deposit feeding lugworm was measured (Besseling et al. 2012) and modelled (Koelmans et al. 2013). By enhancing an established pharmacokinetic model to include the microplastic-influenced processes discussed above, the authors were able to quantitatively address the impact of microplastics as a delivery vehicle for PCBs to marine organism. As their experimental design included extremely high microplastic concentrations (up to 7.4% by weight) in sediment, the work represents a 'worse case', skewing the experiment in favour of finding any possible influences. The laboratory experiment exposed lugworms to three levels of polystyrene and constant total levels of PCBs and measured feeding activity, growth rates, and PCB congener accumulation over time. The model included partitioning of PCB congeners to the added microplastic, the ingestion of microplastic particles, and the dynamics of POPs transfer within the worms.

When simulating a closed system where there is a finite amount of POPs, Koelmans et al. (2013) determined that partitioning competition between the added plastic, sediment and pore-water and the transfer of PCB congeners from the organism to relatively clean plastic during gut transport dominated the overall net POPs exposure. At the end of 30 days, lugworms living in highly altered sediments (7.4% polyethylene) are predicted to accumulate 25 times less PCBs than those living in plastic-free sediments, a difference that was not attributed to changes in exposure moderated by the plastic rather than the minor alterations observed in behaviour, feeding rates, or growth rates. Modelled dissolved PCB exposure from pore-water decreased as PCBs partitioned onto the added plastic.

To explore a more realistic scenario, Koelmans et al. (2013) also model behaviour in an open system, where there is a very large pool of available POPs (i.e. adding plastic to the system does not deplete the dissolved or sediment bound POPs concentrations). In this scenario, addition of even high concentrations (10% by weight) of polystyrene did not affect PCB bioaccumulation (<1% decrease). Because polyethylene has a higher affinity for PCBs, however, the simulation suggests a 2–5 fold decrease in PCB bioaccumulation for polyethylene concentrations in the sediment ranging from 0.1% to 10% by weight. This results from the compensating mechanism of plastic ingestion offsetting PCB uptake by dermal exchange and diet. Although partitioning among pore-water, sediment and plastic begins at equilibrium, feeding and digestion increases the organism PCB concentrations following the standard bioaccumulation paradigm. Since the plastic remains in equilibrium with the surrounding pore-water and sediment, it is now 'clean' relative to the POPs level in biota, resulting in net POPs elimination by back partitioning to ingested plastic particles.

The two important implications of this recent work are (1) when considering all POPs exposure mechanisms simultaneously, addition of POPs-free microplastics if anything decreases bioaccumulation of POPs in deposit feeding organisms, and (2) the magnitude of the calculated impacts on bioaccumulation are quite small relative to typical variation observed in the field.

Laboratory experiments on transfer of POPs from μm -size plastic to lower-trophic-level organisms

In recent years, there have been several studies based on laboratory experiments to evaluate plastic-mediated transfer of POPs to lower-trophic-level organisms. Browne et al. (2013) studied transfer of four kinds of organic contaminants (nonylphenol, phenanthrene, BDE47, and triclosan) from PVC to benthic organisms (lugworm). PVC particles of 230 μm were pre-sorbed with these contaminants at environmentally relevant concentrations and added to clean sands where lugworm was incubated for 10 days and concentrations of the contaminants in the tissue were measured. Concentration of the PVC was 5% of sand. All the contaminants were significantly detected in gut of the lugworm following 10-day incubation, clearly indicating the transfer of the organic chemicals from microplastic to the biological tissue. They also incubated the lugworm in sand presorbed with nonylphenol and phenanthrene without microplastics and found higher concentrations of the pollutants than those on the incubation with microplastics. This could be explained by retention (partitioning) of the hydrophobic pollutants by PVC in sands as predicted by above model approach. Because Browne et al. (2013) utilized clean sands, importance of plastic-mediated transfer relative to natural path (i.e. pollutant uptake through contaminated sediment) was not evaluated. Besseling et al. (2012) studied the relative importance of the transfer of PCBs from natively PCBs-contaminated sediments to similar species of the benthic organisms (lugworm). They added three levels (0.074 %, 0.74 % and 7.4 % dry weight) of polystyrene (PS) particles (400 – 1300 μm) to the sediment to examine the enhancement or depression of bioaccumulation of 19 congeners of PCBs in the lugworm tissue. Microplastic concentration (740 mg/kg or more) on this experiment was one to three order of magnitude higher than those observed in actual environments (up to 81 mg/kg, Reddy et al. 2006). Increased PCB accumulation was observed for several congeners at the lowest PS dose of 0.074 %. Though the increase was statistically significant, the increment was small (i.e. 1.1 – 1.5 times than control (no addition of PS)) and was observed for limited congeners (i.e. CB31, 52, and 105) and no increase was observed for many congeners. These results are basically consistent with the estimation by above model approach. The laboratory experiments and the model approach can conclude that transfer of organic pollutants from μm -sized plastic to tissue of benthic organism occurs but its role to pollutant transfer could be less important relative to natural path at the present condition (i.e. lower concentration of microplastics than natural sedimentary organic matter). However, more studies are necessary to improve the model approach so that it can represent the increase in bioaccumulation of some congeners observed by Besseling et al. (2012). Also field observation to examine the conclusion is neces-

sary. Furthermore, it is concerned that the concentrations of microplastic in sediment and relative importance of microplastics would be increasing in future. Even now, we may underestimate the concentrations of μm -sized or nm-sized plastics in sediment due to less development of the methods for the collection, identification, and counting of the tiny plastic particles in bottom sediments.

Regarding μm -sized plastics in water column, there have been several laboratory experiments to examine the transfer of pollutants to organisms. Chua et al. (2014) exposed amphipod to μm -sized (11 – 700 nm) polyethylene particles which were pre-sorbed with a range of PBDEs (BDE28, 47, 100, 99, 154, 153, and 183). They measured PBDEs concentrations in tissue of the amphipod after 72-h exposure and detected all the BDE congeners in the tissue, indicating the occurrence of transfer of the chemicals from microplastic to the tissue of amphipod. In addition, they observed higher PBDEs uptake by amphipod in control, i.e. system without microplastics, than those with microplastic, again suggesting the retention of hydrophobic pollutants in microplastics which is consistent with above model approach. They fed only microplastics to the amphipod and, therefore, their role of PBDE transfer relative to natural path was not assessed. Rochman et al. (2013a) conducted laboratory experiments where Japanese Medaka was exposed to polyethylene pieces less than 500 μm . The PE pieces were pre-sorbed with variety of POPs through deployment in polluted waters for 3 months and mixed with natural food stuff including cod liver oil and fed to the fish. Statistically significant enhancement of PBDE bioaccumulation was observed for fish fed with contaminated microplastic, suggesting the transfer of PBDEs from microplastic to the organisms. However, no significant increase was observed for PCBs and PAHs mainly due to higher concentrations of POPs in fish fed with natural food stuff only. This may indirectly suggest microplastic play only minor role to bring POPs to biota in comparison to natural path.

A feeding experiment using chicks of Streaked shearwater (*Calonectris leucomelas*) supported this idea. Chicks were fed plastic resin pellets collected from Tokyo containing PCBs together with their natural food (Japanese sand lance) and periodically monitored for PCBs in the preen gland oil excreted from gland at the tail of the bird (Teuten et al. 2009; Yamashita et al. 2007). Significant increases in concentrations of lower-chlorinated biphenyls were observed one week after the feeding of the plastics, suggesting the transfer of the PCBs from the plastics to the tissue of seabird. However, they did not see any significant difference in the PCB concentrations after two weeks. This was also explained by the “natural” exposure of the seabirds through their food (prey) because the plastic was fed only at the beginning of the experiments and PCB amounts exposed through natural food overwhelm the exposure from the ingested plastic in the later stage of the experiment. This is also consistent with the studies that correlated plastic ingestion and PCB concentrations in the fatty tissue of seabirds, Short-tailed shearwater (Yamashita et al. 2011). Their approach is similar to Ryan et al. (1988) and found significant but weak correlation between mass of ingested plastics and tissue concentration of lower-chlorinated congeners.

Field evidence of transfer of contaminants from μ m-size plastic to lower-trophic-level organisms

There has been very limited number of field studies to examine the POPs transfer from microplastic to lower-trophic-level organisms. Rochman et al. (2014b) measured various organic pollutants in fish tissue collected in South Atlantic gyre where accumulation of microplastic was observed. They observed higher concentrations of PBDEs in the tissue of lanternfish collected at areas with higher plastic accumulation in water column. This correlation suggests the important role of plastic-mediated transfer of POPs. The correlation was obtained probably because concentrations of plastic were higher relative to natural food items, e.g. zooplankton. Similarly to the benthic system, the relative contribution of microplastic-mediated POPs transfer depends on the concentration of microplastics relative to natural food items. In some locations, such as the Central Pacific gyre, concentrations of plastic were higher than those of plankton. Similar areas may be observed in coastal zones in proximity to intensive anthropogenic activity, such as Tokyo Bay (Hideshige Takada, unpublished results). Furthermore, there is concern that there will be an increase in the extent of such areas unless there are substantial reductions in the input of plastics to the ocean.

The Koelmans et al. (2013) work applies to the test organism (lugworm) and POPs (PCB congeners) studied in sediments. As several studies have now demonstrated microplastic ingestion in field-caught fish and shellfish (Lusher et al. 2013; Murray and Cowie 2011), the question remains as to the potential role of microplastics in POPs accumulation in the water column. However, the relatively low number of microplastics in marine waters relative to food items coupled with potential selective feeding suggests that microplastics play a minor role among the large variety of natural particles in delivering POPs to higher trophic levels.

Metals

The microplastics collected from the ocean contain metals, although in very small concentrations (Ashton et al. 2010; Holmes et al. 2012). Weathered particles have a higher potential to accumulate metals from the environment compared to new plastic (Ashton et al. 2010; Fotopoulou & Karapanagioti 2012), although in both cases the magnitude of metals accumulation is quite low relative to natural particles. Nakashima et al. (2012) showed that metals in stranded macroplastics may be leached to the surrounding water and release of metals should be enhanced in acid conditions like in the gut of many organisms. The relative importance of microplastic-facilitated transport of metals into aquatic organisms has not been directly measured.

4.3.3 Field evidence of contaminant transfer from mm-size plastics to higher-trophic-level organisms

There have been several field studies to examine the transfer of POPs from ingested plastics to marine organisms with a focus on seabirds. Ryan et al. (1988) measured amounts of plastics in the digestive tracts

of Great shearwaters (*Puffinus gravis*) and concentrations of PCBs in the fat tissue of the bird and examined the correlation between amounts of plastics and PCB concentrations. A positive correlation was observed between the mass of ingested plastic and the PCB concentration in the fat tissue of birds, suggesting the transfer of PCBs in plastics to the organisms. However, correlation was weak because marine organisms, especially higher-trophic animals, are exposed to PCBs through natural prey in addition to ingested plastics.

Tanaka et al. (2013) examined the transfer of PBDEs (an additive used as a flame retardant for certain applications) from the ingested plastics to the tissue of the seabirds with focus on higher brominated congeners, not found in prey items of the seabirds. PBDEs in abdominal adipose of oceanic seabirds (short-tailed shearwaters, *Puffinus tenuirostris*) collected in northern North Pacific Ocean were analysed. In 3 of 12 birds, higher-brominated congeners (viz., BDE209 and BDE183) which were not present in the natural prey (pelagic fish) of the birds were detected. The same compounds were present in plastic found in the stomachs of the 3 birds. These data and their follow-up observations of the same species of seabirds indicated the transfer of plastic-derived chemicals from ingested plastics to the tissues of marine-based organisms. However, the mechanism of the transfer of the chemicals from the plastics to the biological tissue was not revealed. Because the ingested plastics were relatively large (mm to cm-size) and BDE209 and BDE183 are highly hydrophobic, slow release and low bioavailability of the chemicals have been suggested.

4.4 Biological impacts

Section 3 of this report makes apparent that microplastics have become both widespread and ubiquitous in the marine environment. The biological impact of microplastics on organisms in the marine environment is only just emerging (Gregory 1996; Barnes et al. 2009; Ryan et al. 2009; Cole et al. 2011). Adverse effects of microplastics on marine organisms can potentially arise from physical effects (physical obstruction or damage of feeding appendages or digestive tract or other physical harm). In addition, microplastics can act as vectors for chemical transport into marine organisms causing chemical toxicity (additives, monomers, sorbed chemicals). Translocation of microplastics from the gut to other tissues may result in 'internal' exposure of microplastics causing particle toxicity and particle dependent chemical toxicity of leached chemicals. In addition to impaired health, the ingestion of microplastics could potentially cause population, community level effects and affect ecosystem wide processes. Microplastics could also serve as carrier for the dispersal of chemicals and biota (invasive species, pathogens), thus greatly increasing dispersal opportunities in the marine environment, potentially endangering marine biodiversity. The wide range of potential effects of microplastics on marine organisms and ecosystems is further explored below.

4.4.1 Physical effects

The small sizes of microplastics make them available to a wide range of marine organisms posing a potential threat to biota (Derraik 2002; Barnes et al. 2009; Fendall and Sewell 2009; Thompson et al. 2004; Cole et al. 2011). This may be particularly the case for small-sized deposit and suspension feeders, such as zooplankton, polychaetes, crustaceans and bivalves. Microplastics may present a mechanical hazard to small animals once ingested, similar to the effects observed for microplastics and larger animals (Barnes et al. 2009; Cole et al. 2011). The physical effect may be related to entanglement (no records, expected for larger particles), obstruction of feeding organs, e.g. salps (Chan & Witting 2012) and zooplankton (Cole et al. 2013), reduction in the feeding activity/rate/capacity (Besseling et al. 2012; Cole et al. 2013), or adsorption of microplastics on the organism surface, e.g. algae (Bhattacharya et al. 2010) and zooplankton (Cole et al. 2013). Direct effects may occur after ingestion and translocation into tissues, cells and body fluids causing particle toxicity. These studies are summarized in Table 4.2.

So far, only few field studies have attempted to investigate the impacts of microplastics on marine organisms and these refer to larger sizes of microplastics in birds and fish. For example, sub-lethal or lethal effects were difficult to relate to plastic ingestion in fulmars (van Franeker et al. 2011) and plastic characterization seemed unrelated to physical condition in the shear-water species (Codina-García et al. 2013). Foekema et al. (2013) found no clear relation between the condition factor of North Sea fish and the presence of ingested microplastics. However, the authors noted that the particles were probably too small to expect they can cause feelings of satiation and intestinal blockage potentially resulting in a decreased condition factor in the relatively large specimens examined.

In contrast to the field situation, effects of microplastics have been reported in numerous laboratory studies with a wide variety of fish and invertebrate species exposed to relatively high concentrations of virgin/unpolluted microplastics (Table 4.2). Bhattacharya et al. (2010) worked with nano-sized plastic beads and two species of algae (one freshwater and one marine/freshwater species) and found that sorption of nano-plastics to algae hindered algal photosynthesis and appeared to induce oxidative stress. Microplastic adhered to the external carapace and appendages of exposed zooplankton which can significantly decrease function and algal feeding (Cole et al. 2013). A recent study showed that ingestion of microplastics by lug-worms leads to decreased feeding activity in sediments that contain 7.4% polystyrene (Besseling et al. 2012). In the same species Wright et al. (2013b) found statistically significant effects on the organisms' fitness and bioaccumulation, but the magnitude of the effects was not high.

Many marine organisms have the ability to remove unwanted natural materials including sediment, natural detritus and particulates from their body without causing harm. However, once ingested there is the potential for microplastics to be absorbed into the body upon passage through the digestive system via translocation

and causing particle toxicity with associated inflammation and fibrosis. These studies are summarized in Table 4.3 and a number of studies are highlighted below.

It is known that xenobiotic microplastic particles accumulating in organs and tissues may evoke an immune response, foreign body reaction and granuloma formation (Tang & Eaton 1995). A few studies of marine organisms have clearly demonstrated such direct particle toxicity effects of microplastics translocated from gut to body fluids into organs, cells, and even organs. Mussels were exposed to primary HDPE plastic powder >0-80 µm, which was absorbed by digestive gland vacuoles (von Moos et al. 2012). Accumulation of plastic inside of lysosomes coincided with breakdown of the lysosomal membrane and release of degrading enzymes into the cytoplasm causing cell death. A strong immune response towards those HDPE particles which were expelled from the digestive tubules into the surrounding storage tissue was diagnosed. The plastic engulfing macrophages were encapsulated by fibrous tissue, resulting in granulocytoma formation during the exposure time of 96 h. In the same exposure study Höher et al. (2012) evidenced uptake into hemolymph from the digestive system as Browne et al. (2008) with specific uptake of HDPE into proliferating granulocytes and basophilic hemocytes reflecting the strong immune response also seen in the digestive gland.

4.4.2 Comparison with observed effects in mammalian systems

A large body of evidence in the peer-reviewed literature reveals that microparticles of plastic in the human body or mammals are unhealthy. In the absence of studies for marine biota, the effects of particles observed in human cells and tissues or in animal models give an insight into the possible risks of particle exposure in other organisms and in humans. Humans occupy a high trophic level in the marine food chain, and can potentially be exposed to micro- and nano-plastics (especially primary micro- and nano-plastics) while using products that contain them.

Mobility of tiny plastic particles of various size ranges in the human body has been demonstrated in studies of uptake in the gastrointestinal and lymph (Hussain et al. 2001), crossing the human placenta (Wick et al. 2011). The many studies of fine solid particles in air have taught us what particulates can do in human and mammalian tissues and how they negatively impact health through causing allergic reactions, asthma, cancer and heart disease. In the human system much has been learned from PE and PMMA exposures when implants made of these materials degrade and particles are released into the human body causing particle-induced osteolysis (e.g. Martinez et al. 1998; Petit et al. 2002; Nich et al. 2011).

Table 4.3 Selected laboratory studies of biological effects of microplastics exposure in marine organisms

Marine species	Plastic particle exposure and effect	Reference
Phytoplankton	Adsorption of 20 nm PS	Bhattacharya et al. 2010
<i>Scenedesmus</i>	Hindered algal photosynthesis and promotion of algal ROS indicative of oxidative stress	
Zooplankton, various species	Uptake of 1.7–30.6 µm PS beads varying by taxa, life-stage and bead-size. Microplastics adhered to the external carapace and appendages of exposed zooplankton. 7.3 µm microplastics (>4000 mL ⁻¹) significantly decreased algal feeding.	Cole et al. 2013
	Uptake of 0–80 µm MPs into digestive system, into digestive tubules with translocation into cells and cell organelles (lysosomes). Granulocytoma formation (inflammation). Increase in SB haemocytes; decrease in lysosome stability.	von Moos et al. 2012
Blue mussel	Exposure to 10, 30 90 µm MP (incl PS)	Van Cauwenberghe et al. 2013a
<i>Mytilus edulis</i>	Indications for selective uptake of 10 µm MPs. Reduced clearance rate	Wegner et al. 2012
	30-nm PS. Reduced filtering/feeding activity	
Lugworm (<i>Arenicola marina</i>)	PS microplastic. Statistically significant effects on the organisms' fitness and bioaccumulation, but the magnitude of the effects was not high.	Besseling et al. 2012
	Exposure to 130 µm (mean size) PVC conc. 1% of sediment (w/w) reduced total energy reserves by approximately 30%, mainly linked to a reduction in lipid reserves.	Wright et al. 2013b
copepod	Exposure to 0.05, 0.5 and 6 µm PS .100% survival in 96h tox test. Chronic mortality for 0.05 µm PS >12.5 µg/mL.	Lee et al. 2013b
<i>Tigriopus japonicus</i>	Reduced fecundity for 0.5 and 6 µm PS.	
Carp species <i>Carassius carassius</i>	Absorption of 24 nm NPs. Food chain transport of NPs affects behaviour and fat metabolism.	Cedervall et al. 2012
Brown mussel <i>Perna perna</i>	Leached and virgin microplastics (PVC and PE microbeads), in different concentrations (0.5 and 2.5 g/L) and periods of exposure (6, 12, 24, 48, 96 and 144h) indicated physical (and not chemical) impacts.	Preliminary results of master's thesis of Liv Ascer and Marina Santana
	Altered expression levels of AIF and HSP70 proteins and lysosomal stability in the cells in mussels exposed to microbeads during 144h and in mussels exposed to microbeads during 144h, irrespective to concentration.	
Sea urchin <i>Lytechinus variegatus</i>	Plastic pellets (virgins and beach-collected) affect embryonic development, but virgin ones have a stronger effect.	Nobre et al. 2015
copepod	Exposure to ≥4000 particles/mL of 7.3 µm PS significantly decreased copepod feeding on algae.	Cole et al. 2013
<i>Centropages typicus</i>		
copepod	Exposure to 75 particles/mL of 20 µm PS for 24 hrs significantly decreased copepod feeding capacity. Prolonged exposure significantly decreased reproductive output (egg hatching success and survival)	Cole et al. 2013
<i>Calanus helgolandicus</i>		
Japanese medaka <i>Oryzias latipes</i>	Fish fed a diet containing virgin PE pellets and pellets with sorbed PAHs PCBs and PBDE congeners, having been exposed in a contaminated marine environment, show evidence of hepatic stress	Rochman et al. 2013a

In a study of exposure to ultrafine polystyrene particles in rats, lung inflammation and enzyme activity were impacted, in a dose-dependent way, the greater the surface area to volume ratio of the particle. Toxicity increased in direct proportion to a decrease in particle size from 535 nm to 202 nm to 64 nm polystyrene (Brown et al. 2001). Many other effects of ultrafine plastic were measured in vitro in the same study, including induction of increases in IL-8 gene expression in epithelial cells and an increase in cytosolic calcium ion concentration. The authors suggest that these particle-induced calcium changes may be significant in causing pro-inflammatory gene expression, such as chemokines. A large body of literature has been published on the human toxicity of particles, mainly via the inhalation exposure route (e.g. Hesterberg et al. 2010).

More knowledge of the transfer of microparticles, including microplastics and nanoplastics, through biological membranes can also be mined from the drug delivery research literature. There are ongoing investigations of how the bioavailability and uptake of medicines can be improved by way of micro- or nano-particulate carriers (e.g. Hussain et al. 2001 for microplastics and De Jong & Borm 2008; Wesselinova 2011 for some reviews of the emerging field of nanomedicinal applications, including attention to toxicity).

When humans or rodents ingest microplastics (<150 µm) they have been shown to translocate from the gut to the lymph and circulatory systems (Hussain et al. 2001). Wick et al. (2011) recently demonstrated how nano-sized polystyrene particles up to 240 nm in diameter cross the human placenta in perfusion experiments. Synthetic polymers may in some cases be less harmful than the classic engineered nano-plastics. In a recent study, coating toxic carbon nano-tubules with a polystyrene-based polymer was tested with the aim of reducing the cytotoxicity, oxidative stress, and inflammation in an in vivo mice lung test and an in vitro murine macrophage test (Tabet et al. 2011).

These studies from mammals and the medical field issue a warning that when the size of the microparticle approaches the range below approximately a quarter of an mm, adverse effects may start to emerge due to particle interactions with cells and tissues, particle uptake in endosomes, lysosomes, the lymph and circulatory systems and the lungs. These include deleterious effects at cellular level (Berntsen et al. 2010; Fröhlich et al. 2009) or uptake into placental tissue (1et al. 2010) or lymph and circulatory systems (Hussain et al. 2001;). Human exposure is also a concern if seafood containing microplastics is consumed.

4.4.3 Chemical effects

Section 4.3 demonstrated that microplastics serve as a vector of hazardous chemicals including POPs to marine organisms. In many cases, they are also exposed to chemicals through prey and a dominant contribution from microplastic over natural food has been reported for limited species of animals so far. Conclusive evidence of adverse effects caused by chemicals associated with microplastics is difficult to obtain, but progress has been made. Rochman et al. (2013a) demonstrated the transfer of POPs from plas-

tic to fish and adverse effects from environmentally relevant concentrations. Lavers et al. (2014) showed negative correlation of plastic ingestion with some body conditions of a species of seabird (Flesh-footed shearwaters) whose population decrease is of concern. However, contribution of plastic-derived chemicals has not been clearly evidenced to cause declined body conditions and population-level effects.

A laboratory experiment utilizing lugworms (*Arenicola marina*) demonstrated chemicals associated with microplastic cause adverse effects (Browne et al. 2013). The lugworms were exposed to nonylphenol, phenanthrene, BDE-47, and triclosan with PVC and/or Sand. Transfer of the chemicals from microplastics to gut was observed. Reduction of some biological functions was observed. The results indicated that survivorship and feeding were diminished by triclosan associated with PVC. Contribution of plastic-derived nonylphenol to phagocytic activity and lower oxidative status likely induced by nonylphenol and phenanthrene were suggested.

4.4.4 Potential effects on populations, communities and ecosystems

Our knowledge indicating population or higher-level effects caused by micro- and nano-plastics in the marine environment is poor. Microplastic may not only affect species at the organism level; they may also have the capacity to modify population structure with potential impacts on ecosystem dynamics (Wright et al. 2013a,b), including bacteria and viruses. Negative effects on the photosynthesis of primary producers and on the growth of secondary producers, potentially result in a reduced productivity of the whole ecosystem and represent a primary concern.

The bacteria that colonize plastic particles were shown to differ from surrounding water (Zettler et al. 2013) and sediment (Harrison 2012). Biofilm formation on plastic can be temporally variable and is linked to the productivity of the surrounding seawater (Artham et al. 2009). Biofilm is able to be recorded at least one week after exposure, increasing through time, and may influence plastic buoyancy (Lobelle & Cunliffe 2011) and, indeed, microplastics distribution and availability. The increase in biofilm is supposed to increase the food availability to zooplankton (Kirchman and Mitchell 1982), which may have a bottom up effect on plankton communities.

Plastics in the marine environment, floating in the surface or in the water column or in the bottom, may act as new habitats or new food source for marine organisms, the 'plastisphere' sensu (Zettler et al. 2013). In this context, plastics may have an additive effect on floating patches of seaweeds or create new ones in areas where natural floating patches do not occur.

Table 4.3. Selected studies showing translocation of microplastics from the gut to the circulatory system and various tissues and cells in humans and other mammals and mammalian systems

Species	MP exposure and effect	Reference
Human lymph and circulatory system	Absorption of PE particles taken up in lymph and circulatory system from gastro-intestinal tract	Hussain et al. 2001
Human placenta (<i>ex vivo</i>)	Fluorescent 50, 80, 240 and 500 nm PS particles. Particles up to 240 nm were taken up by the placenta	Wick et al. 2011
Rat	535, 202 and 64 nm PS Lung inflammation and enzyme activities were affected, with increasing severity as particle size decreased	Brown et al. 2001
Human airway smooth muscle cell	Fluorescent 40 nm PS particles decreased cell contractility	Berntsen et al. 2010
Human endothelial cells (blood vessels)	Carboxyl PS latex beads in sizes of 20-500 nm were tested. 20 nm PS particles induced cellular damage through apoptosis and necrosis	Fröhlich et al. 2010
Human macrophages	Fluorescent PS microspheres (1, 0.2, and 0.078 μm). Particle uptake of all sizes: on average, $77 \pm 15\%$ (mean $\pm$ SD) of the macrophages contained 0.078 μm , $21 \pm 11\%$ contained 0.2 μm particles, and $56 \pm 30\%$ contained 1 μm particles. Particle uptake steered by non-endocytic processes (diffusion or adhesive interactions).	Geiser et al. 2005
Dog	PVC particles (5–110 μm) appeared in the portal vein and will reach the liver	Volkheimer 1975

Independently from the size of the particle, fouling organisms may find in plastics an additional mean to disperse (Gregory 2009). Individual plastic particles or plastic patches may be considered as stepping-stones (*sensu* MacArthur & Wilson 1967), such as they create new hard bottoms and increase their availability in the ocean. This allows organisms, in a single or multiple generations, to enhance the probability to occur in certain areas in comparison to the situation considering only natural floating debris. A study on the colonization of stranded plastic debris in Arctic and Antarctic islands estimated that human litter more than doubles the rafting opportunities for biota (Barnes 2002).

According to Barnes (2002), Barnes & Milner (2005) and Gregory (2009), this situation may increase the risk of dispersal of aggressive alien and invasive species and thus endanger sensitive coastal habitats. Hypothetically, marine debris (Majer et al. 2012), as also supposed for floating seaweeds (Rothäusler et al. 2012), may also increase gene flow, thus allowing genetic mixing among populations and decreasing genetic variability within populations.

Similarly to floating seaweeds (Vandendriessche et al. 2007), relatively more dense aggregations of floating plastic particles may possibly act as refuges or feeding grounds for fishes. An increase in abundance of microplastics through time was positively correlated to abundance of *Halobates sericeus* and its egg densities (Goldstein et al. 2012). Increasing population densities may have different outcomes at the community and ecosystem level, as discussed for *Halobates micans* (Majer et al. 2012). Depending on the species being favoured by microplastics, one may expect an increase in predatory pressure by *Halobates*, a top-down effect, or in the food supply to *Halobates* consumers, a bottom-up effect.

Depending on the amount and size of the particles, different functional groups may be directly affected by microplastics, compromising ecological process and ecosystem function. As exposed above, adsorption of microplastics in the organism surface, e.g. algae (Bhattacharya et al. 2010) and zooplankton (Cole et al. 2013), was demonstrated to reduce the photosynthetic and feeding rate, respectively. However, what this effect at the base of the food chain could mean for the productivity and resilience of ecosystems in the long term is unknown. Considering that amount of plastics entering the ocean is increasing, plastic degradation produces smaller particle sizes, smaller particles are supposed to be more toxic, and the effect of microplastics at higher levels of organization is supposed to increase, it is possible to suppose an increasing impact of microplastics on marine systems. However, due to their complexity, with species-specific and generalized response from the biota to the presence of microplastics, with external and internal exposures and with physical and chemical effects, which are not well understood, the direction of this effect is hard to be predicted.

Indirect effects may occur due to the presence of small plastic particles in sediments, which were reported to increase permeability, warm more slowly, and reach lower maximum temperatures (Carson et al. 2011).

4.4.5 Potential effects on humans

The potential accumulation of microplastics in the food chain, especially in fish and shellfish (species of molluscs, crustaceans and echinoderms) could have consequences for the health of human consumers. This seems particularly the case for filter-feeding bivalves such as mussels and oysters in Europe but could equally be applicable for deposit-feeding sea cucumber which is more popular in the Asian cuisine.

Microplastics have been shown to be ingested by several commercial species such as mussel, oyster, crab, sea cucumber and fish (Tables 4.1, 4.2). Relatively high concentrations of microplastics were detected in Belgian commercial grown mussels (*Mytilus edulis*) and oysters (*C. gigas*), respectively on average 0.36 ± 0.07 particles g^{-1} wet weight (w.w.), and 0.47 ± 0.16 particles g^{-1} w.w. (Van Cauwenberghe and Janssen 2014). As a result, the annual dietary exposure for European shellfish consumers can amount to 11,000 microplastics per year. Another Belgian study analysed microplastic contamination between consumption mussels and wild type mussels (mainly *M. edulis*), collected at Belgian department stores and Belgian groins and quaysides, respectively (De Witte et al. 2014). The number of total microplastics varied from 2.6 to 5.1 fibres/10 g w.w. of mussel, which compared well with the former study, despite the fact that both Belgian studies used different methods of analysis and size detection limits. A maximum concentration of 105 particles g^{-1} dry weight (d.w.) (d.l. > 10 mm) was reported in wild mussel (*M. edulis*) from the Dutch coast (Leslie et al. 2013); these levels are one order of magnitude higher (13.2 particles g^{-1} ww; using a conversion factor from d.w. to w.w. of 8 assuming a lyophilization rate of 0.12).

Although it is evident that humans are exposed to microplastics through their diet and the presence of microplastics in seafood could pose a threat to food safety (Van Cauwenberghe and Janssen 2014), our understanding of the fate and toxicity of microplastics in humans constitutes a major knowledge gap that deserves special attention. Therefore, an analysis and assessment of the potential health risk of microplastics for humans should comprise dietary exposure from a range of foods across the total diet in order to assess the contributing risk of contaminated marine food items.

4.5 Recommendations for further research

- Examine the extent to which nano-sized plastic particles may cross cell membranes and cause cell damage, under natural conditions, including knowledge and expertise from the medical and pharmaceutical industry (drug delivery).
- Examine the extent to which additive chemicals may cross the gut wall and assess the risk of harm at an individual and population level.
- Examine the extent to which adsorbed organic contaminants may cross the gut wall and assess the risk of harm at an individual and population level.
- Assess the potential health risk of microplastics for humans, including dietary exposure from a range of foods across the total diet in order to assess the contributing risk of contaminated marine food items.
- Examine the potential of microplastics to translocate non-indigenous species, including pathogenic organisms, relative to other transport vectors.
- Examine the potential for accumulations of plastics and microplastics to form additional floating ecosystems.
- Examine species-specific gut conditions that may influence chemical availability and transfer.
- Consider using stable or radioactive labelled compounds (polymers, additive chemicals and absorbed contaminants) to establish the degree of transfer under different conditions.

5 SOCIAL ASPECTS OF MICROPLASTICS IN THE MARINE ENVIRONMENT

5.1 Introduction

The use of plastics has proliferated in recent decades, due to a complex mix of perceived societal and economic benefits. Unfortunately, the rate of increase in use has not been matched by the adoption of suitable systems to control unwanted plastic items. The working group was asked to consider some of the social aspects around the issue of microplastics in the ocean. The scope of this ToR was quite restricted, to explore the potential role of social science approaches for addressing the issue of microplastics, based on the realization that people's perceptions and behaviours contribute to the problem but are also crucial in any solutions suggested. This scoping exercise for the social scientists was rather different from the rest of the report because no established field of research exists in the social sciences that specifically addresses microplastics. Thus, the social scientists were asked

to selectively review relevant approaches rather than producing a quantitative analysis of perceptions and behaviours. The data for a more quantitative analysis are largely lacking, even considering the impact of macro debris, which is a much more recognizable problem.

Perceptions influence the behaviour of:

- a) the general public;
- b) the complex web of industries involved in design, materials science, engineering, manufacturing, advertising and retail;
- c) users of products such as the aquaculture, fisheries, catering, health-care and tourism sectors;
- d) legislators and environmental managers responsible for many different aspects of plastic use and disposal.

people in a US survey believed that the health of the ocean is important for human survival (Ocean Project 1999). When focusing on environmental matters related to the marine environment specifically, several environmental topics are of particular interest to the public, such as climate change, chemical pollution and ocean acidification (e.g. Vignola et al. 2013; Peterlin et al. 2005; Fleming et al. 2006). Similarly, marine debris is commonly noted as one of the most important issues when people are asked whilst visiting the coast (e.g. Santos et al. 2005; Widmer & Reis 2010). In general, microplastics are not mentioned spontaneously in such surveys. This could indicate either a lack of perceived importance, or simply a lack of knowledge and recognition of this particular environmental issue.

One of the largest scientifically based assessments of public perceptions was conducted in Europe, in a survey of 10,000 citizens from ten European countries, where respondents were asked to identify the three most important environment matters regarding the coastline or sea (Buckley and Pinnegar 2011). The survey was conducted in the context of assessing perceptions about climate, but allowed the respondents to express their concerns freely. When stating levels of concern for a number of environmental issues, including overfishing, coastal flooding and ocean acidification, the term 'pollution', particularly water and oil pollution, was mentioned frequently. Marine debris-related terms, such as 'litter', 'rubbish' and 'beach cleanliness' were also reported, but much less frequently (Figure 5.1).

5.2.1 Perceptions of marine litter in general

There is evidence that at least some sections of the public are aware of our dependency on the marine environment. For instance, as early as 1999, 75% of



Figure 5.1. Main responses from a multinational sample from 10 countries ($n = 10,106$) to a qualitative question that asked individuals to state the three main marine environmental matters. Frequency of responses is illustrated by the size of the text, with pollution noted most often (reproduced from Buckley and Pinnegar 2011).

as being more connected to the issue of marine litter and microplastics. With a sample of just under 4,000 respondents from over 16 mostly European countries, the MARLISCO survey found that the majority of respondents were concerned about marine litter and perceived the marine environment as being highly valuable to society. There was a belief that the situation regarding marine litter was worsening, and that most

of marine litter was derived from the sea¹² (B. Hartley unpublished data). This survey also found that all groups significantly underestimated the proportion of marine litter items composed of plastic by about 30% (B. Hartley unpublished data). A separate survey on UK commercial fishers found similar patterns in perception, whereby fishers underestimated the proportion of litter that is plastic, and on average, were unsure whether marine litter was increasing or decreasing (Defra report, forthcoming).

In a Chilean beach visitor survey, most visitors reported that they did not dispose of litter on beaches despite a large proportion of marine debris being left by visitors in general (Eastman et al. 2013; Santos et al. 2005). Even though respondents generally claimed not to be individually responsible, they did identify the overall public to be the main source of debris (Santos et al. 2005; Slavin et al. 2012; Eastman et al. 2013). In terms of the effects and impacts of marine debris, the main problems that beach users identified were related to the impact on marine biota, human health and safety, and attractiveness (B. Hartley unpublished data; Santos et al. 2005; Wyles et al. 2014; Wyles et al. under review). Thus, these findings suggest that beach-users and commercial fishers have a basic understanding of marine litter in general.

5.2.2 Perceptions about microplastics

Individuals' perceptions about macro marine debris are relevant for understanding the overall issue of microplastics, as macro plastic can break down to form secondary microplastics. For the purpose of this report, the extent of individuals' knowledge specifically relating to microplastics is more relevant. Unfortunately, very little has been published in this area. To our knowledge there are only two studies that included any questions specifically related to microplastics. Firstly, a citizen science programme, involving 1,000 school students in Chile, found that the majority of children (aged 10 to 17) had never heard of microplastics before (Hidalgo-Ruz and Thiel 2013). Secondly, in a multinational survey, using a web-based flash survey method (http://ec.europa.eu/public_opinion/flash/fl_388_en.pdf) of over 26,000 Europeans, 78% agreed to the statement that *"the use of micro plastic particles in consumer cosmetic and similar products should be forbidden"* (European Commission 2014).

There has been no published in-depth investigation of individuals' understanding of this issue. Consequently, a pilot study was carried out in 2013, specifically for this report, to begin to explore this area (for the purpose of this report, it will be termed the GESAMP pilot survey; (S. Pahl unpublished data). A sample of 68 adults in a city on the coast of Southwest of England (31 men, 36 females, 1 not stated; average age 43; $SD = 13$ years; age range = 19-71), were questioned about their perceptions of microplastics. Roughly half (53%) had heard of the term 'microplastics'. When asked to define the term, most people described it as *"very small, tiny particles of, miniature pieces of plastic"*. Some people were not able to answer this question, with others giving other answers that were either vague (e.g. *"man-made material"*) or very detailed (e.g. *"carrier bags, lids, toothbrushes"*), or focusing on the degradability (e.g. *"something that never goes away"*). When asked to estimate the size of these particles, the most common responses were that they were microscopic or not visible (25% of responses), less than 1 mm (37%) or less than 5 mm (21%). Of the remaining responses, 12% thought they were larger than 5 mm, with 6% unable to provide an answer. In terms of distribution and abundance, 74% of the respondents believed the quantity of microplastics in the marine environment was increasing, and the majority believed microplastics could be found in the deep sea, surface seawater, on beaches, in marine animals and in polar seas. As illustrated in Figure 5.2, the overall consensus was that a lot can be found on beaches, with less in the deep sea and in polar seas. Consequently, this preliminary research suggests that although about half the participants had heard of the term, the 'public', on the basis of this small rather unrepresentative sub-set, have a limited understanding of the existence and concept of microplastics. Clearly, further research is needed to substantiate this finding.

To explore concern for microplastics specifically, the GESAMP pilot survey also included a question on concern. On a scale from 1 (not at all concerned) to 10 (extremely concerned), respondents were, on average, quite concerned ($M = 6.91$; $SD = 2.63$).¹³ However, respondents were more concerned about other issues such as the health of the natural and marine environment in general, the cost of living and climate change. These findings were consistent with those reported by Potts and colleagues (2011).

¹² The working group considered that there are no reliable estimates of the quantities of plastic entering the ocean. It is conceivable that more than half comes from land but the available evidence suggests that significant regional and local variations in the proportion of sea- and land-derived plastic occur. <http://www.marlisco.eu>

¹³ Not surprisingly, this was lower than the concern expressed by 29 experts in a special workshop on microplastics using the same scale (expert $M = 8.16$, $SD = 1.69$; $t(98) = 2.45$, $p = .02$, $d = .48$ (medium effect)).




Figure 5.2. The frequency of responses to the question about perceived abundance in different areas of the marine environment (perceived distribution) from the GESAMP pilot survey ($n = 68$, S. Pahl unpublished data).

It is worth listing a number of limiting factors that can influence the outcome of a perception survey, and therefore the apparent level of knowledge of or concern over a particular issue. The circumstances of the request may affect the willingness to cooperate, e.g. between educational and organizational settings versus ‘cold-calling’ or a street survey. Younger people with more exposure to social media may be more responsive to using this medium than other members of society. Age, gender, educational attainment and cultural or religious background may influence responses (see below) so larger, ideally representative surveys are desirable. If the respondent has prior knowledge of the topic they may be more willing to express their views. If individuals are active in ‘environmental affairs’ and campaigning then their advocacy may encourage others in their sphere of influence to volunteer their views. As a result, the outcome of any survey of people’s opinions needs to be viewed in the particular context and circumstances of the survey, and conclusions about the wider population cannot necessarily be drawn. When investigating concern specifically, it is important to acknowledge the effect of simply asking questions. For instance, if a general public sample is asked about microplastics, this could suggest that the researcher believes it is something to be concerned about. During data collection for the GESAMP pilot survey, one respondent reported that they had not heard of the term prior to this survey, but as they were being asked about it, they assumed it should be something to be concerned about. Thus, in a field of research in its infancy, a perception survey may at the same time establish opinions as well as trying to measure them.

5.2.3 Public perceptions and coverage in the printed and digital media

An analysis of the reporting of issues in the printed and digital media, or the frequency with which certain words or search terms are used on the internet or in social media, can provide an indication of the level of interest either by the public in general or a particular sub-set of individuals (e.g. a group of ‘activists’). Of course, it would be imprudent to draw exact conclu-

sions as there are many confounding factors that may interfere with extracting reliable information of perceptions of a specific topic. However, usage statistics regarding media resources can indirectly indicate the public’s concern and interest surrounding an issue. For instance, Google Trends can “reflect how many searches have been done for a particular term, relative to the total number of searches done on Google over time” (Google Trends, 2014a). For the purpose of this report, a very basic exploration of the trends relating to marine debris and microplastics was undertaken although some questions remain about how exactly these scores are calculated (see Note in Fig. 5.3 and the results of including slightly different research terms in the subpanels). First, Figure 5.3 (a) shows how searches for ‘marine debris’ and ‘marine litter’ have fluctuated over the past ten years (2004–2014). Second, when ‘Great Pacific Garbage Patch’ was included in the search (Figure 5.3 (b)), the patterns for the former two phrases change drastically, as this new term was evidently more commonly searched and thus had a strong effect (Gaudet 2012; Google Trends 2014b). Third, in terms of ‘microplastics’ and ‘microbeads’, the recent peaks in searches for the terms could potentially suggest an increase in interest and concern in the topic (Figure 5.3 (c)).

Google Trends can offer information on search trends, has good geographical spread (depending on internet access and use of that particular search engine) and is a free source; however, limitations still need to be acknowledged. For instance, the search terms are language dependent, searches cannot combine two terms (e.g. ‘micro’ and ‘plastics’), and results may include unrelated topics (e.g. for hair extensions and as ingredients in products such as pillows and cosmetics). Consequently, this analysis needs to be considered a basic insight in the public’s interest in microplastics. Ideally, results obtained using Google Trends should be validated by correlating them with survey data to see if the same patterns emerge in the more direct approaches (Mellon 2013).




Figure 5.3. The use of the terms (a) marine debris and marine litter; (b) marine debris, marine litter, and the Great Pacific Garbage Patch; and (c) microplastics and microbeads via internet search trends (Google Trends 2014b).

Note. According to Google Trends, “the numbers on the graph reflect how many searches have been done for a particular term, relative to the total number of searches done on Google over time. They don’t represent absolute search volume numbers, because the data is normalized and presented on a scale from 0-100. Each point on the graph is divided by the highest point and multiplied by 100. When we don’t have enough data, 0 is shown” (<https://support.google.com/trends/answer/4355164?hl=en-GB&rd=1>).

In addition to Google Trends, examining newspaper articles offers another perspective, although similar caveats apply in terms of the danger of over-simplifying the relationship between printed media reports and public perceptions. The first scientific publication dedicated to microplastics appeared in *Science* in May 2004 (Thompson et al. 2004). Within a year of this publication, it received regional to international media coverage, with over 50 stories addressing this specific article (see Table 5.1).

Similar to Google Trends, it is possible to review the trends over time in media stories by searching newspaper archives. For the purpose of this report, the terms surrounding ‘micro plastics’ and ‘micro beads’ (with and without spaces and hyphens) were searched within the LexisNexis newspaper archive; however, unlike Google Trends, the results were validated by only including those that made reference to marine pollution and excluded duplicate articles. In total, 29 items were found within the UK national newspapers

between 3 July 2004 and 3 July 2014.¹⁴ Most articles were published in national broadsheet/mid-market newspapers, with a striking increase in interest in the most recent years; however, very few articles appeared in the popular mass newspapers over this time period (‘tabloid’ newspapers, see Figure 5.4).

In addition to these descriptive trends in Google searches and national press articles, other media attention suggests a potential growing interest and concern. For instance, there have been a number of TV documentaries including the Austrian movie “Plastic Planet” (2009) by Werner Boote, the documentary “Midway, Message from the Gyre” (2013) by Chris Jordan and a feature length film “Plastic Seas” by Jeneene Chatowsky (2013). A growing number of websites are dedicated to marine debris and microplastics, such as the NOAA Marine Debris Program (<http://marinedebris.noaa.gov/>); the 5 Gyres Foundation (<http://5gyres.org/>); MARLISCO, Europe (<http://www.marlisco.eu/>), PlasticTides, Bermuda (<http://www.plasticides.org/>), and International Pellet Watch (<http://www.pelletwatch.org/>).

¹⁴ The daily newspapers sampled were: The Times, The Guardian, The Daily Telegraph, The Independent, The Daily Mail, The Express, The Mirror, The Sun, and The Daily Star. The Sunday newspapers sampled were: The Sunday Times, The Observer, The Sunday Telegraph, The Independent on Sunday, The Mail on Sunday, The Sunday Express, The Sunday Mirror and The People.

Finally, dedicated social media campaigns have the potential to bring about change to society. The campaign “Beat the Micro Bead” (www.beatthemicrobead.org) originated in the Netherlands and illustrates public concern regarding microplastics (or microbeads) entering the marine environment from personal care products such as shampoos, toothpaste and lip balm. Up until July 2014, this particular campaign had over 2,000 Twitter followers, 3,200 Facebook ‘likes’ and 17,000 views of YouTube. More recently the campaign has been extended to an international scale and a number of producers have announced their plans to reconsider and potentially phase out the use of microplastics in their products (e.g. Beiersdorf 2014; L’Oréal 2014; Unilever 2014).

Nevertheless, whilst media coverage can influence and indicate the public’s interest and concern, and can lead to changes in the commercial sector, there is a risk that messages may be misinterpreted and perceptions and opinions presented as ‘facts’. A good example is how the observation that plastic accumulated in the North Pacific sub-tropical gyre led to the phrase the “Great Pacific Garbage Patch”, conjuring visions, amongst some, of a huge island of solid garbage floating in

the Pacific.¹⁵ This example is a representation of the media impact on society, by choosing a metaphor concept, as well as an illustration of a dramatic image of certain environmental issue, which can be used as a way to enhance a perceived importance of the issue. In this case, the metaphor succeeded in catching public attention, but fails to carry the accuracy and the complexity of the situation. As O’Neill and Smith (2014) emphasize, images are subject to interpretation by the viewer. Scientific data indicate that there are, in fact, areas of accumulation zones, but these are mainly composed by microplastics floating over an extensive area without an exact size (Goldstein et al. 2013). Some organizations have specifically designed websites to try to clarify some common misconceptions and to give accurate information about it (e.g. NOAA Marine Debris Program).¹⁶ Thus, as well as indicating public interest, the media can be a source of correct (and incorrect) information about a topic, which in turn can influence the general public’s risk perception and understanding of the topic.

¹⁵ <http://www.abc.es/20120416/ciencia/abci-septimo-continente-basurero-flotante201204161033.html>

¹⁶ <http://marinedebris.noaa.gov/marinedebris101>




Figure 5.4. Frequency of newspaper articles with the terms ‘micro plastics’ and ‘micro bead’ within the UK newspapers.

5.2.4 Perceived Risks

Discrepancies in perceived level of risk between experts and the public have been observed in the general risk perception literature. For instance, the public has been found to perceive much greater health risks associated with biotechnology in food than experts have (Savadori et al. 2010; Slovic 1987; 1999). While experts may focus on probabilities or annual fatalities, the public evaluate risks in terms of a number of psychological factors. The two most prominent factors derived from large-scale empirical studies are commonly referred to as ‘dread risk’ and ‘unknown risk’ (Slovic 1987). ‘Dread risk’ is composed of the following perceptions: strong emotional feelings of dread, lack of control over the issue, catastrophic potential and fatal consequences, more risks than benefits, involuntariness and difficulty

reducing the risk. ‘Unknown risk’ is higher when issues are unknown to those exposed and unobservable, with unknown and delayed consequences, and when the issue is seen as new and also relatively unknown to science. From these psychological factors, it is possible to produce quantitative representations of risk perceptions for a range of issues (also known as cognitive maps, see Figure 5.5).

Table 5.1. Examples of media coverage of the article Lost at sea: where is all the plastic? (Thompson et al. 2004)

National Coverage	European Coverage	International Coverage	Commercial Coverage
BBC 1 TV National News	France 2, National TV	BBC World Service	Plastics and Rubber Weekly
BBC 1 TV Newsround	Documentary feature for Complement d'enquete	Canadian Broadcasting Corporation	Chemical and Engineering News, Washington
BBC TV News 24	German Republic Radio SWR2	Canadian West Radio News	Chemistry and Industry Magazine
BBC Radio 4: Today Programme	German Public Radio AR1	Discovery Channel, Canada	Materials Today, UK
BBC Radio 4: You and yours	Liberation, France	National Peoples Radio, USA (2 million)	
BBC Radio 1,2, 3, 4: News	Science et Avenir, France	BBC Radio Falklands	
BBC Radio Scotland	Basler Zeitung, Switzerland	New York Times	
BBC Radio Devon	Frankfurter Allergemine, Germany	Washington Post	
Radio Manchester	Berliner Zeitung, Germany	Washington Times	
The Guardian	Der Spiegel, Germany	Wtop Radio, Washington (1 million)	
Evening Herald, Plymouth	Gazeta Wyborcza, Poland	Wall St Journal	
Western Morning News	Svenska, Sweden	Bloomberg News	
The Scotsman	CORDIS News web site	Boston Globe	
New Scientist web site		Omaha World Herald	
CBBC web site		Houston Chronicle	
BBC Wildlife Web site		Lexington Herald, Kentucky	
		La Hora, Chile	
		Honolulu Advertiser	
		Taipei Times, Taiwan	
		Straits Times, Singapore	
		Pravda, Russia	
		ABC National Network Australia	
		National Radio, New Zealand (250 K)	
		Business Week Magazine	
		National Geographic web site	
		Nature web site	

Moreover, discrepancies in perceived risk between experts and public may also be due to societal processes when dealing with new information, e.g. reported in the media. These have been described as either risk amplification (the public think risks are higher than experts do) or risk attenuation (the public think risks are smaller than experts do). These processes take place in a network of social groups and institutions, including scientists, reporters, the mass media and politicians (Kasperson et al. 2003). Related to this, the level of trust in a source has been shown to influence an individual's risk perception. For example, if individuals distrust the science or management of a hazard, they are likely to perceive a greater risk, which has been found with the use of pesticides and pathogens (Williams & Hammit 2000). This is especially important to consider in the

case of microplastics as the general public cannot assess the abundance and risks of microplastics for themselves (see Anderson & Petersen 2013 for a related observation on nanotechnologies). Consequently, the role of the communicator gains more weight, and factors such as trust in experts become important. From the literature, we know that independent academics are the most highly trusted group of communicators (as opposed to the media or government bodies, for example). Moreover, it is not just expertise that is linked to trust. Trust is highest if a decision-maker is perceived to have both high perceived expertise and warmth (White & Eiser 2006; White & Johnson 2010). 'Warmth' in this context refers to a perception of genuine care and shared values.




Figure 5.5. A cognitive map of perceptions of risks for a number of technological and environmental issues according to dread risk and unknown risk (adapted from Slovic 1987).

In addition to the magnitude of perceived risk of microplastics in general, specific behavioural issues may affect people directly, such as consumption of seafood. As outlined in Section 4, there is a possibility that microplastics may be transferred to humans via seafood. What do we know about the perceived risk and benefits of seafood consumption? Using a cross-sectional survey on 429 Belgium consumers, Verbeke and colleagues (2005) found that respondents believed there were harmful contaminants in fish, explicitly referring to PCBs and heavy metals. We can speculate that microplastics could be seen as another type of contaminant. However, the risks associated with harmful contaminants were rated less important than the beneficial nutrients fish was seen to provide. Overall, fish was seen to offer greater benefits than risks. In other work, risk does not feature highly when considering seafood consumption. Instead, taste and quality of the fish have been found to be the most important predictors in seafood consumption, with price and convenience as the main barriers (Olsen 2004; Weatherell et al. 2003). From this limited literature on seafood

consumption, it seems that people who consume seafood are not highly concerned about the contaminants in seafood, or that people who are very concerned do not consume seafood. In any case, no research has explicitly examined the perceived risks of microplastics in seafood, thus research is required that investigates risk perceptions regarding microplastics in seafood.

5.3 Social and socio-economical impacts

Marine environments offer a range of benefits and services to human society, including recreational uses, food resources, waste management and water quality (Peterson & Lubchenco 1997; UK National Ecosystem Assessment 2011). As a recreational resource alone, visiting the coast or simply viewing images of marine environments can improve a person's mood and cognitive attention, as well as improving physical health indicators such as reducing blood pressure (Felsten 2009; Hipp & Ogunseitan 2011; White et al. 2010; White et al.

2013; Wyles et al. under review). Marine debris is one of the factors that has the potential to undermine the benefits of marine ecosystem services. This immediate, common form of visual pollution has been explicitly rated as unattractive by observers (Tudor & Williams 2006; WHO 2003) and can even be considered as a key reason not to visit particular marine sites (Ballance et al. 2000; McKenna et al. 2011; Moore & Polley 2007; Tudor & Williams 2006; Wilson et al. 1995). As well as being disliked, environments with debris can reduce people's positive mood (Pretty et al. 2005; Roehl & Ditton 1993; Wilson et al. 1995; Wyles et al. under review). While there are these insights about macro marine debris, so far no research has examined the social impacts of microplastics directly. This could be because it is a recent research topic (Depledge et al. 2013) or because microplastics cannot be easily seen with the naked eye (Wilkinson et al. 2007). The smaller the plastic debris gets, the harder it is to recognize.

Cost impacts related to marine debris have already been demonstrated. For instance, it is considered as a problematic and an expensive issue for the coastal tourist industry, which is important revenue for many countries (World Tourism Organization 2013). For the 20,000 km of UK coastline alone, it has been estimated to cost roughly €18 million (\$24 million) each year to remove beach debris to help maintain the tourism revenue (Mouat et al. 2010). The fishing industry also experiences large economic losses due to the time and expense of getting caught in marine debris, clearing nets of rubbish and repairing equipment (Mouat et al. 2010). On the other hand, the ingestion of microplastics by commercially valuable marine species, which has been already reported in fish from the North Pacific Ocean (Choy & Drazen 2013), can also have potential future impact on the fishing industry. Research noted above implies generic contamination of seafood is currently not an influential factor in consumer choice, however, if the general public are later found to be concerned about this issue, the idea of consuming plastics within fish may affect the consumption of seafood, potentially leading to economic losses in the seafood industry.

5.4 The role of individual, group and regional differences

So far, we have described general insights into the perception and social impacts of micro (and macro) plastic and focused on those processes that are common among people. This type of research is similar to marine sciences research describing general, averaged trends, as opposed to comparing different geographical regions or comparing regions that vary in wind or tidal conditions. However, the latter comparative research is also undertaken and complements the picture. In the social sciences this typically focuses on differences in the respondents. It is not within the scope of this assessment to review this comprehensively but we will briefly summarize a few insights into individual, group and geographical differences. First, demographics can play a role in people's understanding, concern and perceived risk. For instance, when examining consumer perceptions of the risks and benefits of fish, younger male respondents with a higher education and without children believed more strongly that seafood

contained harmful contaminants (Verbeke et al. 2005), and women were more aware of the potential health benefits. Individuals with a lower socio-economic status have been found to be less aware of general marine issues and have been reported to leave more litter at the beach (Eastman et al. 2013; Fletcher & Potts 2007; Ocean Project 1999; Santos et al. 2005). A common phenomenon found in the risk perception literature is the "white-male effect" that describes that white men have lower risk perceptions for a broad range of hazards than white females, and than non-white males and females (Finucane et al. 2000; Flynn et al. 1994, see also discussion in Slovic 1999). Various reasons for this difference have been discussed including lower status and power in society and different worldviews. White males were characterized more by hierarchical and individualistic worldviews than the other groups, and reported more trust in technology and less trust in government in a US sample. Finucane et al. (2000) showed that worldview remained important even when age, income, education and political orientation were controlled for statistically, highlighting the importance of general perceptions about the world. The importance of marine litter generally can differ with demographics; for instance, mothers state a high importance of marine litter present on the coast when choosing to visit the coastal environment (Phillips & House 2009).

In addition to these socio-demographic factors, psychological differences in perceptions, worldviews and values are also important. Concern, and also sustainable behaviours, have been found to differ according to value systems. For instance individuals with a greater sense of connectedness to nature state a greater concern regarding environmental issues and perform more pro-environmental behaviours (Davis et al. 2009; Hinds & Spark 2008; Mayer & Frantz 2004; Nisbet et al. 2008). Political orientation has been shown to play a big role for risk perception. For example, Dunlap and MacCright (2008) have shown for the USA that Democrats are more likely to say global warming is happening compared to Republicans, and this gap is widening. Dunlap and MacCright report that 48% of Republican as opposed to 52% of Democrat supporters said global warming was happening in 1997, but this discrepancy had increased to 42% vs. 76% in 2008.

Risk perceptions and knowledge about marine issues can also differ according to stakeholder group. When examining perceptions about marine debris in general, Hartley and colleagues (in preparation) compared different stakeholders including manufacturers, retailers, educators, policy makers and domestic users of the marine environment. Overall, the different stakeholders were similar in their knowledge and concern for marine litter, but there were some subtle differences. For instance, unsurprisingly the environmental groups were more concerned about the issue. All groups also underestimated how much of the marine litter is composed of plastic materials; however, the environmental organizations and coastal/marine industries were closer to the scientific estimate of 75% (OSPAR 2007). Marine litter in coastal habitats was found to be more important among a general public sample, but less so among coastal experts who focused on physical disturbance from recreational activities (Wyles et al. 2014). Other stakeholder differences have also been found in the GESAMP pilot study on microplastics

described above. When compared to the audience poll of scientists and NGO representatives at a symposium dedicated to microplastics, the general public was less concerned than the latter experts (see above).

Finally, in addition to these individual and stakeholder differences, geographical location has also been found to play a role. For instance, within the same country, people who live closer to the coast have been reported to be more aware of marine issues (Buckley and Pinnegar 2011; Fletcher & Potts 2007; Steel et al. 2005a, 2005b). A large-scale survey undertaken by Steel and colleagues in 2003 found generally low levels of ocean literacy (knowledge and understanding) in US adults (Steel et al. 2005a, 2005b). While education, socio-economic status, age and gender were shown to be associated with respondents' levels of ocean literacy, those respondents who lived in coastal states and who visited the coast frequently were found to be more knowledgeable than those who rarely spent time at the coast. This suggests an association between direct experience and familiarity with the issues (Fletcher et al. 2009). Indeed, personal attachment to the marine environment has been shown to be a key factor in generating a sense of marine citizenship among UK citizens (see McKinley & Fletcher 2010).

Differences have also emerged between countries. For instance, a survey that gathered responses from 7,000 individuals from seven European countries found that Germany placed great importance on ocean health, whereas the UK, Poland, Spain and Portugal rated ocean health relatively low compared to other issues (Potts et al. 2011). When deciding to visit a particular coastal site, another cross-national study found differences between countries. For people in Ireland and Wales, cleanliness was seen as the most important factor, whilst Turkish and US samples noted cleanliness to be the second most influential factor, with distance to the coast being more important (McKenna et al. 2011). Whilst stakeholder differences were quite small, numerous national differences were also found in the survey reviewing individuals' understanding of marine litter, with countries expressing differences in perception of distribution of marine litter, source, experience of marine litter and composition of the rubbish (Hartley et al. in preparation). Some variations in concern was also reported, with Portugal, Slovenia and the UK reporting a greater concern about marine litter, and Romania, Cyprus, Denmark and the Netherlands the least. For the one question examining microplastics in a cross-national European sample, similar differences were found whereby respondents agreeing with the statement that microplastics should be forbidden from cosmetic products varied between countries 53% (Estonia) to 85% (France and Croatia) (European Commission 2014). The GESAMP pilot survey did not find reliable effects of age and gender (S. Pahl unpublished data). However, little research has been dedicated to microplastics specifically, and this gap needs to be filled.

5.5 Overcoming barriers and towards solutions

Selected insights from the psychological and more generic social science literature have been described above, and, where possible, applied to microplastics

specifically. Although only a snapshot of some relevant themes, this body of work highlights how we can integrate social aspects when working towards solutions. It is advisable to integrate science and social science efforts early on in order to make sure the public is informed of the issue and potential solutions. Certainly, many initiatives can contribute to the goal of increasing the awareness on the impacts of microplastics. There is no one perfect solution to the issue, but there are numerous approaches, such as changing the legislation, improving plastic waste facilities and management, changing plastic use and consumption, and education and public engagement should be part of these. A combination of these factors will help address the issue of microplastics in the oceans. Overall, understanding risk (and benefit) perception is important, as these have been found to influence behaviour and acceptability of regulatory approaches (see Zlatev et al. 2010 for an example in the domain of health). For instance, Klöckner (2013) recently reviewed the influence of a number of factors contributing to a range of environmental behaviours. He highlighted that psychological factors play a fundamental role in people's behaviour. Specifically, the lower people's perceived responsibility and capability of addressing the issue, the less likely they were to take action. This could be relevant for driving solutions towards reducing microplastics. In terms of marine debris in general, the MARLISCO survey found that Governments and policy makers were seen as the agents most responsible for tackling the issue, whilst those who were the most capable were seen to be the environmental groups rather than the Governments and policy makers (Hartley et al. in preparation). People may also ascribe responsibility and capability to technological innovations. Rather than focus on personal solutions (e.g. reduce litter), there is often a fondness for technological solutions that can fix problems (Slovic 1999). This could include microplastics, as there has been recent interest in a potential technological solution of mechanical cleaning of the open ocean. Unfortunately, ascribing responsibility and capabilities to technological solutions could encourage a negative spill-over effect, promoting an unintended behaviour of people not managing their waste responsibly. Such a spill-over effect would be particularly concerning if the technological fix was actually ineffective. Worryingly, this has already been indicated in marine debris in general. In the US, it was found that people litter more when they perceive the item to be biodegradable (Keep Los Angeles Beautiful 2009).

Consequently, any solution to this global multidimensional environmental issue would need to consider people's current perceptions and apply multiple approaches that complement one another, addressing both the microplastics entering the environment (e.g. management of waste) and those already in the environment (e.g. mass clean ups). Additional social research is required in order to understand people's current perceptions of the issue of microplastics. In addition to reviewing the limited research on microplastics and making inferences from related research, this chapter has highlighted where work dedicated to microplastics explicitly is missing. For instance, very little is known about individuals' knowledge and understanding, perceived risks, and the associated consequences of microplastics specifically on humans.

Future research needs to consider methodological aspects. For instance, studies on individuals' knowledge and understanding of microplastics should consider regional, demographic, and individual differences. Also, surveys should include a wider geographical coverage, since to date only a select number of countries is represented (e.g. US and Europe). Finally, new studies should address the economic consequences of microplastics. At the same time, attention should be focused on potential solutions and their acceptability, including social 'solutions' such as education, public engagement and behaviour change campaigns.

5.5.1 *Education and Public Engagement*

Education and public engagement are often referred to as ways of improving public understanding and working towards social solutions for environmental problems such as microplastic accumulation. In terms of education, there are two main ways to increase the public understanding of environmental issues: formal and informal education (Dori & Tal 2000). Formal education refers to the inclusion of scientific topics in centralized curricula in a country's educational systems. Informal education refers, amongst other contexts, to volunteering projects where self-directed, voluntary learning is guided by individual interests and needs. For instance, beach clean-ups occur across the world, organized by several environmental organizations (e.g. Ocean Conservancy, Project AWARE). An example is the annual international beach clean-up day, where in 2013 approximately 650,000 participants from 92 different countries were involved (Ocean Conservancy 2014). These activities represent educational opportunities that have been useful for enhancing local engagement of the issue (Storrier & McGlashan 2006), and increases in marine awareness have recently been demonstrated (Wyles et al. under review; Hartley et al. in press). It can also encourage other pro-environmental behaviour, for instance fishers engaged in the Fishing for Litter scheme run by KIMO (where KIMO pays for the waste disposal of the rubbish caught by the registered fishers during their working day; KIMO 2014) reported reducing waste entering the marine environment, both at work and during their leisure time, than fishers not involved in this scheme (Defra report, forthcoming). Thus, engaging stakeholder groups with marine debris through informal education and engagement can have numerous benefits.

On the other hand, volunteer participation can also be useful for scientific research, a field referred to as citizen science (Cohn 2008; Bonney et al. 2009). Working with volunteers is beneficial in many respects: it allows for more information to be collected when resources are limited, it can be used as an educational tool, and it can promote environmental concern and stewardship in participants (Anderson & Alford 2013). Certain projects have already been looking at marine debris, predominantly focusing on determining the distribution and composition of the debris and their impact on marine biota. Some examples are the Marine Conservation Society from the UK, Our Sea of East Asia (OSeA) from South Korea (see Hong et al. 2013; 2014) and the National Marine Debris Monitoring Program from US (see Ribic et al. 2011; 2012). The only programme that has been working specifically with the

topic of microplastics, is the programme Científicos de la Basura (Litter Scientist) from Chile (Bravo et al. 2009; Eastman et al. 2013; Hidalgo-Ruz & Thiel 2013). This programme is an outreach citizen science project with school children throughout Chile and Easter Island. The information provided by these volunteers has contributed to a better knowledge about large-scale patterns of marine debris and microplastics in the SE-Pacific and also to raising environmental awareness in the region (Hidalgo-Ruz & Thiel 2013). Four main steps have been applied: (a) Contact and commitment of the participants, (b) Development and explanation of sampling protocols and motivational materials, (c) Application of activities and recovery of the obtained data, and (d) Writing of the report and release of the information to local and national press. The experience of these projects suggests that involving the public in studies of marine debris and therefore microplastics can support the enhancement of scientific literacy about microplastics, but can also bring about awareness and an attitude change of this ecological issue in society. Nevertheless, more research is needed specifically evaluating the long term effect and influence of these activities on public understanding of science and environmental awareness (Hidalgo-Ruz & Thiel 2013).

Although these education and public engagement activities are necessary and important parts of engaging society, it is important to allow for proper interactions between scientists, experts and the public rather than the one-way notion that scientists ought to feed the right information to the public. People's perceptions, views and behaviours are influenced by a complex set of factors (as noted above) that can appear to be irrational to someone outside of that group. It is important, therefore, to recognize these differences and take them into account when seeking explanations for the status quo and mechanisms for delivering change (Slovic 1999). Consequently, if we want to engage people, we need to meet them on their terms and acknowledge their construction of the risk issue. Only by building on people's perceptions and using their strengths will we achieve lasting change and consensual solutions (e.g. Pahl et al. 2014).

5.6 Recommendations for further research

To conduct empirical social research on microplastics to address: a) individuals' knowledge and understanding; b) perceived risks; and, c) the associated consequences on humans. Social perceptions are linked to behaviour and support of measures addressing the issue.

To improve the geographical representativeness of this work – outside North and South America and Europe – to identify needs and tailor information to account for social, economic and other cultural differences, and promote effective mitigation strategies.

To analyse the economic impacts of microplastics, in terms of cost-benefit to forecast future effects in response to any changes in microplastic use/input.

Promote the collection and evaluation of examples of public engagement programmes (e.g. citizen science; beach cleans) in terms of their effects on perceptions and actions, including longitudinal follow-ups.

6 KEY OBSERVATIONS AND CONCLUSIONS

The following represent some of the key observations and conclusions that emerged during the assessment process. They are intended to be understandable by a non-technical audience. Readers are encouraged to refer to the preceding Sections 3, 4 and 5 for further explanation, discussion and a summary of the evidence. A confidence level of 'high', 'medium' and 'low' has been given to each statement, based on a consensus within the working group.

6.1 Sources, distribution and fate of microplastics

High confidence

1. The term 'microplastics' has been adopted to describe small plastic particles, generally <5 mm in diameter, sampled in the environment.
2. Commonly available techniques restrict the minimum particle size sampled, currently 10s to 100s microns in diameter. However, it is likely that plastic particles a few nanometres in diameter are present in the environment; hence microplastic fragments span a size range of over 5 orders of magnitude.
3. Microplastics may be manufactured for particular applications or result from fragmentation of larger items. They can be released as a result of many different human activities, but there are no reliable estimates of the quantities entering the marine environment, at a regional or global scale.
4. Most microplastic particles are composed of the six major polymer types. Those composed of polyethylene, polypropylene and expanded polystyrene are more likely to float, and those composed of polyvinyl chloride, polyamide (nylon) and polyethylene terephthalate (PET) are more likely to sink.
5. The surface of any solid object rapidly becomes coated with inorganic and organic compounds and biofilms when immersed in seawater. This may cause floating plastic particles to sink.
6. Plastics will tend to absorb and concentrate hydrophobic contaminants from the surrounding seawater. In addition, additive chemicals incorporated during manufacture may represent a significant proportion of the particle composition.
7. After entry into the ocean microplastics can become globally distributed and have been found on beaches, in surface waters, seabed sediments and in a wide variety of biota (invertebrates, fish, birds, mammals), from the Arctic to Antarctic. They become concentrated in some locations such as ocean gyres, following long-distance transport, but also close to population centres, shipping routes and other major sources.
8. A very high degree of spatial and temporal variability in particle distribution has been observed, partly linked to small-scale circulation and mixing processes in the upper ocean.

9. It seems unlikely that a cost-effective technical solution can be developed and maintained to allow the large-scale removal of significant quantities of floating microplastics from the ocean. Any proposed scheme would be ineffective as long as plastics and microplastics continue to enter the ocean.

10. Better control of the sources of plastic waste, through applying the principles of the 3 Rs (Reduce, Re-use, Recycle), and improving the overall management of plastics via the circular economy, represents the most efficient and cost-effective way of reducing the quantity of plastic objects and microplastic particles accumulating in the ocean. The working group agreed the urgency of implementing effective measures to reduce all inputs of plastic to the ocean.

11. Even if all releases of plastic to the environment were to cease immediately, the number of microplastics in the ocean would be expected to continue to increase as a result of continuing fragmentation, on the basis of current evidence.

Medium confidence

12. Although it appears likely that the quantities of microplastics in the ocean are increasing, due to the continuing fragmentation of existing plastic objects, there is very limited reliable evidence about variations in the abundance of microplastics in space time, despite the availability of some long-term datasets.

13. Dissolved metals will become adsorbed to the surface of plastic particles, in a similar manner as metals adsorb to inorganic sediment particles.

6.2 Effects

High confidence

1. Microplastics are ingested by a wide range of marine organisms including invertebrates, fish and birds and in some organisms the incidence of ingestion is widespread across populations.
2. The movement, storage and elimination of microplastics by marine organisms will depend on the size of the particle. Particles at the smaller end of the size spectrum (nano scales) have been shown to cross membranes into cells, in controlled laboratory experiments.
3. When microplastics cross cell membranes, some tissues have been shown, in vitro, to exhibit a response to the presence of particles; i.e. causing inflammation and cell damage, followed by healing responses and fibrous encapsulation of particles.
4. The risk of associated effects following exposure to microplastics will depend on: i) the number of particles; ii) the size distribution, shape, surface properties, polymer composition and density of the particles; iii) the duration of exposure; iv) the kinetics of absorption and desorption of contaminants, with respect to the plastic and the organism; and, v) the biology of the organism.

Medium confidence

5. Marine organisms are exposed to microplastics via the same pathways used to food, including filtration, active grazing and deposit feeding, and via transport across the gills.

6. Emerging evidence suggests that some additive chemicals, which can be present in relatively high concentrations in some particles, transfer across the gut and concentrate in tissue, under natural conditions. Absorbed contaminants have been shown to exhibit similar behaviour in the laboratory, but there is not yet published, unequivocal evidence to demonstrate that this occurs under natural conditions. The relative importance of contaminant exposure mediated by microplastics compared to other exposure pathways remains unknown.

7. Microplastics may be transferred from prey to predator, but the process will be species-specific. Currently there is no evidence to support or refute potential bio-magnification of particles or associated chemicals.

8. Among the various types of seafood, consumption of filter feeding invertebrates, such as mussels or oysters, appears the most likely route of human exposure to microplastics. However, there is no evidence to confirm this is occurring.

9. The ingestion of microplastics may have an effect on the feeding, movement, growth and breeding success of the host organism.

Low confidence

10. If there are effects on individuals, there is a potential to have impacts at a population level for some species, but this is very uncertain.

6.3 Social aspects

High confidence

1. The presence of macro debris has been recorded to have negative social and economic impacts, reducing the ecosystem services and compromising perceived benefits.

2. Public engagement and education is a useful tool to help raise awareness and promote positive behaviour change, whilst we further develop our scientific knowledge.

Medium confidence

3. Even though there appears to be little awareness in terms of microplastics specifically, there is a general awareness of marine litter and marine threats as a broader concept.

4. Based on the risk assessment literature, perceptions of risk and impacts are influenced by individual, stakeholder group and cultural and other factors. There is no reason to think reactions to the risk of microplastics will be any different.

5. Understanding individuals' perceptions (knowledge, concern, perceived risk) and the impact of microplastics on individuals are important as these have influential consequences. For instance, they influence political pressure, personal behaviour change, acceptance of new products and influence commercial impacts (e.g. to stop using products that contain microplastics).

Low confidence

6. Based on limited survey reports, there appears to be little awareness of the specific issue of microplastics amongst the general public. However, according to on-line media trends, there appears to be increasing attention being paid to this topic. There is evidence of the increasing use of smartphone applications and social media in relation to plastic issues, but it is not clear what proportion of the public is engaged, and the global significance of such initiatives.

7. Based on inferences from similar environmental issues and the generic risk perception literature, it is likely that the perception of risk will increase if public knowledge of microplastics increases.

8. Microplastics are thought to have the potential to have negative socio-economic impacts. For example, if people perceive a high risk e.g. of transfer through seafood (influencing the fishing industry).

7 KEY POLICY-RELATED RECOMMENDATIONS

7.1 Rationale

This assessment represents the first attempt, at a global scale, to identify the main sources, fate and effects of microplastics in the ocean. As a result we have an improved understanding of the scale of the problem posed by the presence of microplastics, and of the link between larger plastic items (macro- and mega-plastics) and the generation of 'secondary' microplastics. It has been possible to state, with more confidence, what is known (Section 6), although there remain con-

siderable areas of uncertainty that will require further investigation.

Policy-makers, and other decision-makers in the public (e.g. municipalities) and private sectors (e.g. manufacturing, retail, tourism, fisheries), need guidance now on how best to target the microplastics issue. This was considered at some length, recognizing that there is a need for decisions to be made before all possible evidence has been collected. In addition, it was recognized that a number of outstanding issues remain

which would benefit from more detailed assessment. For these reasons two sets of policy-related recommendations have been proposed:

1. Action-orientated recommendations addressing marine microplastics
2. Recommendations to improve a future assessment(s)

These policy-related recommendations are based on the detailed analysis described in the main sections of the report. Each is preceded by a specific challenge to be addressed, and is followed by a series of potential solutions considered to be the most cost-effective. They are in addition to a number of technical recommendations included in the main body of the report, which are largely concerned with identifying research needs. They are also linked to the key conclusions (Section 6) that summarize what is known with reasonable certainty.

Any litter reduction measures that are proposed need to be targeted, effective, equitable and cost-efficient. It is critical that measures are designed to minimize unintended, adverse consequences. For example, replacement of plastic by glass bottles on tourist beaches may increase the incidence of injury, if littering behaviour persists. Inadequate separation of waste streams during plastics recycling may result in the contamination of consumer goods, such as children's toys, with unnecessary additives.

Reducing the input of marine debris to the marine environment is a complex 'wicked' problem (Brown et al. 2010), with many possible part-solutions requiring input, commitment, acceptance and resources from a wide range of individuals, institutions,

industries and socio-economic sectors, such as manufacturing, retail, tourism, fisheries, aquaculture and shipping, as well as society at large. These recommendations are designed to support this process.

7.2 Action-orientated recommendations addressing marine microplastics

7.2.1 Challenge 1

Comprehensive improvement to waste generation and management practices is essential in order to reduce the entry of plastics and microplastics into the marine environment. This requires an adequate understanding of the relative importance of different types of materials and sources at global, regional and sub-regional scales, and the socio-economic sectors involved. The input of macro-plastics and microplastics is highly variable and poorly quantified on a regional basis, presenting great difficulties in designing and implementing cost-effective mitigation strategies. Reducing the input of macro-plastics represents the most effective way of minimizing the increase in the abundance of microplastics in the ocean.

Recommendation 1: identify the main sources and categories of plastics and microplastics entering the ocean

Suggested solution/response:

Identify probable 'hotspots' of land- and sea-based sources for plastic and microplastics, using a combination of targeted modelling, knowledge of actual and potential sources (e.g. coastal tourism, aquaculture, fisheries, riverine inputs, urban inputs), environmental and societal data. This will allow mitigation measures to be better targeted, and used to predict and verify their effectiveness. Examples of 'hot spots', from available evidence, include: the Bay of Bengal, Mediterranean Sea, Gulf of Mexico, Japan Sea and other far eastern seas.¹⁷ This can help to inform the development of effective measures in other regions.

Risk of not addressing this challenge:

Efforts to reduce plastics entering the ocean may be poorly targeted, and fail to yield tangible results. Scarce resources may be wasted which could have been better directed.

7.2.2 Challenge 2

There is growing acceptance, especially at an institutional level, that land- or sea-based inputs of plastic waste to the ocean should be reduced. A lack of capacity, technical or financial resources is often cited as a barrier to the implementation of effective waste reduction strategies. In addition, decision-makers are faced with many other demands, affecting the local environment, economy, society and political process, which may be given higher priority in allocating effort and funding.

Recommendation 2: utilize end-of-life plastic as a valuable resource rather than a waste product

Suggested solution/response:

There is great potential in promoting the 3 Rs (Reduction, Re-use and Recycling) as a key contribution to reducing plastic waste generation and reducing the input of plastic to the oceans. This will be aided by the development of innovative and effective solutions as an intrinsic part of the circular economy.¹⁸ In this way 'unwanted' plastic can be seen as a useful resource, with commercial value, rather than a waste problem requiring the allocation of scarce public and private sector resources. Such action reduces our reliance on non-renewable reserves of oil and gas to produce plastics and reduces the need for waste management, for example via landfill. This is a rapidly

¹⁷ Ocean gyres represent accumulation zones but, in general, are too remote to directly link debris with a particular source.

¹⁸ <http://www.ellenmacarthurfoundation.org/business/reports>

developing field that is being embraced by business and institutions, and it needs to be encouraged at a global level. However, adequate controls have to be in place to ensure that plastic waste streams are separated appropriately, to reduce the potential for unnecessary and unwanted cross-contamination, especially of consumer products made with recycled plastic. Commercially available 'biodegradable'¹⁹ plastics do not offer a viable alternative, and in most cases will not lead to a reduction in microplastic formation.

With this change in philosophy, other approaches from the business and commercial sectors may be useful, such as the use of value-chain models. This can help to guide the optimum use of resources, identify intervention points and provide opportunities for economic incentives throughout society,²⁰ with an end-point being the reduction of inputs of marine debris.

Risk of not addressing this challenge:

It will be more difficult to bring about a significant reduction in plastics entering the ocean, and this may come to be seen as an inevitable consequence of economic growth. When faced with hard choices about allocating public and private sector resources, it may be difficult to justify expenditure on 'ocean pollution', which is avoidable, compared with more immediate societal needs (e.g. health service, education and other economic investment).

7.2.3 Challenge 3

Legislation alone will be insufficient to substantially reduce the input of plastic waste into the ocean. It will require a change in perceptions in the public and private sectors, as well as society more widely, as these play a key role in influencing decisions and behaviour. This applies in many areas of risk perception and management. For example, emerging evidence suggests that perceptions influence waste generation and management in general and, more specifically, littering behaviour. It will also influence whether society will be willing to support better waste management practices, such as recycling schemes.

Recommendation 3: promote greater awareness of the impacts of plastics and microplastics in the marine environment

Suggested solution/response:

Utilize expertise from the social sciences, including psychological studies, to better understand perceptions of risk, social responsibilities and the drivers of behaviours in the public and private sectors. Facilitate the transfer of complex and uncertain scientific findings into a language that can be understood by target stakeholder groups (e.g. industrial production, manufacturing, retail, fisheries, aquaculture, coastal tourism,

¹⁹ see Section 3.2.4 for an explanation of 'biodegradable' plastics

²⁰ for example: plastic bottle deposits, fishing net return schemes

shipping). Take proper account of regional, cultural, gender, economic, educational and other demographic differences, in assessing perceptions and behaviours. This can embrace the outputs from the assessments of social resilience and governance, conducted as part of the GEF Transboundary Waters Assessment.²¹

Risk of not addressing this challenge:

It will be more difficult to gain political agreement, public and private sector commitment, and public acceptance, to pursue direct mitigation measures for litter reduction, as well as encouraging the introduction of the circular economy and the benefits of treating unwanted plastics as a resource. It will be more difficult to encourage particular key socio-economic sectors, and the public in general, to adapt their behaviours and contribute to the overall goal of reducing marine plastics.

7.3 Recommendations to improve a future assessment

7.3.1 Challenge 4

The word 'microplastics' has tended to be used as an 'umbrella' term covering particles ranging in size over several orders of magnitude, from particles several mm in diameter to those in the nano size range (<1 µm). Particles in these size ranges may be introduced directly or may gradually form by fragmentation. The sampling methods normally used to collect microplastics tend to exclude material <330 µm. This means there is very limited information about the occurrence of finer plastic particles, including nano-plastics, in the ocean and particularly the degree to which nano-plastics interact with biota internally.

The available evidence from the medical, pharmaceutical and toxicology literature suggests that nano-sized particles are much more likely to cross cell membranes and induce a response that may adversely affect marine life. Therefore they have the potential to pose a greater risk to the organism than micro-sized plastics. Unfortunately, at present it is not possible to provide a credible assessment of the extent to which nano-plastics do present a risk.

Recommendation 4: include particles in the nano-size range in future assessments of the impact of plastics in the ocean

Suggested solution/response:

Encourage the inclusion of expertise on pharmacology, mammalian toxicology, nano-polymer sciences and nano-engineering in future assessments. Critically review laboratory-based experiments examining the behaviour and potential effects of nano-plastics and assess their relevance to the natural environment. Assess current sampling and detection methods for nano-sized plastic particles, particularly in biota.

²¹ Transboundary Waters Assessment Programme, a full-size GEF project, 2012–2014; <http://geftwap.org>

Risk of not addressing this challenge:

The extent to which nano-size plastics pose a significant risk to the marine environment and human health will remain unknown. Meanwhile, there is a very high probability that the quantities of nano-plastics in the marine environment will increase substantially, due to the fragmentation of larger plastic particles and, potentially, due to the direct introduction of particle in the nano-size range.

7.3.2 Challenge 5

The surface of any object exposed to seawater rapidly becomes coated with a variety of inorganic and biological coatings. This mechanism, utilizing natural materials such as timber as a vector for the transfer of organisms, has been taking place for millions of years. However, the rapid increase in floating plastics, which do not disintegrate in transit, has the potential to bring about a rapid increase in the importance of this vector. Colonization of plastic objects by larger sessile organisms is observed frequently. There is emerging concern that microplastics may also act as a vector for microorganisms, including pathogenic species of bacteria, resulting in an increase in the occurrence of non-indigenous species (NIS).²²

Recommendation 5: evaluate the potential significance of plastics and microplastics as a vector for organisms in future assessments

Suggested solution/response:

Review the published evidence on NIS introductions and potential vectors (e.g. ship hull transfer, ballast water transfer), to estimate the relative importance of plastics and microplastics as a transport vector for macro- and micro-organisms. Assess potential consequences in terms of biodiversity and ecosystem functioning. Undertake a targeted risk assessment based on existing data on NIS introductions, and utilize existing circulation models to identify key transport routes, and the conditions favourable for growth, including for pathogenic and invasive organisms.

Risk of not addressing this challenge:

We will be unsure whether plastics and microplastics do represent a significant risk for the introduction of NIS, including pathogenic microorganisms. This may have serious consequences for both ecosystem and potentially human health.

7.3.3 Challenge 6

There is emerging evidence that some organic compounds present as additives, in relatively high concentrations in some categories of plastic, are capable of transferring into the body tissue of fish-eating sea birds. This has been demonstrated for PBDE²³ flame-retardants using a chemical fingerprinting technique, which can distinguish the composition of PBDEs in the plastic particles and the tissue of prey species (Tanaka et al. 2013). The extent to which other chemicals are bioavailable to species of interest is unknown. The role of digestive fluids in influencing transfer rates (e.g. utilizing fish oil for digestion in birds) is also unclear. The extent to which this poses a risk to individual organisms, or to the population as a whole, and to predators of the affected species, including humans, remains untested.

Recommendation 6: future assessments should address the chemical risk posed by ingested microplastics in greater depth

Suggested solution/response:

Compare information from laboratory-based experiments of the bioavailability of the target chemicals with field-based observations of their distribution in the tissue of marine organisms. Include expertise on animal behaviour and physiology for target species, including important commercial species. Take account of gut retention times and the gut environment when assessing risk. Include a consideration of particle size and shape when assessing risk of damage.

Risk of not addressing this challenge:

We will remain unsure as to whether microplastics do pose a significant additional risk to organisms in terms of chemical composition, particle shape and size.

²² Non-indigenous species are sometimes referred to as 'alien' species. Where NIS become established and out compete native species they may be referred to as 'invasive' species.

²³ PBDE – Polybrominated diphenyl ethers, chemicals classified according to the number of bromine rings, can act as endocrine disruptors, affecting fertility.

REFERENCES

- Abu-Hilal, A. H. & Al-Najjar, T. H. (2009) 'Plastic pellets on the beaches of the northern Gulf of Aqaba, Red Sea'. *Aquatic Ecosystem Health & Management*, 12 (4), pp 461-470.
- Adams, R. G., Lohmann, R., Fernandez, L. A. & MacFarlane, J.K., 2007. Polyethylene Devices: Passive Samplers for Measuring Dissolved Hydrophobic Organic Compounds in Aquatic Environments'. *Environmental Science & Technology*, 41 (4), pp 1317-1323.
- Ainley, D., Fraser, W. & Spear, L. (1990) 'The incidence of plastic in the diets of Antarctic seabirds' in Shomura, R. & Godfrey, M. (Eds.). Proceedings of the Second International Conference on Marine Debris. 1990 U.S. Department of Commerce, NOAA Technical Memorandum. NMFS. NOAA-TM-NMFS-SWFSC-154, pp. 653-664.
- Aliani, S., A. Griffo and A. Molcard (2003). "Floating debris in the Ligurian Sea, north-western Mediterranean." *Marine Pollution Bulletin* 46(9): 1142-1149.
- Amos, A., Balazs, G. (1985) Impact of ocean debris on marine turtles: entanglement and ingestion in Shomura, R., Yoshida, H. (Eds.), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 27-29 November 1984. Department of Commerce (1985), pp. 387-429, NOAA Technical Memorandum NMFS SWFC-54., Honolulu, US.
- Anderson, A. & Peterson, A. (2013) 'Nanotechnologies and Trust' in Candlin, C.N. and Crichton, J. (eds.) Discourses of trust. Basingstoke: Palgrave Macmillan, pp. 237-251.
- Anderson, J. A. & Alford, A. B. (2014) 'Ghost fishing activity in derelict blue crab traps in Louisiana'. *Marine Pollution Bulletin*, 79 (1), pp. 261-267.
- Andrady, A. L., H. S. Hamid and A. Torikai (2003). 'Effects of climate change and UV-B on materials'. *Photochemical and Photobiological Sciences* 2(1): 68-72.
- Andrady, A. L. and M. A. Neal (2009). 'Applications and societal benefits of plastics'. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364(1526): 1977-1984.
- Andrady, A. L., J. E. Pegram and N. D. Searle (1996). 'Wavelength sensitivity of enhanced photodegradable polyethylenes, ECO, and LDPE/MX.' *Journal of Applied Polymer Science* 62(9): 1457-1463.
- Andrady, A. L. (2011) 'Microplastics in the marine environment'. *Marine Pollution Bulletin*, 62(8). pp 1596-1605.
- Antunes, J., Frias, J., Micaelo, A. & Sobral, P. (2013) 'Resin pellets from beaches of the Portuguese coast and adsorbed persistent organic pollutants'. *Estuarine, Coastal and Shelf Science*, 130 pp 62-69.
- Arnot, J. A. & Gobas, F. A. P. C. (2004) 'A food web bioaccumulation model for organic chemicals in aquatic ecosystems'. *Environmental Toxicology and Chemistry*, 23 (10). pp 2343-2355.
- Artham, T., Sudhakar, M., Venkatesan, R., Madhavan Nair, C., Murty, K. V. G. K. & Doble, M. (2009) 'Biofouling and stability of synthetic polymers in sea water'. *International Biodeterioration & Biodegradation*, 63 (7). pp 884-890.
- Arthur, C. & Baker, J. (2012) *Proceedings of the Second Research Workshop on Microplastic Marine Debris*. U.S. Department of Commerce, Silver Spring, MD: National Oceanic and Atmospheric Administration. (NOS-OR&R-39).
- Arthur, C., Bamford, H. & Baker, J. (2009) 'Proceedings of the International Research Workshop on the Occurrence, Effects and Fate of Microplastic Marine Debris. Sept 9-11, 2008'. NOAA Technical Memorandum NOS-OR&R-30.
- Arthur, C. D., Opfer, S. E. & Bamford, H. A. (2009) *Standardization of field methods for collection of microplastic marine debris: pelagic, shoreline and sediment protocols*. New Orleans, USA, 19-23 November 2009: Setac 30th North America Meeting. Available.
- Ashton, K., Holmes, L. & Turner, A. (2010) 'Association of metals with plastic production pellets in the marine environment'. *Marine Pollution Bulletin*, 60 (11). pp 2050-2055.
- Azzarello, M. Y. & Van Vleet, E. S. (1987) 'Marine birds and plastic pollution'. *Marine Ecology Progress Series*, 37 pp 295-303.
- Bakir, A., Rowland, S. J. & Thompson, R. C. (2012) 'Competitive sorption of persistent organic pollutants onto microplastics in the marine environment'. *Marine Pollution Bulletin*, 64 (12). pp 2782-2789.
- Bakir, A., Rowland, S. J. & Thompson, R. C. (2014a) 'Enhanced desorption of persistent organic pollutants from microplastics under simulated physiological conditions'. *Environmental Pollution*, 185 (0). pp 16-23.
- Bakir, A., Rowland, S. J. & Thompson, R. C. (2014b) 'Transport of persistent organic pollutants by microplastics in estuarine conditions'. *Estuarine, Coastal and Shelf Science*, 140 pp 14-21.

- Balazs, G. (1985) 'Impact of ocean debris on marine turtles: entanglement and ingestion', Shomura, R. and Yoshida, H. eds.). *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 27–29 November 1984. Honolulu, US Department of Commerce (1985), pp. 387–429, NOAA Technical Memorandum NMFS SWFC-54.
- Ballance, A., Ryan, P. G. & Turpie, J. K. (2000) 'How much is a clean beach worth? The impact of litter on beach users in the Cape Peninsula, South Africa'. *South African Journal of Science*, 96 (5). pp 210-213.
- Ballent, A., Pando, S., Purser, A., Juliano, M. & Thomsen, L. (2013) 'Modelled transport of benthic marine microplastic pollution in the Nazaré Canyon'. *Biogeosciences*, 10 (12). pp 7957-7970.
- Ballent, A., Purser, A., Mendes, P., Pando, S. & Thomsen, L. (2012) 'Physical transport properties of marine microplastic pollution'. *Biogeosciences Discussions*, 9 pp 18755-18798.
- Baltz, D. M. & Morejohn, G. V. (1976) 'Evidence from seabirds of plastic particle pollution off central California'. *Western Birds*, 7 pp 111-112.
- Barnes, D. K., Galgani, F., Thompson, R. C. & Barlaz, M. (2009) 'Accumulation and fragmentation of plastic debris in global environments'. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364 (1526). pp 1985-1998.
- Barnes, D. K. A. (2002) 'Invasions by marine life on plastic debris'. *Nature*, 416 pp 808-809.
- Barnes, D. K. A. & Milner, P. (2005) 'Drifting plastic and its consequences for sessile organism dispersal in the Atlantic Ocean'. *Marine Biology*, 146 (4). pp 815-825.
- Barry, B. & Smith, J. (2014) 'Microplastics in the Mohawk-Hudson watershed', Cockburn, J.M.H. and Garver, J.I. (eds.). *Proceedings of the 2014 Mohawk Watershed Symposium*. Union College, Schenectady, NY, March 21, 2014.
- Bauer, L. J., M. S. Kendall and C. F. G. Jeffrey (2008). 'Incidence of marine debris and its relationships with benthic features in Gray's Reef National Marine Sanctuary, Southeast USA'. *Marine Pollution Bulletin* 56(3): 402-413.
- Baun, A., Hartmann, N. B., Grieger, K. & Kusk, K. O. (2008) 'Ecotoxicity of engineered nanoparticles to aquatic invertebrates: a brief review and recommendations for future toxicity testing'. *Ecotoxicology*, 17 (5). pp 387-395.
- Baztan, J., Carrasco, A., Chouinard, O., Cleaud, M., Gabaldon, J. E., Huck, T., Jaffrès, L., Jorgensen, B., Miguelez, A., Paillard, C. & Vanderlinden, J.-P. (2014) 'Protected areas in the Atlantic facing the hazards of microplastic pollution: First diagnosis of three islands in the Canary Current'. *Marine Pollution Bulletin*, 80 pp 302-311.
- Beiersdorf (2014) 'Raw materials: Making the most of raw materials'. Retrieved August 5, 2014, <http://www.beiersdorf.com/sustainability/products/raw-materials> [Online].
- Bern, L. (1990). 'Size-Related Discrimination Of Nutritive And Inert Particles By Fresh-Water Zooplankton'. *Journal of Plankton Research* 12(5): 1059-1067.
- Berntsen, P., C. Y. Park, B. Rothen-Rutishauser, A. Tsuda, T. M. Sager, R. M. Molina, T. C. Donaghey, A. M. Alencar, D. I. Kasahara, T. Ericsson, E. J. Millet, J. Swenson, D. J. Tschumperlin, J. P. Butler, J. D. Brain, J. J. Fredberg, P. Gehr and E. H. Zhou (2010). 'Biomechanical effects of environmental and engineered particles on human airway smooth muscle cells'. *Journal of the Royal Society Interface* 7: S331-S340.
- Besseling, E., Wegner, A., Foekema, E. M., van den Heuvel-Greve, M. J. & Koelmans, A. A. (2012) 'Effects of Microplastic on Fitness and PCB Bioaccumulation by the Lugworm *Arenicola marina* (L.)'. *Environmental Science & Technology*, 47 (1). pp 593-600.
- Betts, K. (2008) 'Why small plastic particles may pose a big problem in the oceans'. *Environmental Science & Technology*, 42 (24). pp 8995-8995.
- Bhattacharya, P., Lin, S. J., Turner, J. P. & Ke, P. C. (2010) 'Physical Adsorption of Charged Plastic Nanoparticles Affects Algal Photosynthesis'. *Journal of Physical Chemistry C*, 114 (39). pp 16556- 16561.
- Blight, L. K. & Burger, A. E. (1997) 'Occurrence of plastic particles in seabirds from the eastern North Pacific'. *Marine Pollution Bulletin*, 34 (5). pp 323-325.
- Boerger, C. M., Lattin, G. L., Moore, S. L. & Moore, C. J. (2010) 'Plastic ingestion by planktivorous fishes in the North Pacific Central Gyre'. *Marine Pollution Bulletin*, 60 (12). pp 2275-2278.
- Bolton, T. F. & Havenhand, J. N. (1998) 'Physiological versus viscosity-induced effects of an acute reduction in water temperature on microsphere ingestion by trochophore larvae of the serpulid polychaete *Galeolaria caespitosa*'. *Journal of Plankton Research*, 20 (11). pp 2153-2164.
- Bond, A. L., Provencher, J. F., Elliot, R. D., Ryan, P. C., Rowe, S., Jones, I. L., Robertson, G. J. & Wilhelm, S. I. (2013) 'Ingestion of plastic marine debris by Common and Thick-billed Murres in the northwestern Atlantic from 1985 to 2012'. *Marine Pollution Bulletin*, 77 (1–2). pp 192-195.

- Bond, S. (1971) 'Red phalarope mortality in southern California'. *California Birds*, 2 pp 97.
- Bonney, R., Cooper, C. B., Dickinson, J., Kelling, S., Phillips, T., Rosenberg, K. V. & Shirk, J. (2009) 'Citizen science: a developing tool for expanding science knowledge and scientific literacy'. *BioScience*, 59 (11). pp 977-984.
- Bourne, W. & Imber, M. (1980) 'Plastic pellets collected by a prion on Gough Island, central south Atlantic Ocean'. *Marine Pollution Bulletin*, 13 pp 20-21.
- Brandão, M. L., Braga, K. M. & Luque, J. L. (2011) 'Marine debris ingestion by Magellanic penguins, *Spheniscus magellanicus* (Aves: Sphenisciformes), from the Brazilian coastal zone'. *Marine Pollution Bulletin*, 62 (10). pp 2246-2249.
- Bravo, M., de los Angeles Gallardo, M., Luna-Jorquera, G., Núñez, P., Vásquez, N. & Thiel, M. (2009) 'Anthropogenic debris on beaches in the SE Pacific (Chile): Results from a national survey supported by volunteers'. *Marine Pollution Bulletin*, 58 (11). pp 1718-1726.
- Brilliant, M. & MacDonald, B. (2002) 'Postingestive selection in the sea scallop (*Placopecten magellanicus*) on the basis of chemical properties of particles'. *Marine Biology*, 141 (3). pp 457-465.
- Brown, D. M., M. R. Wilson, W. MacNee, V. Stone and K. Donaldson (2001). 'Size-dependent proinflammatory effects of ultrafine polystyrene particles: A role for surface area and oxidative stress in the enhanced activity of ultrafines.' *Toxicology and Applied Pharmacology* 175 (3): 191-199.
- Brown, R. G. B., Barker, S. P., Gaskin, D. E. & Sandeman, M. R. (1981) 'The foods of Great and Sooty Shearwaters *Puffinus gravis* and *P. griseus* in Eastern Canadian waters'. *Ibis*, 123 (1). pp 19-30.
- Brown, V.A., J. Harris and J. Russell (2010). *Tackling Wicked Problems : Through the Transdisciplinary Imagination*. Hoboken, Taylor and Francis.
- Browne, M. A. (2007) *Environmental and biological consequences of microplastic within marine habitats*. Plymouth University.
- Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T. & Thompson, R. (2011) 'Accumulation of Microplastic on Shorelines Worldwide: Sources and Sinks'. *Environmental Science & Technology*, 45 (21). pp 9175-9179.
- Browne, M. A., Dissanayake, A., Galloway, T. S., Lowe, D. M. & Thompson, R. C. (2008) 'Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.)'. *Environmental Science & Technology*, 42 (13). pp 5026-5031.
- Browne, M. A., Galloway, T. S. & Thompson, R. C. (2010) 'Spatial Patterns of Plastic Debris along Estuarine Shorelines'. *Environmental Science & Technology*, 44 (9). pp 3404-3409.
- Browne, M. A., Niven, S. J., Galloway, T. S., Rowland, S. J. & Thompson, R. C. (2013) 'Microplastic Moves Pollutants and Additives to Worms, Reducing Functions Linked to Health and Biodiversity'. *Current Biology*, 23 (23). pp 2388-2392.
- Buckley, P., Pinnegar, J., Terry, G., Chilvers, J., Lorenzoni, I., Gelcich, S., Dudek, A. & Arquati, A. (2011) 'Report on European public awareness and perception of marine climate change risks and impacts'. CLAMER: Climate Change and European Marine Ecosystem Research. *Centre for Environment, Fisheries & Aquaculture Science (Cefas), Lowestoft, Suffolk*.
- Buckley, P. and Pinnegar, J.K. (2011) European public awareness and perception of marine climate risks and impacts. Deliverable Report 2.2. European Seventh Framework Programme, Climate Change and European Marine Ecosystem Research (CLAMER), <http://www.clamer.eu> 64pp.
- Canesi, L., C. Ciacci, R. Fabbri, A. Marcomini, G. Pojana and G. Gallo (2012). 'Bivalve molluscs as a unique target group for nanoparticle toxicity'. *Marine Environmental Research* 76: 16-21.
- Carey, M. J. (2011) 'Intergenerational transfer of plastic debris by Short-tailed Shearwaters (*Ardenna tenuirostris*)'. *Emu*, 111 (3). pp 229-234.
- Carpenter, E. J., Anderson, S. J., Harvey, G. R., Miklas, H. P. & Peck, B. B. (1972) 'Polystyrene spherules in coastal waters'. *Science*, 178 (4062). pp 749-750.
- Carpenter, E. J. & Smith, K. (1972) 'Plastics on the Sargasso Sea surface'. *Science*, 175 (4027). pp 1240-1241.
- Carson, H. S., Colbert, S. L., Kaylor, M. J. & McDermid, K. J. (2011) 'Small plastic debris changes water movement and heat transfer through beach sediments'. *Marine Pollution Bulletin*, 62 (8). pp 1708-1713.
- Carson, H. S., Nerheim, M. S., Carroll, K. A. & Eriksen, M. (2013) 'The plastic-associated microorganisms of the North Pacific Gyre'. *Marine Pollution Bulletin*, 75 (1-2). pp 126-132.
- Cedervall, T., L.-A. Hansson, M. Lard, B. Frohm and S. Linse (2012). 'Food Chain Transport of Nanoparticles Affects Behaviour and Fat Metabolism in Fish'. *PLoS ONE*, 7(2).

- Celina, M. C. (2013). 'Review of polymer oxidation and its relationship with materials performance and lifetime prediction'. *Polymer Degradation and Stability* 98(12): 2419-2429.
- Chan, W. Y. & Witting, J. (2012) 'The Impact of Microplastics on Salp Feeding in the Tropical Pacific'. *The ANU Undergraduate Research Journal*, pp 129.
- Cheng, L. & Pitman, R. L. (2002) 'Mass oviposition and egg development of the ocean-skater *Halobates sobrinus* (Heteroptera: Gerridae)'. *Pacific science*, 56 (4). pp 441-447.
- Choy, C. A. & Drazen, J. C. (2013) 'Plastic for dinner? Observations of frequent debris ingestion by pelagic predatory fishes from the central North Pacific'. *Marine Ecology Progress Series*, 485. pp 155-163.
- Chua, E. M., J. Shimeta, D. Nugagoda, P. D. Morrison and B. O. Clarke (2014). 'Assimilation of Polybrominated Diphenyl Ethers from Microplastics by the Marine Amphipod, *Allorchestes compressa*'. *Environmental Science & Technology* 48(14): 8127-8134.
- Claessens, M., De Meester, S. & Janssen, C. R. (2009) 'Occurrence of microplastics in the Belgian coastal zone'. Belgium, 27 November 2009. VLIZ Special Publication, 43. Vlaams Instituut voor de Zee (VLIZ): Oostende, Belgium. xiii + 221p.
- Claessens, M., De Meester, S., Van Landuyt, L., De Clerck, K. & Janssen, C. R. (2011) 'Occurrence and distribution of microplastics in marine sediments along the Belgian coast'. *Marine Pollution Bulletin*, 62 (10). pp 2199-2204.
- Claessens, M. & Janssen, C. R. (2011) 'Plastics on your plate? Detecting microplastics in sediments and organisms', Mees, J. and Seys, J. (eds.). Book of abstracts. VLIZ Young Scientist's Day. Bruges, Belgium, 25 February 2011. VLIZ Special Publication, 48. Vlaams Insituut voor de Zee (VLIZ): Oostende, Belgium. xiv + 143p.
- Claessens, M., Van Cauwenberghe, L., Goffin, A., Dewitte, E., Braarup Cuykens, A., Maelfait, H., Vanhecke, V., Mees, J., Stienen, E. & Janssen, C. (2013a) *Assessment of marine debris on the Belgian continental shelf: occurrence and effects' AS-MADE'*: final report. Brussels, Belgium: Belgian Science Policy Office. 79 p pp. Available.
- Claessens, M., Van Cauwenberghe, L., Vandegehuchte, M. B. & Janssen, C. R. (2013b) 'New techniques for the detection of microplastics in sediments and field collected organisms'. *Marine Pollution Bulletin*, 70 (1-2). pp 227-233.
- Codina-García, M., Militão, T., Moreno, J. & González-Solís, J. (2013) 'Plastic debris in Mediterranean seabirds'. *Marine Pollution Bulletin*, 77 (1-2). pp 220-226.
- Cohn, J. P. (2008) 'Citizen science: Can volunteers do real research?'. *BioScience*, 58 (3). pp 192- 197.
- Colabuono, F. I., Barquete, V., Domingues, B. S. & Montone, R. C. (2009) 'Plastic ingestion by Procellariiformes in southern Brazil'. *Marine Pollution Bulletin*, 58 (1). pp 93-96.
- Colabuono, F. I., Taniguchi, S. & Montone, R. C. (2010) 'Polychlorinated biphenyls and organochlorine pesticides in plastics ingested by seabirds'. *Marine Pollution Bulletin*, 60 (4). pp 630-634.
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J. & Galloway, T. S. (2013) 'Microplastic Ingestion by Zooplankton'. *Environmental Science & Technology*, 47 (12). pp 6646- 6655.
- Cole, M., Lindeque, P., Halsband, C. & Galloway, T. S. (2011) 'Microplastics as contaminants in the marine environment: A review'. *Marine Pollution Bulletin*, 62 (12). pp 2588-2597.
- Cole, M., Webb, H., Lindeque, P. K., Fileman, E. S., Halsband, C. & Galloway, T. S. (2014) 'Isolation of microplastics in biota-rich seawater samples and marine organisms'. *Scientific Reports*, 4 pp 4528.
- Collignon, A., Hecq, J.-H., Galgani, F., Collard, F. & Goffart, A. (2014) 'Annual variation in neustonic micro- and meso-plastic particles and zooplankton in the Bay of Calvi (Mediterranean-Corsica)'. *Marine Pollution Bulletin*, 79. pp 293-298.
- Collignon, A., Hecq, J.-H., Galgani, F., Voisin, P., Collard, F. & Goffart, A. (2012) 'Neustonic microplastic and zooplankton in the North Western Mediterranean Sea'. *Marine Pollution Bulletin*, 64 (4). pp 861-864.
- Colton, J. B., Knapp, F. D. & Burns, B. R. (1974) 'Plastic particles in surface waters of the northwestern Atlantic'. *Science*, 185 (4150). pp 491-497.
- Connors, P. G. & Smith, K. G. (1982) 'Oceanic plastic particle pollution: Suspected effect on fat deposition in red phalaropes'. *Marine Pollution Bulletin*, 13 (1). pp 18-20.
- Cooper, D. A. & Corcoran, P. L. (2010) 'Effects of mechanical and chemical processes on the degradation of plastic beach debris on the island of Kauai, Hawaii'. *Marine Pollution Bulletin*, 60 (5). pp 650-654.
- Cooper, P. (2013). Socio-ecological accounting: DPSWR, a modified DPSIR framework, and its application to marine ecosystems. *Ecological Economics* 94: 106-115.

- Corcoran, P. L., Biesinger, M. C. & Grifi, M. (2009) 'Plastics and beaches: A degrading relationship'. *Marine Pollution Bulletin*, 58 (1). pp 80-84.
- Cornelissen, G., Pettersen, A., Broman, D., Mayer, P. & Breedveld, G. D. (2008) 'Field testing of equilibrium passive samplers to determine freely dissolved native polycyclic aromatic hydrocarbon concentrations'. *Environmental Toxicology and Chemistry*, 27 (3). pp 499-508.
- Costa, M. F., do Sul, J. A. I., Silva-Cavalcanti, J. S., Araújo, M. C. B., Spengler, Â. & Tourinho, P. S. (2010) 'On the importance of size of plastic fragments and pellets on the strandline: a snapshot of a Brazilian beach'. *Environmental Monitoring and Assessment*, 168 (1-4). pp 299-304.
- Cózar, A., F. Echevarría, J. I. González-Gordillo, X. Irigoien, B. Úbeda, S. Hernández-León, Á. T. Palma, S. Navarro, J. García-de-Lomas, A. Ruiz, M. L. Fernández-de-Puelles and C. M. Duarte (2014). 'Plastic debris in the open ocean'. *Proceedings of the National Academy of Sciences*, 111 10239-10244.
- Creet, S., van Franeker, J., van Spanje, T. & Wolff, W. (1994) 'Diet of the pintado petrel *Daption capense* at King George Island, Antarctica, 1990/91'. *Marine Ornithology*, 22 (2). pp 221-229.
- Cruz-Pinto, J. J. C., Carvalho, M. E. S. and Ferreira, J. F. A. (1994). The kinetics and mechanism of polyethylene photo-oxidation. *Angew Makromol Chem.* 216:113-33.
- Davidson, T. M. (2012) 'Boring crustaceans damage polystyrene floats under docks polluting marine waters with micro-plastic'. *Marine Pollution Bulletin*, 64 (9). pp 1821-1828.
- Davis, J. L., Green, J. D. & Reed, A. (2009) 'Interdependence with the environment: Commitment, interconnectedness, and environmental behavior'. *Journal of Environmental Psychology*, 29 (2). pp 173-180.
- Davison, P. & Asch, R. G. (2011) 'Plastic ingestion by mesopelagic fishes in the North Pacific Subtropical Gyre'. *Marine Ecology Progress Series*, 432 pp 173-180.
- Day, R.H., Wehle, D.H.S., Coleman, F. (1985). Ingestion of plastic pollutants by marine birds, in: Shomura, R., Yoshida, H. (Eds.), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 27-29 November 1984 Department of Commerce (1985), pp. 387-429, NOAA Technical Memorandum NMFS SWFC-54, Honolulu, US.
- Day, R. H. & Shaw, D. G. (1987) 'Patterns in the abundance of pelagic plastic and tar in the north Pacific Ocean, 1976-1985'. *Marine Pollution Bulletin*, 18 (6, Supplement B). pp 311-316.
- De Jong, W. H. & Borm, P. J. (2008) 'Drug delivery and nanoparticles: applications and hazards'. *International Journal of Nanomedicine*, 3 (2). pp 133.
- De Jorge, J. 'El «séptimo continente»: un basurero flotante en el Pacífico. Ciencia, ABC.es '. Retrieved August 19, 2014 <http://www.abc.es/20120416/ciencia/abci-septimo-continente-basurero-flotante-201204161033.html>.
- de Lucia, G. A., Caliani, I., Marra, S., Camedda, A., Coppa, S., Alcaro, L., Campani, T., Giannetti, M., Coppola, D., Cicero, A. M., Panti, C., Baini, M., Guerranti, C., Marsili, L., Massaro, G., Fossi, M. C. & Matiddi, M. (In Press) 'Amount and distribution of neustonic micro-plastic off the western Sardinian coast (Central-Western Mediterranean Sea)'. *Marine Environmental Research*, (0).
- de Stephanis, R., J. Gimenez, E. Carpinelli, C. Gutierrez-Exposito and A. Canadas (2013). 'As main meal for sperm whales: Plastics debris'. *Marine Pollution Bulletin*, 69(1-2): 206-214.
- De Witte, B., Devriese, L., Bekaert, K., Hoffman, S., Vandermeersch, G., Cooreman, K. & Robbens, J. (2014) 'Quality assessment of the blue mussel (*Mytilus edulis*): Comparison between commercial and wild types'. *Marine Pollution Bulletin*, 85 (1). pp 146-155.
- Dekiff, J. H., Remy, D., Klasmeier, J. & Fries, E. (2014) 'Occurrence and spatial distribution of microplastics in sediments from Norderney'. *Environmental Pollution*, 186 (0). pp 248-256.
- Den Hartog, J. & Van Nierop, M. (1984) *A study on the gut contents of six leathery turtles Dermochelys coriacea (Linnaeus) (Reptilia: Testudines: Dermochelyidae) from British waters and from the Netherlands*. Rijksmuseum van Natuurlijke Historie.
- Depledge, M., Galgani, F., Panti, C., Caliani, I., Casini, S. & Fossi, M. (2013) 'Plastic litter in the sea'. *Marine Environmental Research*, 92 pp 279-281.
- Denuncio, P., R. Bastida, M. Dassis, G. Giardino, M. Gerpe and D. Rodriguez (2011). 'Plastic ingestion in Franciscana dolphins, *Pontoporia blainvillei* (Gervais and d'Orbigny, 1844), from Argentina'. *Marine Pollution Bulletin* 62(8): 1836-1841.
- Derraik, J. G. (2002) 'The pollution of the marine environment by plastic debris: a review'. *Marine Pollution Bulletin*, 44 (9). pp 842-852.

- Desforbes, J.-P. W., Galbraith, M., Dangerfield, N. & Ross, P. S. (2014) 'Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean'. *Marine Pollution Bulletin*, 79. pp 94-99.
- Detloff, K. C. (2012) 'Marine Litter: Projects–Threats–Solutions'. *Progress in Marine Conservation in Europe 2012*, 18 pp 135.
- Donahue, S. C. (1996) 'Clean ships, clean ports, clean oceans: Controlling garbage and plastic wastes at sea : Edited by the Committee on Shipborne Wastes, Marine Board, Commission on Engineering and Technical Systems, National Research Council. National Academy Press'. *Ocean & Coastal Management*, 32 (2). pp 127-128.
- Dori, Y. J. & Tal, R. T. (2000) 'Formal and informal collaborative projects: Engaging in industry with environmental awareness'. *Science Education*, 84 (1). pp 95-113.
- Doughty, R. & Eriksen, M. (2014) 'Plastic Pollution: The Case for a Ban on Microplastics in Personal Care Products'. *Tul. Env'tl. LJ*, 27. pp 277-393.
- Doyle, M. J., Watson, W., Bowlin, N. M. & Sheavly, S. B. (2011) 'Plastic particles in coastal pelagic ecosystems of the Northeast Pacific ocean'. *Marine Environmental Research*, 71 (1). pp 41-52.
- Dubaish, F. & Liebezeit, G. (2013) 'Suspended Microplastics and Black Carbon Particles in the Jade System, Southern North Sea'. *Water Air and Soil Pollution*, 224 (2). pp 1352.
- Dufault, S. & Whitehead, H. (1994) 'Floating marine pollution in 'the Gully' on the continental slope, Nova Scotia, Canada'. *Marine Pollution Bulletin*, 28 (8). pp 489-493.
- Duffus, J. H., Nordberg, M. & Templeton, D. M. (2007) 'Glossary of terms used in toxicology, (IUPAC recommendations 2007)'. *Pure and Applied Chemistry*, 79 (7). pp 1153-1344.
- Dunlap, R. E. and A. M. McCright (2008). 'A Widening Gap: Republican and Democratic Views on Climate Change'. *Environment: Science and Policy for Sustainable Development* 50(5): 26-35.
- Eastman, L. B., Núñez, P., Crettier, B. & Thiel, M. (2013) 'Identification of self-reported user behavior, education level, and preferences to reduce littering on beaches—A survey from the SE Pacific'. *Ocean & Coastal Management*, 78. pp 18-24.
- Edwards, M., Beaugrand, G., Johns, D.G., Helaouet, P., Licandro, P., McQuatters-Gollop, A., Wootton, M. (2011) *Ecological Status Report: results from the CPR survey 2009/2010*. Plymouth, UK: SAHFOS Technical Report. Available.
- Endo, S., Takizawa, R., Okuda, K., Takada, H., Chiba, K., Kanehiro, H., Ogi, H., Yamashita, R. & Date, T. (2005) 'Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets: Variability among individual particles and regional differences'. *Marine Pollution Bulletin*, 50 (10). pp 1103-1114.
- Endo, S., Yuyama, M. & Takada, H. (2013) 'Desorption kinetics of hydrophobic organic contaminants from marine plastic pellets'. *Marine Pollution Bulletin*, 74 (1). pp 125-131.
- Eriksen, M., L. C. M. Lebreton, H. S. Carson, M. Thiel, C. J. Moore, J. C. Borerro, F. Galgani, P. G. Ryan and J. Reisser (2014). 'Plastic Pollution in the World's Oceans: More than 5 Trillion Plastic Pieces Weighing over 250,000 Tons Afloat at Sea'. *PLoS ONE*, 9(12).
- Eriksen, M., Mason, S., Wilson, S., Box, C., Zellers, A., Edwards, W., Farley, H. & Amato, S. (2013a) 'Microplastic pollution in the surface waters of the Laurentian Great Lakes'. *Marine Pollution Bulletin*, 77 (1-2). pp 177-182.
- Eriksen, M., Maximenko, N., Thiel, M., Cummins, A., Lattin, G., Wilson, S., Hafner, J., Zellers, A. & Rifman, S. (2013b) 'Plastic pollution in the South Pacific subtropical gyre'. *Marine Pollution Bulletin*, 68 (1-2). pp 71-76.
- Eriksson, C. & Burton, H. (2003) 'Origins and Biological Accumulation of Small Plastic Particles in Fur Seals from Macquarie Island'. *Ambio*, 32 (6). pp 380-384.
- Eubeler, J. P., S. Zok, M. Bernhard and T. P. Knepper (2009). 'Environmental biodegradation of synthetic polymers I. Test methodologies and procedures'. *TrAC Trends in Analytical Chemistry* 28(9): 1057-1072.
- European Commission (2014) 'Flash Eurobarometer 388: Attitudes of Europeans towards Waste Management and Resource Efficiency'. Retrieved July 8, 2014 from http://ec.europa.eu/public_opinion/flash/fl_388_en.pdf.
- Farrell, P. & Nelson, K. (2013) 'Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus maenas* (L.)'. *Environmental Pollution*, 177 (0). pp 1-3.
- Farrington, J. W. & Takada, H. (2014) 'Persistent Organic Pollutants (POPs), Polycyclic Aromatic Hydrocarbons (PAHs), and Plastics Examples of the Status, Trend, and Cycling of Organic Chemicals of Environmental Concern in the Ocean'. *Oceanography*, 27 (1). pp 196-213.
- Felsten, G. (2009) 'Where to take a study break on the college campus: An attention restoration theory perspective'. *Journal of Environmental Psychology*, 29 (1). pp 160-167.

- Fendall, L. S. & Sewell, M. A. (2009) 'Contributing to marine pollution by washing your face: Microplastics in facial cleansers'. *Marine Pollution Bulletin*, 58 (8). pp 1225-1228.
- Finucane, M. L., Slovic, P., Mertz, C. K., Flynn, J. & Satterfield, T. A. (2000) 'Gender, race, and perceived risk: The 'white male' effect'. *Health, risk & society*, 2 (2). pp 159-172.
- Fischer, V., Elsner, N.O., Brenke, N., Schwabe, E. and Brandt, A. (2015). Plastic pollution of the Kuril–Kamchatka Trench area (NW pacific) . *Deep-Sea Research II*, 111, 399-405.
- Fisner, M., Taniguchi, S., Majer, A. P., Bicego, M. C. & Turra, A. (2013a) 'Concentration and composition of polycyclic aromatic hydrocarbons (PAHs) in plastic pellets: Implications for small-scale diagnostic and environmental monitoring'. *Marine Pollution Bulletin*, 76 (1-2). pp 349-354.
- Fisner, M., Taniguchi, S., Moreira, F., Bicego, M. C. & Turra, A. (2013b) 'Polycyclic aromatic hydrocarbons (PAHs) in plastic pellets: Variability in the concentration and composition at different sediment depths in a sandy beach'. *Marine Pollution Bulletin*, 70 (1-2). pp 219-226.
- Fleming, L., Broad, K., Clement, A., Dewailly, E., Elmir, S., Knap, A., Pomponi, S., Smith, S., Solo Gabriele, H. & Walsh, P. (2006) 'Oceans and human health: emerging public health risks in the marine environment'. *Marine Pollution Bulletin*, 53 (10). pp 545-560.
- Fletcher, S. & Potts, J. (2007) 'Ocean citizenship: an emergent geographical concept'. *Coastal management*, 35 (4). pp 511-524.
- Fletcher, S., Potts, J. S., Heeps, C. & Pike, K. (2009) 'Public awareness of marine environmental issues in the UK'. *Marine Policy*, 33 (2). pp 370-375.
- Flynn, J., Slovic, P. & Mertz, C. K. (1994) 'Gender, race, and perception of environmental health risks'. *Risk analysis*, 14 (6). pp 1101-1108.
- Foekema, E. M., De Gruijter, C., Mergia, M. T., van Franeker, J. A., Murk, A. J. & Koelmans, A. A. (2013) 'Plastic in North sea fish'. *Environmental Science & Technology*, 47 (15). pp 8818-8824.
- Fossi, M., Panti, C., Coppola, D., Matteo, M., Marsili, L., Minutoli, R. & Guerranti, C. (2012a) 'Are baleen whales exposed to microplastics threat? The case study of the Mediterranean Fin whale'. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 163 pp S25-S26.
- Fossi, M. C., Coppola, D., Baini, M., Giannetti, M., Guerranti, C., Marsili, L., Panti, C., de Sabata, E. & Clò, S. (In Press) 'Large filter feeding marine organisms as indicators of microplastic in the pelagic environment: The case studies of the Mediterranean basking shark (*Cetorhinus maximus*) and fin whale (*Balaenoptera physalus*)'. *Marine Environmental Research*, 100. pp 17-24.
- Fossi, M. C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L. & Minutoli, R. (2012c) 'Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*)'. *Marine Pollution Bulletin*, 64 (11). pp 2374-2379.
- Fotopoulou, K. N. & Karapanagioti, H. K. (2012) 'Surface properties of beached plastic pellets'. *Marine Environmental Research*, 81 (0). pp 70-77.
- Free, C. M., Jensen, O. P., Mason, S. A., Eriksen, M., Williamson, N. J. & Boldgiv, B. 'High-levels of microplastic pollution in a large, remote, mountain lake'. *Marine Pollution Bulletin*, (0).
- Frias, J., Otero, V. & Sobral, P. (2014) 'Evidence of microplastics in samples of zooplankton from Portuguese coastal waters'. *Marine Environmental Research*, 95 pp 89-95.
- Frias, J. P., Antunes, J. C. & Sobral, P. (2013) 'Local marine litter survey-A case study in Alcobaça municipality, Portugal'. *RGCI-Revista de Gestão Costeira Integrada*, 13 (2).
- Frias, J. P. G. L., Sobral, P. & Ferreira, A. M. (2010) 'Organic pollutants in microplastics from two beaches of the Portuguese coast'. *Marine Pollution Bulletin*, 60 (11). pp 1988-1992.
- Fries, E., Dekiff, J. H., Willmeyer, J., Nuelle, M.-T., Ebert, M. & Remy, D. (2013) 'Identification of polymers types and additives in marine microplastic particles using pyrolysis-GC/MS and scanning electron microscopy'. *Environmental Science: Processes & Impacts*, 15 pp 1949-1956.
- Fries, E. & Zarfl, C. (2012) 'Sorption of polycyclic aromatic hydrocarbons (PAHs) to low and high density polyethylene (PE)'. *Environmental Science and Pollution Research*, 19 (4). pp 1296-1304.
- Fröhlich, E., C. Samberger, T. Kueznik, M. Absenger, E. Roblegg, A. Zimmer and T. R. Pieber (2009). 'Cytotoxicity of nanoparticles independent from oxidative stress'. *Journal of Toxicological Sciences* 34(4): 363-375.

- Fry, D. M., Fefer, S. I. & Sileo, L. (1987) 'Ingestion of plastic debris by Laysan Albatrosses and Wedge-tailed Shearwaters in the Hawaiian Islands'. *Marine Pollution Bulletin*, 18 (6, Supplement B). pp 339-343.
- Furness, B. L. (1983) 'Plastic particles in three procellariiform seabirds from the Benguela Current, South Africa'. *Marine Pollution Bulletin*, 14 (8). pp 307-308.
- Furness, R. W. (1985a) 'Ingestion of plastic particles by seabirds at Gough Island, south Atlantic Ocean'. *Environmental Pollution Series A, Ecological and Biological*, 38 (3). pp 261-272.
- Furness, R. W. (1985b) 'Plastic particle pollution: Accumulation by procellariiform seabirds at Scottish Colonies'. *Marine Pollution Bulletin*, 16 (3). pp 103-106.
- Gabrys, J. (2013) '12 Plastic and the work of the biodegradable'. *Accumulation: The Material Politics of Plastic*, pp 208.
- Galgani, F., Claro, F., Depledge, M. & Fossi, C. (In Press) 'Monitoring the impact of litter in large vertebrates in the Mediterranean Sea within the European Marine Strategy Framework Directive (MSFD): Constraints, specificities and recommendations'. *Marine Environmental Research*, (0).
- Galgani, F., Ellerbrake, K., Fries, E. & Goreux, C. (2011) 'Marine pollution: Let us not forget beach sand'. *Environmental Sciences Europe*, 23 (1). pp 1-7.
- Galgani, F., Hanke G, Werner S., and Piha, H. 2011. MSFD GES Technical Subgroup on Marine Litter. Technical Recommendations for the Implementation of MSFD Requirements. JRC Scientific and Technical Report, EUR 25009 EN – 2011. 93 pp.
- Galgani, F., Hanke, G., Werner, S. & De Vrees, L. (2013) 'Marine litter within the European Marine Strategy Framework Directive'. *ICES Journal of Marine Science: Journal du Conseil*, 70 (6). pp 1055-1064.
- Galgani, F., Zampoukas, N., Fleet, D., Franeker, J. v., Katsanevakis, S., Maes, T., Mouat, J., Oosterbaan, L., Poitou, I. & Hanke, G. (2010) Marine Strategy Framework Directive: Task Group 10 Report Marine Litter. Office for Official Publications of the European Communities.
- Garza, S. & Smith, C. A. (2010) 'Analysis of microplastics in Corpus Christi's marine environment'. *Abstracts of Papers of the American Chemical Society*, 240
- Gaudet, J. J. (2012) 'Garbage Patch'e-flows: Exploring on-line plastics in the ocean knowledge politics', *University of Waterloo-Wilfrid Laurier University* 2012.
- Gaylor, Michael O., Ellen Harvey, and Robert C. Hale. 'House crickets can accumulate polybrominated diphenyl ethers (PBDEs) directly from polyurethane foam common in consumer products'. *Chemosphere* 86.5 (2012): 500-505.
- Geiser, M., B. Rothen-Rutishauser, N. Kapp, S. Schurch, W. Kreyling, H. Schulz, M. Semmler, V. I. Hof, J. Heyder and P. Gehr (2005). 'Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells'. *Environmental Health Perspectives* 113(11): 1555-1560.
- George, A. R. (2006) *A New Spectroscopic Method for the Non-Destructive Characterization of Weathering Damage In Plastics*. Brigham Young University.
- GESAMP (2001). Protecting the oceans from land-based activities. IMO, FAO, UNESCO-IOC, UNIDO, WMO, IAEA, UN, UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection, GESAMP Rep. Stud. No. 71, 162pp.
- GESAMP (2010). Bowmer, T. and Kershaw, P.J., 2010 (Eds.) 'Proceedings of the GESAMP International Workshop on micro-plastic particles as a vector in transporting persistent, bio-accumulating and toxic substances in the oceans'. IMO, FAO, UNESCO-IOC, UNIDO, WMO, IAEA, UN, UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection, GESAMP Rep. Stud. No. 82, 68pp.
- Gilfillan, L. R., Doyle, M. J., Ohman, M. D. & Watson, W. (2009) *Occurrence of Plastic Micro-debris in the Southern California Current System*. CalCOFI. Available.
- Goldstein, M. C. (2012) *Abundance and Ecological Implications of Microplastic Debris in the North Pacific Subtropical Gyre*. University of California, San Diego.
- Goldstein, M. C. & Goodwin, D. S. (2013) 'Gooseneck barnacles (*Lepas* spp.) ingest microplastic debris in the North Pacific Subtropical Gyre'. *PeerJ*, 1 pp e184.
- Goldstein, M. C., Rosenberg, M. & Cheng, L. (2012) 'Increased oceanic microplastic debris enhances oviposition in an endemic pelagic insect'. *Biology Letters*, 8 (5). pp 817-820.
- Goldstein, M. C., Titmus, A. J. & Ford, M. (2013) 'Scales of Spatial Heterogeneity of Plastic Marine Debris in the Northeast Pacific Ocean'. *PLoS ONE*, 8 (11). pp e80020.

- Google Trends (2014a) 'About Trends Graphs'. Accessed July 9, 2014 from <https://support.google.com/trends/answer/4355164?hl=en&rd=1>.
- Google Trends (2014b) 'http://www.google.com/trends/'. Accessed July 14, 2014 from <http://www.google.com/trends/>.
- Gorycka, M. (2009) 'Environmental risks of microplastics'. *Investigation Project. Vrije Universiteit. Faculteit der Aard-en Levenswetenschappen. Holland*, Project Report 468017, 171pp.
- Gouin, T., Roche, N., Lohmann, R. & Hodges, G. (2011) 'A Thermodynamic Approach for Assessing the Environmental Exposure of Chemicals Absorbed to Microplastic'. *Environmental Science & Technology*, 45 (4). pp 1466-1472.
- Graham, E. R. & Thompson, J. T. (2009) 'Deposit- and suspension-feeding sea cucumbers (Echinodermata) ingest plastic fragments'. *Journal of Experimental Marine Biology and Ecology*, 368 (1). pp 22-29.
- Gregory, M. R. (1977) 'Plastic pellets on New Zealand beaches'. *Marine Pollution Bulletin*, 8 (4). pp 82-84.
- Gregory, M. R. (1978) 'Accumulation and distribution of virgin plastic granules on New Zealand beaches'. *New Zealand Journal of Marine and Freshwater Research*, 12 (4). pp 399-414.
- Gregory, M. R. (1983) 'Virgin plastic granules on some beaches of eastern Canada and Bermuda'. *Marine Environmental Research*, 10 (2). pp 73-92.
- Gregory, M. R. (1996) 'Plastic "scrubbers" in hand cleansers: a further (and minor) source for marine pollution identified'. *Marine Pollution Bulletin*, 32 (12). pp 867-871.
- Gregory, M. R. (2009) 'Environmental implications of plastic debris in marine settings - entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions'. *Philosophical Transactions of the Royal Society B*, 364 pp 2013- 2025.
- Gregory, M. R. and Andrady, A. L. (2003) *Plastics in the Marine Environment*, in *Plastics and the Environment* (ed. A. L. Andrady), John Wiley & Sons, Inc., Hoboken, NJ, USA. doi: 10.1002/0471721557.ch10
- Hagmann, P., Faure, F., Galgani, F., Potter, G., de Alencastro, L. & Saini, C. (2013) 'An evaluation of surface micro and meso plastic pollution in pelagic ecosystems of western Mediterranean Sea', *6th International Conference on Water Resources and Environment Research (ICWRER)*.
- Halden, R. U. Plastics and health risks. *Annual Review of Public Health*. 31: 179-194.
- Harms, J. (1990) 'Marine plastic litter as an artificial hard bottom fouling ground'. *Helgoländer Meeresuntersuchungen*, 44 (3-4). pp 503-506.
- Harper, P. C. & Fowler, J. A. (1987) 'Plastic pellets in New Zealand storm-killed prions (*Pachyptila* spp.), 1958-1977'. *Notornis*, 34 pp 65-70.
- Harrison, J. P. (2012) *The spectroscopic detection and bacterial colonisation of synthetic microplastics in coastal marine sediments*. PhD thesis, University of Sheffield.
- Harrison, J. P., Ojeda, J. J. & Romero-Gonzalez, M. E. (2012) 'The applicability of reflectance micro-Fourier-transform infrared spectroscopy for the detection of synthetic microplastics in marine sediments'. *Science of The Total Environment*, 416 pp 455-463.
- Harrison, J. P., Sapp, M., Schratzberger, M. & Osborn, A. M. (2011) 'Interactions Between Microorganisms and Marine Microplastics: A Call for Research'. *Marine Technology Society Journal*, 45 (2). pp 12-20.
- Hart, M. W. (1991) 'Particle Captures and the Method of Suspension Feeding by Echinoderm Larvae'. *The Biological Bulletin*, 180 (1). pp 12-27.
- Hays, H. & Cormons, G. (1974) 'Plastic particles found in tern pellets, on coastal beaches and at factory sites'. *Marine Pollution Bulletin*, 5 (3). pp 44-46.
- Heo, N. W., S. H. Hong, G. M. Han, S. Hong, J. Lee, Y. K. Song, M. Jang and W. J. Shim (2013). 'Distribution of Small Plastic Debris in Cross-section and High Strandline on Heungnam Beach, South Korea'. *Ocean Science Journal* 48(2): 225-233.
- Heskett, M., Takada, H., Yamashita, R., Yuyama, M., Ito, M., Geok, Y. B., Ogata, Y., Kwan, C., Heckhausen, A., Taylor, H., Powell, T., Morishige, C., Young, D., Patterson, H., Robertson, B., Bailey, E. & Mermoz, J. (2012) 'Measurement of persistent organic pollutants (POPs) in plastic resin pellets from remote islands: Toward establishment of background concentrations for International Pellet Watch'. *Marine Pollution Bulletin*, 64 (2). pp 445-448.
- Hesterberg, T. W., C. M. Long, C. A. Lapin, A. K. Hamade and P. A. Valberg (2010). 'Diesel exhaust particulate (DEP) and nanoparticle exposures: What do DEP human clinical studies tell us about potential human health hazards of nanoparticles?' *Inhalation Toxicology* 22(8): 679-694.

- Hidalgo-Ruz, V., Gutow, L., Thompson, R. C. & Thiel, M. (2012) 'Microplastics in the Marine Environment: A Review of the Methods Used for Identification and Quantification'. *Environmental Science & Technology*, 46 (6). pp 3060-3075.
- Hidalgo-Ruz, V. & Thiel, M. (2013) 'Distribution and abundance of small plastic debris on beaches in the SE Pacific (Chile): A study supported by a citizen science project'. *Marine Environmental Research*, 87-88. pp 12-18.
- Hinds, J. & Sparks, P. (2008) 'Engaging with the natural environment: The role of affective connection and identity'. *Journal of Environmental Psychology*, 28 (2). pp 109-120.
- Hipp, J. A. & Ogunseitan, O. A. (2011) 'Effect of environmental conditions on perceived psychological restorativeness of coastal parks'. *Journal of Environmental Psychology*, 31 (4). pp 421-429.
- Hinds, J., & Spark, P. (2008). Engaging with the natural environment: The role of affective connection and identity. *Journal of Environmental Psychology*, 28, 109-120.
- Hirai, H., Takada, H., Ogata, Y., Yamashita, R., Mizukawa, K., Saha, M., Kwan, C., Moore, C., Gray, H., Laursen, D., Zettler, E. R., Farrington, J. W., Reddy, C. M., Peacock, E. E. & Ward, M. W. (2011) 'Organic micropollutants in marine plastics debris from the open ocean and remote and urban beaches'. *Marine Pollution Bulletin*, 62 (8). pp 1683-1692.
- Hodgkin, J. H., Simon, G. P. & Varley, R. J. (1998) 'Thermoplastic Toughening of Epoxy Resins: a Critical Review'. *Polymers for Advanced Technologies*, 9. pp 3-10.
- Höher, N., A. Koehler, J. Strand and K. Broeg (2012). 'Effects of various pollutant mixtures on immune responses of the blue mussel (*Mytilus edulis*) collected at a salinity gradient in Danish coastal waters'. *Marine Environmental Research* 75: 35-44.
- Hoet, P. H., Brüske-Hohlfeld, I. & Salata, O. V. (2004) 'Nanoparticles – known and unknown health risks'. *Journal of nanobiotechnology*, 2 (1). pp 12.
- Hoffmann, J., Reznickova, I., Kozakova, J., Ruzicka, J., Alexy, P., Bakos, D. & Precnerova, L. (2003) 'Assessing biodegradability of plastics based on poly(vinyl alcohol) and protein wastes'. *Polymer Degradation and Stability*, 79 (3). pp 511-519.
- Holmes, L. A. (2013) *Interactions of Trace Metals with Plastic Production Pellets in the Marine Environment*. PhD Thesis, University of Plymouth.
- Holmes, L. A., Turner, A. & Thompson, R. C. (2012) 'Adsorption of trace metals to plastic resin pellets in the marine environment'. *Environmental Pollution*, 160 (0). pp 42-48.
- Hong, S., Lee, J., Jang, Y. C., Kim, Y. J., Kim, H. J., Han, D., Hong, S. H., Kang, D. & Shim, W. J. (2013) 'Impacts of marine debris on wild animals in the coastal area of Korea'. *Marine Pollution Bulletin*, 66 (1). pp 117-124.
- Hong, S., Lee, J., Kang, D., Choi, H.-W. & Ko, S.-H. (2014) 'Quantities, composition, and sources of beach debris in Korea from the results of nationwide monitoring'. *Marine Pollution Bulletin*, 84 (1). pp 27-34.
- Hopewell, J., Dvorak, R. & Kosior, E. (2009) 'Plastics recycling: challenges and opportunities'. *Philosophical Transactions of the Royal Society B*, 364 pp 2115-2126.
- Hughes, G. (1970) *Further studies on marine turtles in Tongaland, III. Lammergeyer*, 12, 7-25. Available.
- Hughes, G. (1974) *The sea turtles of South-East Africa. I. Status, morphology and distribution. South African Association of Marine Biological Research, and Oceanographic Research. Institute Investigational Report*, 35, 144p. Available.
- Hussain, N., V. Jaitley and A. T. Florence (2001). 'Recent advances in the understanding of uptake of microparticulates across the gastrointestinal lymphatics'. *Advanced Drug Delivery Reviews* 50(1-2): 107-142.
- Hutton, I., Carlile, N. & Priddel, D. (2008) 'Plastic ingestion by flesh-footed (Puffinus carneipes) and wedge-tailed (P. pacificus) shearwaters', *Papers and Proceedings of the Royal Society of Tasmania*. pp. 1.
- Imber, M. J. In Day, R., Wehle, D. & Coleman, F. (1985) Ingestion of plastic pollutants by marine birds. In Shomura, R. & Yoshida, H. (Eds.). *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 27–29 November 1984, Honolulu, US Department of Commerce (1985), pp. 387–429 NOAA Technical Memorandum NMFS SWFC-54.
- Imhof, H. K., Ivleva, N. P., Schmid, J., Niessner, R. & Laforsch, C. (2013) 'Contamination of beach sediments of a sub-alpine lake with microplastic particles'. *Current Biology*, 23 (19). pp R867- R868.
- Imhof, H. K., Schmid, J., Niessner, R., Ivleva, N. P. & Laforsch, C. (2012) 'A novel, highly efficient method for the separation and quantification of plastic particles in sediments of aquatic environments'. *Limnology and Oceanography-Methods*, 10 pp 524-537.
- Ivar do Sul, J. A. & Costa, M. F. (2014) 'The present and future of microplastic pollution in the marine environment'. *Environmental Pollution*, 185 (0). pp 352-364.

- Ivar do Sul, J. A., Costa, M. F., Barletta, M. & Cysneiros, F. J. A. (2013) 'Pelagic microplastics around an archipelago of the Equatorial Atlantic'. *Marine Pollution Bulletin*, 75 pp 305-309.
- Ivar do Sul, J. A., Spengler, Å. & Costa, M. F. (2009) 'Here, there and everywhere. Small plastic fragments and pellets on beaches of Fernando de Noronha (Equatorial Western Atlantic)'. *Marine Pollution Bulletin*, 58 (8). pp 1236-1238.
- Jantz, L. A., Morishige, C. L., Bruland, G. L. & Lepczyk, C. A. (2013) 'Ingestion of plastic marine debris by longnose lancetfish (*Alepisaurus ferox*) in the North Pacific Ocean'. *Marine Pollution Bulletin*, 69 (1-2). pp 97-104.
- Jayasiri, H., Purushothaman, C. & Vennila, A. (2013) 'Quantitative analysis of plastic debris on recreational beaches in Mumbai, India'. *Marine Pollution Bulletin*, 77 (1). pp 107-112.
- Jayasiri, H., Purushothaman, C. & Vennila, A. (2014) 'Occurrence of Polycyclic Aromatic Hydrocarbons (Pahs) in Beached Plastic Pellets from Mumbai Coast, India', *Proceedings of International Forestry and Environment Symposium*.
- Kaberi, H., Tsangaris, C., Zeri, C., Mousdis, G., Papadopoulos, A. & Streftaris, N. (2013) 'Microplastics along the shoreline of a Greek island (Kea isl., Aegean Sea): types and densities in relation to beach orientation, characteristics and proximity to sources', *Proceedings of the 4th International Conference on Environmental Management, Engineering, Planning and Economics (CEMEPE) and SECOTOX Conference*. Mykonos Island, Greece, June 24-28, 2013.
- Kako, S., A. Isobe and S. Magome 'Sequential monitoring of beach litter using webcams'. *Marine Pollution Bulletin* 60(5): 775-779.
- Kaminuma, T., Ohtake, C. & Kabuyama, N. (2000) 'Distribution and origin of plastic resin pellets as environmental pollutants at the East China Sea area'. *Kokuritsu Iyakuhiin Shokuhin Eisei Kenkyusho Hokoku*, 118 pp 90-99.
- Kaposi, K. L., Mos, B., Kelaher, B. & Dworjanyn, S. A. (2014) 'Ingestion of microplastic has limited impact on a marine larva'. *Environmental Science & Technology*, 48 (3). pp 1638-1645.
- Karapanagioti, H., Endo, S., Ogata, Y. & Takada, H. (2011) 'Diffuse pollution by persistent organic pollutants as measured in plastic pellets sampled from various beaches in Greece'. *Marine Pollution Bulletin*, 62 (2). pp 312-317.
- Karapanagioti, H. & Klontza, I. (2007) 'Investigating the properties of plastic resin pellets found in the coastal areas of Lesvos island'. *Global nest. The international journal*, 9 (1). pp 71-76.
- Karapanagioti, H. K. & Klontza, I. (2008) 'Testing phenanthrene distribution properties of virgin plastic pellets and plastic eroded pellets found on Lesvos island beaches (Greece)'. *Marine Environmental Research*, 65 (4). pp 283-290.
- Kartar, S., Abou-Seedo, F. & Sainsbury, M. (1976) 'Polystyrene spherules in the Severn Estuary – a progress report'. *Marine Pollution Bulletin*, 7 (3). pp 52.
- Kasperson, J. X., Kasperson, R. E., Pidgeon, N. & Slovic, P. (2003) 'The social amplification of risk: assessing fifteen years of research and theory'. In Pidgeon, N., Kasperson, R.E. and Slovic, P. (eds.) *The social amplification of risk*. UK: Cambridge University Press.
- Keep Los Angeles Beautiful (2009) *Littering and the iGeneration: City-wide intercept study of youth litter behaviour in Los Angeles*. congresopsicamb2015.com/ Available.
- Kelly, B. C., Gobas, F. A. & McLachlan, M. S. (2004) 'Intestinal absorption and biomagnification of organic contaminants in fish, wildlife, and humans'. *Environmental Toxicology and Chemistry*, 23 (10). pp 2324-2336.
- Khordagui, H. K. & Abu-Hilal, A. H. (1994) 'Industrial plastic on the southern beaches of the Arabian Gulf and the western beaches of the Gulf of Oman'. *Environmental Pollution*, 84 (3). pp 325-327.
- KIMO (2014) 'Fishing for Litter'. <http://www.kimointernational.org/FishingforLitter.aspx>. Klöckner, C. A. (2013) 'A comprehensive model of the psychology of environmental behaviour—A meta-analysis'. *Global Environmental Change*, 23 (5). pp 1028-1038.
- Kirchman, D. and R. Mitchell (1982). 'Contribution of particle-bound bacteria to total microheterotrophic activity in 5 ponds and 2 marshes'. *Applied and Environmental Microbiology* 43(1): 200-209.
- Klöckner, C.A. (2013). A comprehensive model of the psychology of environmental behaviour—A meta-analysis. *Global Environmental Change*, 23(5), 1028-1038.
- Koch, H. M. & Calafat, A. M. (2009) 'Human body burdens of chemicals used in plastic manufacture'. *Philosophical Transactions of the Royal Society B*, 364 pp 2063-2078.
- Koelmans, A. A., Besseling, E. & Foekema, E. M. (2014) 'Leaching of plastic additives to marine organisms'. *Environmental Pollution*, 187 (0). pp 49-54.
- Koelmans, A. A., Besseling, E., Wegner, A. & Foekema, E. M. (2013) 'Plastic as a Carrier of POPs to Aquatic Organisms: A Model Analysis'. *Environmental Science & Technology*, 47 (14). pp 7812- 7820.

- Koelmans, A. A., Gouin, T., Thompson, R., Wallace, N. & Arthur, C. (2014) 'Plastics in the marine environment'. *Environmental Toxicology and Chemistry*, 33 (1). pp 5-10.
- Kripa, V., Nair, P. G., Dhanya, A., Pravitha, V., Abhilash, K., Mohammed, A. A., Vijayan, D., Vishnu, P., Mohan, G. & Anilkumar, P. (2014) 'Microplastics in the gut of anchovies caught from the mud bank area of Alappuzha, Kerala'. *Marine Fisheries Information Service; Technical and Extension Series*, (219). pp 27-28.
- Kukulka, T., G. Proskurowski, S. Moret-Ferguson, D. W. Meyer and K. L. Law (2012). 'The effect of wind mixing on the vertical distribution of buoyant plastic debris'. *Geophysical Research Letters* 39.
- Kupryianchyk, D., Rakowska, M., Roessink, I., Reichman, E., Grotenhuis, J. & Koelmans, A. (2013) 'In situ treatment with activated carbon reduces bioaccumulation in aquatic food chains'. *Environmental Science & Technology*, 47 (9). pp 4563-4571.
- Kuriyama, Y., Konishi, K., Kanehiro, H., Otake, C., Kaminuma, T., Mato, Y., Takada, H. & Kojima, A. (2002) 'Plastic pellets in the marine environment of Tokyo Bay and Sagami Bay'. *Nippon Suisan Gakkaishi*, 68 (2). pp 164-171.
- Kusui, T. & Noda, M. (2003) 'International survey on the distribution of stranded and buried litter on beaches along the Sea of Japan'. *Marine Pollution Bulletin*, 47 (1-6). pp 175-179.
- Kwan, C. S., Takada, H., Boonyatumanond, R., Kato, Y., Mizukawa, K., Ito, M., Dung, L. Q., Zakaria, M. P. & Santiago, E. C. (2014) 'Historical occurrences of polybrominated diphenyl ethers and polychlorinated biphenyls in Manila Bay, Philippines, and in the upper Gulf of Thailand'. *Science of The Total Environment*, 470-471 (0). pp 427-437.
- L'Oreal (2014) 'News: L'Oréal commits to phase out all polyethylene microbeads from its scrubs by 2017'. <http://www.loreal.com/news/loreal-commits-to-phase-out-all-polyethylene-microbeads-from-its-scrubs-by-2017.aspx>. (Accessed: 5th August 2014). Laboratoire Boidoxis/Foundation CIMA.
- Lambert, S., Sinclair, C. J., Bradley, E. L. & Boxall, A. B. A. (2013) 'Effects of environmental conditions on latex degradation in aquatic systems'. *Science of the Total Environment*, 447 (0). pp 225-234.
- Lattin, G. L., Moore, C. J., Zellers, A. F., Moore, S. L. & Weisberg, S. B. (2004) 'A comparison of neustonic plastic and zooplankton at different depths near the southern California shore'. *Marine Pollution Bulletin*, 49 (4). pp 291-294.
- Lavers, J. L., A. L. Bond and I. Hutton (2014). 'Plastic ingestion by Flesh-footed Shearwaters (*Puffinus carneipes*): Implications for fledgling body condition and the accumulation of plastic-derived chemicals'. *Environmental Pollution* 187: 124-129.
- Law, K. L., Moret-Ferguson, S., Maximenko, N. A., Proskurowski, G., Peacock, E. E., Hafner, J. & Reddy, C. M. (2010) 'Plastic Accumulation in the North Atlantic Subtropical Gyre'. *Science*, 329 (5996). pp 1185-1188.
- Law, K. L., Moret-Ferguson, S. E., Goodwin, D. S., Zettler, E. R., De Force, E., Kukulka, T. & Proskurowski, G. (2014) 'Distribution of Surface Plastic Debris in the Eastern Pacific Ocean from an 11-Year Data Set'. *Environmental Science & Technology*, 48 (9). pp 4732-4738.
- Lebreton, L. C., S. D. Greer and J. C. Borrero (2012). 'Numerical modelling of floating debris in the world's oceans'. *Marine Pollution Bulletin* 64(3): 653-661.
- Lee, H., Shim, W. J. & Kwon, J.-H. (2014) 'Sorption capacity of plastic debris for hydrophobic organic chemicals'. *Science of the Total Environment*, 470-471 pp 1545-1552.
- Lee, J., Hong, S., Song, Y. K., Hong, S. H., Jang, Y. C., Jang, M., Heo, N. W., Han, G. M., Lee, M. J. & Kang, D. (2013a) 'Relationships among the abundances of plastic debris in different size classes on beaches in South Korea'. *Marine Pollution Bulletin*, 77 (1). pp 349-354.
- Lee, K.-W., Shim, W. J., Kwon, O. Y. & Kang, J.-H. (2013b) 'Size-dependent effects of micro polystyrene particles in the marine copepod *Tigriopus japonicus*'. *Environmental Science & Technology*, 47 (19). pp 11278-11283.
- Leichter, J. J. (2011) 'Investigating the Accumulation of Plastic Debris in the North Pacific Gyre' in Omori, K., Guo, X., Yoshie, N., Fujii, N., Handoh, I. C., Isobe, A. and Tanabe, S. (eds.) *Interdisciplinary Studies on Environmental Chemistry – Modeling & Analysis of Marine Environmental Problems*. TERRAPUB, pp 251-259.
- Leslie, H., van der Meulen, M., Kleissen, F. & Verhaak, A. (2011) *Microplastic litter in the Dutch marine environment*. 1203772-000 © Deltares, 2011. Available.
- Leslie, H. A., van Velzen, M. J. M., Vethaak, A. D (2013) *Microplastic survey of the Dutch environment. Novel data set of microplastics in North Sea sediments, treated wastewater effluents and marine biots*. Amsterdam, The Netherlands: IVM Institute for Environmental Studies, Final report R-13/11. Available.
- Liebezeit, G. & Dubaish, F. (2012) 'Microplastics in Beaches of the East Frisian Islands Spiekeroog and Kachelotplate'. *Bulletin of Environmental Contamination and Toxicology*, 89 (1). pp 213-217.

- Lima, A. R. A., Costa, M. F. & Barletta, M. (2014) 'Distribution patterns of microplastics within the plankton of a tropical estuary'. *Environmental Research*, 132 (0). pp 146-155.
- Lobelle, D. & Cunliffe, M. (2011) 'Early microbial biofilm formation on marine plastic debris'. *Marine Pollution Bulletin*, 62 (1). pp 197-200.
- Löder, M. & Gerdtz, G. (2013) 'FT-IR analysis for monitoring marine microplastics'. YOUMARES 4, Oldenburg, 11 September 2013 - 13 September 2013.
- Lusher, A. L., McHugh, M. & Thompson, R. C. (2013) 'Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel'. *Marine Pollution Bulletin*, 67 (1-2). pp 94-99.
- MacArthur, R.H. and Wilson, E.O. (1967) The theory of island biogeography. Princeton University Press. 224pp.
- Macfadyen, G., T. Huntington and R. Cappell (2009). Abandoned, lost or otherwise discarded fishing gear. UNEP Regional Seas Reports and Studies 185; FAO Fisheries and Aquaculture Technical Paper 523.
- Magnusson, K. (2014) *Microplastic and other microscopic anthropogenic particles in the sea area off Rauma and Turku, Finland*. Thesis, Swedish Environmental Research Institute.
- Majer, A. P., Vedolin, M. C. & Turra, A. (2012) 'Plastic pellets as oviposition site and means of dispersal for the ocean-skater insect Halobates'. *Marine Pollution Bulletin*, 64 (6). pp 1143-1147.
- Mallory, M. L. (2008) 'Marine plastic debris in northern fulmars from the Canadian high Arctic'. *Marine Pollution Bulletin*, 56 (8). pp 1501-1504.
- MARPOL (1973). International Convention for the Prevention of Pollution from Ships. <http://www.imo.org/About/Conventions/ListOfConventions/Pages/International-Convention-for-the-Prevention-of-Pollution-from-Ships-%28MARPOL%29.aspx>
- Marris, C., Wynne, B., Simmons, P. & Weldon, S. (2001) 'Public perceptions of agricultural biotechnologies in Europe'. Lancaster, UK: Lancaster University, Report of the PABE project funded by the European Commission, DG Research (contract number: FAIR CT98-3844 (DG12 - SSMI). 113pp.
- Martinez, M. E., S. Medina, M. T. del Campo, M. J. Sanchez-Cabezudo, M. Sanchez and L. Munuera (1998). 'Effect of polyethylene on osteocalcin, alkaline phosphatase and procollagen secretion by human osteoblastic cells'. *Calcified Tissue International* 62(5): 453-456.
- Martins, J. & Sobral, P. (2011) 'Plastic marine debris on the Portuguese coastline: A matter of size?'. *Marine Pollution Bulletin*, 62 (12). pp 2649-2653.
- Mascarenhas, R., Santos, R. & Zeppelini, D. (2004) 'Plastic debris ingestion by sea turtle in Paraiba, Brazil'. *Marine Pollution Bulletin*, 49 (4). pp 354-355.
- Mathalon, A. & Hill, P. (2014) 'Microplastic fibers in the intertidal ecosystem surrounding Halifax Harbor, Nova Scotia'. *Marine Pollution Bulletin*, 81 (1). pp 69-79.
- Mato, Y., Isobe, T., Takada, H., Kanehiro, H., Ohtake, C. & Kaminuma, T. (2001) 'Plastic Resin Pellets as a Transport Medium for Toxic Chemicals in the Marine Environment'. *Environmental Science and Technology*, 35 (2). pp 318-324.
- Mato, Y., Takada, H., Zakaria, M., Kuriyama, Y. & Kanehiro, H. (2002) 'Toxic chemicals contained in plastic resin pellets in the marine environment: Spatial difference in pollutant concentrations and the effects of resin type'. *Environmental Science*, 15 pp 415-423.
- Matthies, M., Schulz, M., Neumann, D., Klasmeier, J., Nülle, M.-T., Dekiff, J., Fleet, D., Guse, N., Markones, N. & Weiel, S. (2013) 'Compilation and assessment of selected anthropogenic pressures in the context of the Marine Strategy Framework Directive Descriptor 10-marine litter', *International Conference on Prevention and Management of Marine Litter in European Seas*. Berlin, 10 April 2013 – 12 April 2013.
- Maximenko, N., Hafner, J. & Nüller, P. (2012) 'Pathways of marine debris derived from trajectories of Lagrangian drifters'. *Marine Pollution Bulletin*, 65 (1-3). pp 51-62.
- Mayer, F. S. & Frantz, C. M. (2004) 'The connectedness to nature scale: A measure of individuals' feeling in community with nature'. *Journal of Environmental Psychology*, 24 (4). pp 503-515.
- McDermid, K. J. & McMullen, T. L. (2004) 'Quantitative analysis of small-plastic debris on beaches in the Hawaiian archipelago'. *Marine Pollution Bulletin*, 48 (7-8). pp 790-794.
- McDonnell, A. M. P. and K. O. Buesseler (2010). 'Variability in the average sinking velocity of marine particles'. *Limnology and Oceanography* 55(5): 2085-2096.

- McKenna, J., Williams, A. T. & Cooper, J. A. G. (2011) 'Blue Flag or Red Herring: Do beach awards encourage the public to visit beaches?'. *Tourism Management*, 32 (3). pp 576-588.
- McKinley, E. & Fletcher, S. (2010) 'Individual responsibility for the oceans? An evaluation of marine citizenship by UK marine practitioners'. *Ocean & Coastal Management*, 53 (7). pp 379- 384.
- Meeker, J. D., Sathyanarayana, S. & Swan, S. H. (2009) 'Phthalates and other additives in plastics: human exposure and associated health outcomes'. *Philosophical Transactions of the Royal Society B*, 364 pp 2097-2113.
- Mellon, J. (2013) 'Where and When Can We Use Google Trends to Measure Issue Salience?'. *PS: Political Science & Politics*, 46 (02). pp 280-290.
- Meylan, A. (1984) *Feeding ecology of the hawksbill turtle (Eretmochelys imbricata)*. Ph.D. Thesis, University of Florida, Gainesville, 115 p. Available.
- Mizukawa, K., Takada, H., Ito, M., Geok, Y. B., Hosoda, J., Yamashita, R., Saha, M., Suzuki, S., Miguez, C. & Frias, J. (2013) 'Monitoring of a wide range of organic micropollutants on the Portuguese coast using plastic resin pellets'. *Marine Pollution Bulletin*, 70. pp 296-302.
- Mohamed Nor, N. H. & Obbard, J. P. (2014) 'Microplastics in Singapore's coastal mangrove ecosystems'. *Marine Pollution Bulletin*, 79. pp 278-283.
- Mohee, R. & Unmar, G. (2007) 'Determining biodegradability of plastic materials under controlled and natural composting environments'. *Waste Management*, 27 (11). pp 1486-1493.
- Moore, M. N. (2006). 'Do nanoparticles present ecotoxicological risks for the health of the aquatic environment?' *Environment International* 32(8): 967-976.
- Moore, C., Lattin, G. L. & Zellers, A. F. (2011) 'Quantity and type of plastic debris flowing from two urban rivers to coastal waters and beaches of Southern California'. *Journal of Integrated Coastal Zone Management*, 11 (1). pp 65-73.
- Moore, C. J. (2008) 'Synthetic polymers in the marine environment: A rapidly increasing, long-term threat'. *Environmental Research*, 108 (2). pp 131-139.
- Moore, C. J., Moore, S. L., Leecaster, M. K. & Weisberg, S. B. (2001) 'A Comparison of Plastic and Plankton in the North Pacific Central Gyre'. *Marine Pollution Bulletin*, 42 (12). pp 1297-1300.
- Moore, C. J., Moore, S. L., Weisberg, S. B., Lattin, G. L. & Zellers, A. F. (2002) 'A comparison of neustonic plastic and zooplankton abundance in southern California's coastal waters'. *Marine Pollution Bulletin*, 44 (10). pp 1035-1038.
- Moore, M. (2006) 'Do nanoparticles present ecotoxicological risks for the health of the aquatic environment?'. *Environment International*, 32 (8). pp 967-976.
- Moore, S. A. & Polley, A. (2007) 'Defining indicators and standards for tourism impacts in protected areas: Cape Range National Park, Australia'. *Environmental Management*, 39 (3). pp 291-300.
- Morét-Ferguson, S., Law, K. L., Proskurowski, G., Murphy, E. K., Peacock, E. E. & Reddy, C. M. (2010) 'The size, mass, and composition of plastic debris in the western North Atlantic Ocean'. *Marine Pollution Bulletin*, 60 (10). pp 1873-1878.
- Morris, R. J. (1980) 'Plastic debris in the surface waters of the South Atlantic'. *Marine Pollution Bulletin*, 11 (6). pp 164-166.
- Moser, M. L. & Lee, D. S. (1992) 'A Fourteen-Year Survey of Plastic Ingestion by Western North Atlantic Seabirds'. *Colonial Waterbirds*, 15 (1). pp 83-94.
- Mouat, J., Lopez Lozano, R. & Bateson, H. (2010) *Economic impacts of marine litter*. UK: Kommunenenes Internasjonale Miljøorganisasjon (KIMO International). Available.
- Murray, F. & Cowie, P. R. (2011) 'Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758)'. *Marine Pollution Bulletin*, 62 (6). pp 1207-1217.
- Muthukumar, T., A. Aravinthan, K. Lakshmi, R. Venkatesan, L. Vedaprakash and M. Doble (2011). 'Fouling and stability of polymers and composites in marine environment'. *International Biodeterioration & Biodegradation* 65(2): 276-284.
- Nakashima, E., A. Isobe, S. I. Kako, T. Itai and S. Takahashi (2012). 'Quantification of Toxic Metals Derived from Macroplastic Litter on Ookushi Beach, Japan'. *Environmental Science & Technology* 46(18): 10099-10105.
- Ness, B., S. Anderberg and L. Olsson (2010). 'Structuring problems in sustainability science: The multi-level DPSIR framework'. *Geoforum* 41 (3). pp 479-488.
- Ng, K. & Obbard, J. (2006) 'Prevalence of microplastics in Singapore's coastal marine environment'. *Marine Pollution Bulletin*, 52 (7). pp 761-767.

- Nich, C., J. Langlois, A. Marchadier, C. Vidal, M. Cohen-Solal, H. Petite and M. Hamadouche (2011). 'Oestrogen deficiency modulates particle-induced osteolysis'. *Arthritis research & therapy* 13 (3). pp R100-R100.
- Nisbet, E. K., Zelenski, J. M. & Murphy, S. A. (2008) 'The nature relatedness scale: Linking individuals' connection with nature to environmental concern and behavior'. *Environment and Behavior*, 41. pp 715-740.
- Nobre, C. R., M. F. M. Santana, A. Maluf, F. S. Cortez, A. Cesar, C. D. S. Pereira and A. Turra (2015). 'Assessment of microplastic toxicity to embryonic development of the sea urchin *Lytechinus variegatus* (Echinodermata: Echinoidea)'. *Marine Pollution Bulletin*.
- Norén, F. (2008) *Small plastic particles in coastal Swedish waters*. N-Research report, commissioned by KIMO Sweden. 11 pp. Available.
- Norén, F. & Ekendahl, S. (2009) *Microscopic Anthropogenic Particles in Swedish Waters: many more than believed*. Schwerin, Germany: Helsinki Commission. 3 pp. Available.
- Norén, F., Naustvoll, L.J. (2011) *Survey of microplastic anthropogenic particles in Skagerrak*. Flødevigen, Norway: Institute of Marine Research. 21 pp. Available.
- Nuelle, M.-T., Dekiff, J. H., Remy, D. & Fries, E. (2014) 'A new analytical approach for monitoring microplastics in marine sediments'. *Environmental pollution (Barking, Essex: 1987)*, 184 pp 161- 169.
- Obbard, R. W., Sadri, S., Wong, Y. Q., Khitun, A. A., Baker, I. & Thompson, R. C. (2014) 'Global warming releases microplastic legacy frozen in Arctic Sea ice'. *Earth's Future*, pp 2014EF000240.
- Ocean Conservancy (2014) *Turning the tide on trash*. International Coastal Cleanup 2014. 28 pp. Available.
- Ocean Project (1999) Questionnaire and topline results from a national poll regarding oceans. Retrieved October 16th 2013 from http://theoceanproject.org/wp-content/uploads/2012/1999/Poll_Questionnaire_1999.pdf Available.
- Oehlmann, J., U. Schulte-Oehlmann, W. Kloas, O. Jagnytsch, I. Lutz, K. O. Kusk, L. Wollenberger, E. M. Santos, G. C. Paull, K. J. W. VanLook and C. R. Tyler (2009). 'A critical analysis of the biological impacts of plasticizers on wildlife'. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364(1526): 2047-2062.
- Ogata, Y., Takada, H., Mizukawa, K., Hirai, H., Iwasa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M. P., Dung, L. Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagioti, H. K., Weerts, S., McClurg, T., Burres, E., Smith, W., Velkenburg, M. V., Lang, J. S., Lang, R. C., Laursen, D., Danner, B., Stewardson, N. & Thompson, R. C. (2009) 'International Pellet Watch: Global monitoring of persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs'. *Marine Pollution Bulletin*, 58 (10). pp 1437-1446.
- Ogi, H., Baba, N., Ishihara, S. & Shibata, Y. (1999) 'Sampling of plastic pellets by two types of neuston net and plastic pollution in the sea'. *Bulletin of the Faculty of Fisheries, Hokkaido University*, 50 (2). pp 77-91.
- Ogi, H. & Fukumoto, Y. (2000) 'A sorting method for small plastic debris floating on the sea surface and stranded on sandy beaches'. *Bulletin of the Faculty of Fisheries, Hokkaido University*, 51 (2). pp 71-93.
- Ohlendorf, H., Risebrough, R. & Vermeer, K. (1978) Exposure of marine birds to environmental pollutants. U.S. *Fish & Wildlife Service, Wildlife Research Report*, 9, 1-40. Available.
- Oliveira, M., Ribeiro, A. & Guilhermino, L. (2012a) 'Effects of short-term exposure to microplastics and pyrene on *Pomatoschistus microps* (Teleostei, Gobiidae)'. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 163, Supplement (0). pp S20.
- Oliveira, M., Ribeiro, A. & Guilhermino, L. (2012b) 'Effects of exposure to microplastics and PAHs on microalgae *Rhodomonas baltica* and *Tetraselmis chuii*'. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 163, Supplement (0). pp S19-S20.
- Oliveira, M., Ribeiro, A., Hylland, K. & Guilhermino, L. (2013) 'Single and combined effects of microplastics and pyrene on juveniles (0+ group) of the common goby *Pomatoschistus microps* (Teleostei, Gobiidae)'. *Ecological Indicators*, 34 pp 641-647.
- Olsen, S. O. (2004) 'Antecedents of seafood consumption behavior: an overview'. *Journal of Aquatic Food Product Technology*, 13 (3). pp 79-91.
- O'Neill, S. J. and N. Smith (2014). 'Climate change and visual imagery'. *Wiley Interdisciplinary Reviews: Climate Change* 5(1): 73-87.
- OSPAR (2007) *OSPAR Pilot Project on Monitoring Marine Beach Litter: Monitoring of marine litter on beaches in the OSPAR region*. London: OSPAR Commission. Available.

- Paerl, H. W. (1975). 'Microbial Attachment to Particles in Marine and Freshwater Ecosystems.' *Microbial Ecology* 2(1): 73-83.
- Pahl, S., Sheppard, S., Boomsma, C. & Groves, C. (2014) 'Perceptions of time in relation to climate change'. *Wiley Interdisciplinary Reviews: Climate Change*, 5(3). pp 375-388.
- Pegram, J. E. and A. L. Andrady (1989). 'Outdoor weathering of selected polymeric materials under marine exposure conditions.' *Polymer Degradation and Stability* 26(4): 333-345.
- Peterlin, M., Kontic, B. & Kross, B. C. (2005) 'Public perception of environmental pressures within the Slovene coastal zone'. *Ocean & Coastal Management*, 48(2). pp 189-204.
- Peterson, C. H. & Lubchenco, J. (1997) 'Marine ecosystem services'. *Nature's Services: Societal Dependence on Natural Ecosystems. Edited by GC Daily. Island Press (Washington, DC)*, pp 177-194.
- Petit, A., I. Catelas, J. Antoniou, D. J. Zukor and O. L. Huk (2002). 'Differential apoptotic response of J774 macrophages to alumina and ultra-high-molecular-weight polyethylene particles'. *Journal of Orthopaedic Research* 20(1): 9-15.
- Petry, M. V. & Fonseca, V. d. S. (2002) 'Effects of human activities in the marine environment on seabirds along the coast of Rio Grande do Sul, Brazil'. *Ornitol. Neotrop.*, 13 pp 137-142.
- Pettit, T. N., Grant, G. S. & Whittow, G. C. (1981) 'Ingestion of plastics by Laysan albatross'. *The Auk*, 98 (4). pp 839-841.
- Pfeiffer, T. J. & Wills, P. S. (2011) 'Evaluation of three types of structured floating plastic media in moving bed bio-filters for total ammonia nitrogen removal in a low salinity hatchery recirculating aquaculture system'. *Aquacultural Engineering*, 45 (2). pp 51-59.
- Pham, C.K., et al. (2014) 'Marine litter distribution and density in European seas, from the shelves to deep basins'. *PLoS ONE*, 9(4). pp e95839-e95839.
- Phillips, M. & House, C. (2009) 'An evaluation of priorities for beach tourism: Case studies from South Wales, UK'. *Tourism Management*, 30 (2). pp 176-183.
- Pichel, W. G., J. H. Churnside, T. S. Veenstra, D. G. Foley, K. S. Friedman, R. E. Brainard, J. B. Nicoll, Q. Zheng and P. Clemente-Colon (2007). 'Marine debris collects within the North Pacific Subtropical Convergence Zone'. *Marine Pollution Bulletin*, 54(8): 1207-1211.
- Pidgeon, N., Kasperson, R. E. & Slovic, P. (2003) *The social amplification of risk*. Cambridge, UK: Cambridge University Press.
- Pinnegar, J.K. (2014) DAPSTOM – An integrated database and portal for fish stomach records. Version 4.7. Cefas Contract Reprot DP332, C3746, ME1228. The Centre for Environment, Fisheries & Aquaculture Science, Lowestoft, UK.
- Plastic Soup Foundation (2013) 'Colgate-Palmolive, L'Oreal and Beiersdorf halt use of micro beads'. <http://plasticsoup-foundation.org/eng/veen-categorie/colgate-palmolive-loreal-and-beiersdorf-halt-use-of-micro-beads/>. (Accessed: 30th July 2013).
- Possatto, F. E., Barletta, M., Costa, M. F., Ivar do Sul, J. A. & Dantas, D. V. (2011) 'Plastic debris ingestion by marine catfish: An unexpected fisheries impact'. *Marine Pollution Bulletin*, 62 (5). pp 1098-1102.
- Potts, T., O'Higgins, T., Mee, L. & Pita, C. (2011) *Public perceptions of Europe's seas: A policy brief*. EU FP7 KNOWSEAS Project. ISBN 0-9529089-3-X. Available.
- Poulicek, M. and C. Jeuniaux (1991). 'Chitin biodegradation in marine environments - an experimental approach'. *Biochemical Systematics and Ecology* 19(5): 385-394.
- Powers, K. W., Brown, S. C., Krishna, V. B., Wasdo, S. C., Moudgil, B. M. & Roberts, S. M. (2006) 'Research Strategies for Safety Evaluation of Nanomaterials. Part VI. Characterization of Nanoscale Particles for Toxicological Evaluation'. *Toxicological Sciences*, 90 (2). pp 296-303.
- Pretty, J., Peacock, J., Sellens, M. & Griffin, M. (2005) 'The mental and physical health outcomes of green exercise'. *International Journal of Environmental Health Research*, 15 (5). pp 319-337.
- Pruter, A. T. (1987) 'Sources, quantities and distribution of persistent plastics in the marine environment'. *Marine Pollution Bulletin*, 18 (6, Supplement 2). pp 305-310.
- Qayyum, M. M. and J. R. White (1993). 'Effect of stabilizers on failure mechanisms in weathered polypropylene'. *Polymer Degradation and Stability* 41 (2): 163-172.
- Quik, J. T. K., M. C. Stuart, M. Wouterse, W. Peijnenburg, A. J. Hendriks and D. van de Meent (2012). 'Natural colloids are the dominant factor in the sedimentation of nanoparticles'. *Environmental Toxicology and Chemistry* 31(5): 1019-1022.

- Rebolledo, E. L. B., J. A. van Franeker, O. E. Jansen and S. M. J. M. Brasseur (2013). 'Plastic ingestion by harbour seals (*Phoca vitulina*) in The Netherlands'. *Marine Pollution Bulletin* 67(1-2): 200-202.
- Reddy, M. S., Shaik, B., Adimurthy, S. & Ramachandraiah, G. (2006) 'Description of the small plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India'. *Estuarine, Coastal and Shelf Science*, 68 (3-4). pp 656-660.
- Redford, D. P., Trulli, H. K. & Trulli, W. R. (1997) 'Sources of plastic pellets in the aquatic environment'. *Marine Debris*. Springer, pp 335-343.
- Reed, S. (1981) *Wreck of kerguelen and blue petrels*. *Notornis*, 28, 239-240. 239-240 pp. Available.
- Reisser, J., Shaw, J., Hallegraeff, G., Proietti, M., Barnes, D. K. A., Thums, M., Wilcox, C., Hardesty, B. D. & Pattiaratchi, C. (2014) 'Millimeter-Sized Marine Plastics: A New Pelagic Habitat for Microorganisms and Invertebrates'. *PLoS ONE*, 9 (6). pp e100289.
- Reisser, J., Shaw, J., Wilcox, C., Hardesty, B. D., Proietti, M., Thums, M. & Pattiaratchi, C. (2013) 'Marine Plastic Pollution in Waters around Australia: Characteristics, Concentrations, and Pathways'. *PLoS ONE*, 8 (11). pp e80466.
- Ribic C. A., S. B. Sheavly & D. J. Rugg (2011). Trends in Marine Debris in the U.S. Caribbean and the Gulf of Mexico 1996-2003. *Journal of Integrated Coastal Zone Management*, 11(1):7-19.
- Ribic, C. A., Sheavly, S. B., Rugg, D. J. & Erdmann, E. S. (2012) 'Trends in marine debris along the US Pacific Coast and Hawai'i 1998-2007'. *Marine Pollution Bulletin*, 64 (5). pp 994-1004.
- Rillig, M. C. (2012) 'Microplastic in Terrestrial Ecosystems and the Soil?'. *Environmental Science & Technology*, 46 (12). pp 6453-6454.
- Rios, L. M., Moore, C. & Jones, P. R. (2007) 'Persistent organic pollutants carried by synthetic polymers in the ocean environment'. *Marine Pollution Bulletin*, 54 (8). pp 1230-1237.
- Rios, L. M., P. R. Jones, C. Moore and U. V. Narayan (2010). 'Quantitation of persistent organic pollutants adsorbed on plastic debris from the Northern Pacific Gyre's "eastern garbage patch"'. *Journal of Environmental Monitoring*, 12(12): 2226-2236.
- Robards, M. D., Piatt, J. F. & Wohl, K. D. (1995) 'Increasing frequency of plastic particles ingested by seabirds in the subarctic North Pacific'. *Marine Pollution Bulletin*, 30 (2). pp 151-157.
- Rochman, C. M., Hentschel, B. T. & Teh, S. J. (2014a) 'Long-Term Sorption of Metals Is Similar among Plastic Types: Implications for Plastic Debris in Aquatic Environments'. *PLoS ONE*, 9 (1). pp e85433.
- Rochman, C. M., Hoh, E., Hentschel, B. T. & Kaye, S. (2012) 'Long-Term Field Measurement of Sorption of Organic Contaminants to Five Types of Plastic Pellets: Implications for Plastic Marine Debris'. *Environmental Science & Technology*, 47 (3). pp 1646-1654.
- Rochman, C. M., E. Hoh, B. T. Hentschel and S. Kaye (2013b). 'Long-term field measurement of sorption of organic contaminants to five types of plastic pellets: implications for plastic marine debris'. *Environmental Science & Technology* 47(3): 1646-1654.
- Rochman, C. M., Hoh, E., Kurobe, T. & Teh, S. J. (2013a) 'Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress'. *Scientific Reports*, 3; 3263.
- Rochman, C. M., Lewison, R. L., Eriksen, M., Allen, H., Cook, A.-M. & Teh, S. J. (2014b) 'Polybrominated diphenyl ethers (PBDEs) in fish tissue may be an indicator of plastic contamination in marine habitats'. *Science of the Total Environment*, 476-477 (0). pp 622-633.
- Roehl, W. S. & Ditton, R. B. (1993) 'Impacts of the offshore marine industry on coastal tourism: the case of Padre Island National Seashore'. *Coastal management*, 21 (1). pp 75-89.
- Rossi, G., Barnoud, J. & Monticelli, L. (2013) 'Polystyrene Nanoparticles Perturb Lipid Membranes'. *The Journal of Physical Chemistry Letters*, 5 (1). pp 241-246.
- Rothäusler, E., Gutow, L. & Thiel, M. (2012) 'Floating seaweeds and their communities'. *Seaweed Biology*. Springer, pp 359-380.
- Rothstein, S. I. (1973) 'Plastic Particle Pollution of the Surface of the Atlantic Ocean: Evidence from a Seabird'. *The Condor*, 75 (3). pp 344-345.
- Ruppert, E. E., Nash, T. R. & Smith, A. J. (2000) 'The size range of suspended particles trapped and ingested by the filter-feeding lancelet *Branchiostoma floridae* (Cephalochordata: Acrania)'. *Journal of the Marine Biological Association of the United Kingdom*, 80 (02). pp 329-332.

- Ryan, P. & Jackson, S. (1987) 'The lifespan of ingested plastic particles in seabirds and their effect on digestive efficiency'. *Marine Pollution Bulletin*, 18 (5). pp 217-219.
- Ryan, P. G. (1987) 'The incidence and characteristics of plastic particles ingested by seabirds'. *Marine Environmental Research*, 23 (3). pp 175-206.
- Ryan, P. G. (1988a) 'The characteristics and distribution of plastic particles at the sea-surface off the southwestern Cape Province, South Africa'. *Marine Environmental Research*, 25 (4). pp 249-273.
- Ryan, P. G. (1988b) 'Effects of ingested plastic on seabird feeding: Evidence from chickens'. *Marine Pollution Bulletin*, 19 (3). pp 125-128.
- Ryan, P. G. (1988c) 'Intraspecific Variation in Plastic Ingestion by Seabirds and the Flux of Plastic Through Seabird Populations'. *The Condor*, 90 (2). pp 446-452.
- Ryan, P. G. (2008) 'Seabirds indicate changes in the composition of plastic litter in the Atlantic and south-western Indian Oceans'. *Marine Pollution Bulletin*, 56 (8). pp 1406-1409.
- Ryan, P. G. (2013) 'A simple technique for counting marine debris at sea reveals steep litter gradients between the Straits of Malacca and the Bay of Bengal'. *Marine Pollution Bulletin*, 69 (1-2). pp 128-136.
- Ryan, P. G. (2014) 'Litter survey detects the South Atlantic 'garbage patch''. *Marine Pollution Bulletin*, 79 (1-2). pp 220-224.
- Ryan, P. G. and C. L. Moloney (1990). 'Plastic and other artefacts on South_african beaches – temporal trends in abundance and composition'. *South African Journal of Science* 86(7-10): 450-452.
- Ryan, P. G., Bouwman, H., Moloney, C. L., Yuyama, M. & Takada, H. (2012) 'Long-term decreases in persistent organic pollutants in South African coastal waters detected from beached polyethylene pellets'. *Marine Pollution Bulletin*, 64 (12). pp 2756-2760.
- Ryan, P. G., Connell, A. D. & Gardner, B. D. (1988) 'Plastic ingestion and PCBs in seabirds: Is there a relationship?'. *Marine Pollution Bulletin*, 19 (4). pp 174-176.
- Ryan, P. G., Moore, C. J., van Franeker, J. A. & Moloney, C. L. (2009) 'Monitoring the abundance of plastic debris in the marine environment'. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364 (1526). pp 1999-2012.
- Sadri, S. S. & Thompson, R. C. (2014) 'On the quantity and composition of floating plastic debris entering and leaving the Tamar Estuary, Southwest England'. *Marine Pollution Bulletin*, 81 (1). pp 55-60.
- Saido, K., Itagaki, T., Sato, H., Kodera, Y., Abe, O., Ogawa, N., Chung, S.-Y. & Miyashita, K. (2009) 'New contamination derived from marine debris plastics'. ACS. Washington, DC. Available at: <http://oasys2.confex.com/acs/238nm/techprogram/P1304877.HTM>
- Sajiki, J. & Yonekubo, J. (2003) 'Leaching of bisphenol A (BPA) to seawater from polycarbonate plastic and its degradation by reactive oxygen species'. *Chemosphere*, 51 (1). pp 55-62.
- Sajiki, J. & Yonekubo, J. (2004) 'Leaching of bisphenol A (BPA) from polycarbonate plastic to water containing amino acids and its degradation by radical oxygen species'. *Chemosphere*, 55 (6). pp 861-867.
- Sanchez, W., Bender, C. & Porcher, J.-M. (2014) 'Wild gudgeons (*Gobio gobio*) from French rivers are contaminated by microplastics: Preliminary study and first evidence'. *Environmental Research*, 128 (0). pp 98-100.
- Santos, I. R., Friedrich, A. C., Wallner-Kersanach, M. & Fillmann, G. (2005) 'Influence of socio- economic characteristics of beach users on litter generation'. *Ocean & Coastal Management*, 48 (9). pp 742-752.
- Satir, F. (2009) 'Microplastics, Macroproblem'. *Terrain*, [Online]. Available at: <http://www.tacoma.washington.edu/terrain/microplastics.html>.
- Savadori, L., Savio, S., Nicotra, E., Rumiati, R., Finucane, M. & Slovic, P. (2004) 'Expert and public perception of risk from biotechnology'. *Risk analysis*, 24 (5). pp 1289-1299.
- Savadori, L., Savio, S., Nicotra, E., Rumiati, R., Finucane, M. L. & Slovic, P. (2010) 'Expert and public perception of risk from biotechnology'. in Slovic, P. (ed.) *The Feeling of Risk: New Perspectives on Risk Perception*. Oxon, UK: Earthscan, pp 245-260.
- Schuyler, Q., Hardesty, B. D., Wilcox, C. & Townsend, K. (2012) 'To Eat or Not to Eat? Debris Selectivity by Marine Turtles'. *PLoS ONE*, 7 (7). pp e40884.
- Schuyler, Q. A., Wilcox, C., Townsend, K., Hardesty, B. D. & Marshall, N. J. (2014) 'Mistaken identity? Visual similarities of marine debris to natural prey items of sea turtles'. *BMC Ecology*, 14 (14).

- Schwemmer, P., Guepner, F., Guse, N. & Garthe, S. (2012) 'Food choice of birds from the German North Sea coast'. *Vogelwarte*, 50 (3). pp 141-154.
- Setälä, O., Fleming-Lehtinen, V. & Lehtiniemi, M. (2014) 'Ingestion and transfer of microplastics in the planktonic food web'. *Environmental Pollution*, 185 (0). pp 77-83.
- Shaw, D. G. & Day, R. H. (1994) 'Colour- and form-dependent loss of plastic micro-debris from the North Pacific Ocean'. *Marine Pollution Bulletin*, 28 (1). pp 39-43.
- Sheavly, S. B. & Register, K. M. (2007) 'Marine Debris & Plastics: Environmental Concerns, Sources, Impacts and Solutions'. *Journal of Polymers and the Environment*, 15 (4). pp 301-305.
- Shehada, A. & Land, M. (2014) *What are the effects of plastic particles on growth and mortality of marine organisms?* Mistra EviEM Pilot Study PS5 (www.eviem.se). Available.
- Shiber, J. (1979) 'Plastic pellets on the coast of Lebanon'. *Marine Pollution Bulletin*, 10 (1). pp 28- 30.
- Shiber, J. (1982) 'Plastic pellets on Spain's 'Costa del Sol' beaches'. *Marine Pollution Bulletin*, 13 (12). pp 409-412.
- Shiber, J. (1987) 'Plastic pellets and tar on Spain's Mediterranean beaches'. *Marine Pollution Bulletin*, 18 (2). pp 84-86.
- Shiber, J. & Barrales-Rienda, J. (1991) 'Plastic pellets, tar, and megalitter on Beirut beaches, 1977-1988'. *Environmental Pollution*, 71 (1). pp 17-30.
- Shin, J. S. and S. N. Abraham (2001). 'Cell biology - Caveolae - Not just craters in the cellular landscape'. *Science* 293(5534): 1447-1448.
- Sileo, L., Sievert, P. R., Samuel, M. D. & Fefer, S. I. (1990) 'Prevalence and characteristics of plastic ingested by Hawaiian seabirds', Shomura, R. and Godfrey, M. (eds.). *Proceedings of the Second International Conference on Marine Debris*. Honolulu, Hawaii US Department of Commerce, NOAA Technical Memorandum. NMFS. , pp. 665-681.
- Slavin, C., Grage, A. & Campbell, M. L. (2012) 'Linking social drivers of marine debris with actual marine debris on beaches'. *Marine Pollution Bulletin*, 64 (8). pp 1580-1588.
- Slovic, P. (1987) 'Perception of risk'. *Science*, 236 (4799). pp 280-285.
- Slovic, P. (1999) 'Trust, emotion, sex, politics, and science: Surveying the risk-assessment battlefield'. *Risk analysis*, 19 (4). pp 689-701.
- Song, Y. K., S. H. Hong, M. Jang, J.-H. Kang, O. Y. Kwon, G. M. Han and W. J. Shim (2014). 'Large Accumulation of Micro-sized Synthetic Polymer Particles in the Sea Surface Microlayer'. *Environmental Science & Technology* 48(16): 9014-9021.
- Spear, L. B., Ainley, D. G. & Ribic, C. A. (1995) 'Incidence of plastic in seabirds from the tropical Pacific, 1984-91: Relation with distribution of species, sex, age, season, year and body weight'. *Marine Environmental Research*, 40 (2). pp 123-146.
- Steel, B., Lovrich, N., Lach, D. & Fomenko, V. (2005a) 'Correlates and consequences of public knowledge concerning ocean fisheries management'. *Coastal management*, 33 (1). pp 37-51.
- Steel, B. S., Smith, C., Opsommer, L., Curiel, S. & Warner-Steel, R. (2005b) 'Public ocean literacy in the United States'. *Ocean & Coastal Management*, 48 (2). pp 97-114.
- Storrier, K. & McGlashan, D. (2006) 'Development and management of a coastal litter campaign: the voluntary coastal partnership approach'. *Marine Policy*, 30 (2). pp 189-196.
- Tabet, L., C. Bussy, A. Setyan, A. Simon-Deckers, M. J. Rossi, J. Boczkowski and S. Lanone (2011). 'Coating carbon nanotubes with a polystyrene-based polymer protects against pulmonary toxicity'. *Particle and Fibre Toxicology*, 8.
- Takada, H. (2006) 'Call for pellets! International Pellet Watch Global Monitoring of POPs using beached plastic resin pellets'. *Marine Pollution Bulletin*, 52 (12). pp 1547-1548.
- Takahashi, S., Mukai, H., Tanabe, S., Sakayama, K., Miyazaki, T. & Masuno, H. (1999) 'Butyltin residues in livers of humans and wild terrestrial mammals and in plastic products'. *Environmental Pollution*, 106 (2). pp 213-218.
- Tanaka, K., Takada, H., Yamashita, R., Mizukawa, K., Fukuwaka, M.-a. & Watanuki, Y. (2013) 'Accumulation of plastic-derived chemicals in tissues of seabirds ingesting marine plastics'. *Marine Pollution Bulletin*, 69 (1-2). pp 219-222.
- Tang, L. P. and J. W. Eaton (1995). 'Inflammatory responses to biomaterials'. *American Journal of Clinical Pathology*, 103(4): 466-471.
- Teuten, E. L., Rowland, S. J., Galloway, T. S. & Thompson, R. C. (2007) 'Potential for Plastics to Transport Hydrophobic Contaminants'. *Environmental Science & Technology*, 41 (22). pp 7759- 7764.

- Teuten, E. L., Saquing, J. M., Knappe, D. R., Barlaz, M. A., Jonsson, S., Björn, A., Rowland, S. J., Thompson, R. C., Galloway, T. S. & Yamashita, R. (2009) 'Transport and release of chemicals from plastics to the environment and to wildlife'. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364 (1526). pp 2027-2045.
- Thomann, R. V., Connolly, J. P. & Parkerton, T. F. (1992) 'An equilibrium model of organic chemical accumulation in aquatic food webs with sediment interaction'. *Environmental Toxicology and Chemistry*, 11 (5). pp 615-629.
- Thompson, R., C. Moore, A. Andrady, M. Gregory, H. Takada and S. Weisberg (2005). 'New directions in plastic debris'. *Science* 310(5751): 1117-1117.
- Thompson, R. C., Moore, C. J., Vom Saal, F. S. & Swan, S. H. (2009a) 'Plastics, the environment and human health: current consensus and future trends'. *Philosophical Transactions of the Royal Society B*, 364 pp 2153-2166.
- Thompson, R. C., Olsen, Y., Mitchell, R. P., Davis, A., Rowland, S. J., John, A. W., McGonigle, D. & Russell, A. E. (2004) 'Lost at sea: where is all the plastic?'. *Science*, 304 (5672). pp 838-838.
- Thompson, R. C., Swan, S. H., Moore, C. J. & Vom Saal, F. S. (2009b) 'Our plastic age'. *Philosophical Transactions of the Royal Society B*, 364 pp 1973-1976.
- Tourinho, P. S., Ivar do Sul, J. A. & Fillmann, G. (2010) 'Is marine debris ingestion still a problem for the coastal marine biota of southern Brazil?'. *Marine Pollution Bulletin*, 60 (3). pp 396-401.
- Tudor, D. & Williams, A. (2006) 'A rationale for beach selection by the public on the coast of Wales, UK'. *Area*, 38 (2). pp 153-164.
- Tudor, D. & Williams, A. (2008) 'Important aspects of beach pollution to managers: Wales and the Bristol Channel, UK'. *Journal of Coastal Research*, pp 735-745.
- Turner, A. & Holmes, L. (2011) 'Occurrence, distribution and characteristics of beached plastic production pellets on the island of Malta (central Mediterranean)'. *Marine Pollution Bulletin*, 62 (2). pp 377-381.
- Turra, A., Manzano, A. B., Dias, R. J. S., Mahiques, M. M., Barbosa, L., Balthazar-Silva, D. & Moreira, F. T. (2014) 'Three-dimensional distribution of plastic pellets in sandy beaches: shifting paradigms'. *Nature Scientific Reports*, 4 (4435).
- Ugolini, A. & Ungherese, G. (2011) 'Supralittoral amphipods and ingestion of microplastic debris'. *Biologia Marina Mediterranea*, 18 (1). pp 396-397.
- Ugolini, A., Ungherese, G., Ciofini, M., Lapucci, A. & Camaiti, M. (2013) 'Microplastic debris in sandhoppers'. *Estuarine, Coastal and Shelf Science*, 129. pp 19-22.
- UK National Ecosystem Assessment (2011) Technical report: Understanding nature's value to society. Retrieved 30th July 2014 from <http://uknea.unep-wcmc.org/Resources/tabid/82/Default.aspx>. Available.
- UK National Ecosystem Assessment [NEA] (2011) The UK National Ecosystem Assessment Technical Report. <http://uknea.unep-wcmc.org/Resources/tabid/82/Default.aspx>: Cambridge, UK: UNEP-WCMC. Available.
- UNEP (1995). Global programme of action for the protection of the marine environment from land-based activities. UNEP(OCA)/LBA/IG.2/7
- UNEP (2005). Marine litter, an analytical overview.
- UNEP (2009). Global threat, global challenge: review and analysis of UNEP's Global Initiative on Marine Litter.
- UNEP (2011). Year Book 2011. Emerging issues in our global environment.
- UNEP (2014). Proceedings of the United Nations Environment Assembly of the United Nations. Environment Programme at its first session, 23-27 June 2014. UNEP/EA.1/10
- UNGA (2005). Oceans and the law of the sea, Agenda item 75 (a), Sixtieth session United Nations General Assembly, UNGA.
- Unilever (2014) 'Micro-plastics'. Accessed: 5th August 2014 from <http://www.unilever.com/sustainable-living-2014/our-approach-to-sustainability/responding-to-stakeholder-concerns/micro-plastics>
- Van, A., Rochman, C. M., Flores, E. M., Hill, K. L., Vargas, E., Vargas, S. A. & Hoh, E. (2012) 'Persistent organic pollutants in plastic marine debris found on beaches in San Diego, California'. *Chemosphere*, 86 (3). pp 258-263.
- Van Cauwenberghe, L., Claessens, M. & Janssen, C. (2012b) 'Selective uptake of microplastics by a marine bivalve (*Mytilus edulis*)'. *VLIZ Special Publication*, 55
- Van Cauwenberghe, L., Claessens, M. & Janssen, C. (2014) 'Microplastic on our plate, also in our shops...', Mees, J. and Seys, J. (eds). Book of abstracts: VLIZ Young Scientists' Day. Brugge, Belgium, 7 March 2014. *VLIZ Special Publication*, 67. Vlaams Instituut voor de Zee - Flanders Marine Institute (VLIZ): Oostende. xii, 178 pp.

- Van Cauwenberghe, L., Claessens, M. & Janssen, C. R. (2013a) 'Selective uptake of microplastics by a marine bivalve (*Mytilus edulis*)'. *Communications in agricultural and applied biological sciences*, 78 (1). pp 25-27.
- Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M. & Janssen, C. (2012a) 'Occurrence of microplastics in mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) collected along the French-Belgian-Dutch coast'. *VLIZ Special Publication*, 55
- Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M. B., Mees, J. & Janssen, C. R. (2013b) 'Assessment of marine debris on the Belgian Continental Shelf'. *Marine Pollution Bulletin*, 73 (1). pp 161-169.
- Van Cauwenberghe, L. & Janssen, C. R. (2014) 'Microplastics in bivalves cultured for human consumption'. *Environmental Pollution*, 193 (0). pp 65-70.
- Van Cauwenberghe, L., Vanreusel, A., Mees, J. & Janssen, C. (2014) 'Microplastics... one league under the sea...', Mees, J. and Seys, J. eds.). *Book of abstracts: VLIZ Young Scientists' Day*. Brugge, Belgium, 7 March 2014. *VLIZ Special Publication*, 67. Vlaams Instituut voor de Zee - Flanders Marine Institute (VLIZ): Oostende. xii, 178 pp.
- Van Cauwenberghe, L., Vanreusel, A., Mees, J. & Janssen, C. R. (2013c) 'Microplastic pollution in deep-sea sediments'. *Environmental Pollution*, 182 (0). pp 495-499.
- van der Goot, F. Gisou, and Jean Gruenberg. 'Oiling the wheels of the endocytic pathway'. *Trends in cell biology* 12.7 (2002): 296-299
- van Franeker, J. (1983) Plastics—een bedreiging voor zeevogels. *Nieuwbrief NSO* 4, 41-61. Available.
- van Franeker, J. A. (1985) 'Plastic ingestion in the North Atlantic fulmar'. *Marine Pollution Bulletin*, 16 (9). pp 367-369.
- van Franeker, J. A. & Bell, P. J. (1988) 'Plastic ingestion by petrels breeding in Antarctica'. *Marine Pollution Bulletin*, 19 (12). pp 672-674.
- van Franeker, J. A., Blaize, C., Danielsen, J., Fairclough, K., Gollan, J., Guse, N., Hansen, P.-L., Heubeck, M., Jensen, J.-K., Le Guillou, G., Olsen, B., Olsen, K.-O., Pedersen, J., Stienen, E. W. M. & Turner, D. M. (2011) 'Monitoring plastic ingestion by the northern fulmar *Fulmarus glacialis* in the North Sea'. *Environmental Pollution*, 159 (10). pp 2609-2615.
- van Franeker, J. A., Heubeck, M., Fairclough, K., Turner, D. M., Stienen, E. W. M., Guse, N., Pedersen, J., Olsen, K. O., Andersson, P. J. & Olsen, B. (2005) 'Save the North Sea' *Fulmar Study 2002-2004: a regional pilot project for the Fulmar-Litter-EcoQO in the OSPAR area*. Den Berg (Texel), The Netherlands: ALTErrA, (Report No. Alterra-rapport 1162). Available.
- van Franeker, J. A. & Meijboom, A. (2003) *Marine Litter Monitoring by Northern Fulmars - Progress Report 2002*. Den Berg (Texel), The Netherlands: ALTErrA - Texel, Marine and Coastal Zone Research, (Report No. Alterra-rapport 622). 49 pp. Available.
- van Franeker, J. A., Meijboom, A. & de Jong, M. L. (2004) *Marine litter monitoring by Northern Fulmars in the Netherlands 1982-2003*. Den Burg (Texel), The Netherlands: ALTErrA, (Report No. Alterra-rapport 1093). 48 pp. Available.
- van Sebille, E., M. H. England and G. Froyland (2012). 'Origin, dynamics and evolution of ocean garbage patches from observed surface drifters'. *Environmental Research Letters* 7(4).
- Vandegehuchte, M., Van Cauwenberghe, L., Claessens, M. & Janssen, C. R. (2012) 'Plastic waste in the Belgian coastal waters: where and how much?', Mees, J. and Seys, J. eds.). *Book of abstracts VLIZ Young Scientist's Day*. Bruges, Belgium, 25 February 2011 VLIZ Special Publication, 48. Vlaams Insituut voor de Zee (VLIZ): Oostende, Belgium xiv + 143p.
- Vandendriessche, S., E. W. M. Stienen, M. Vincx and S. Degraer (2007). 'Seabirds foraging at floating seaweeds in the Northeast Atlantic.' *Ardea* 95(2): 289-298.
- Velzeboer, I., Kwadijk, C. J. A. F. & Koelmans, A. A. (2014) 'Strong Sorption of PCBs to Nanoplastics, Microplastics, Carbon Nanotubes, and Fullerenes'. *Environmental Science & Technology*, 48 (9). pp 4869-4876.
- Verbeke, W., Sioen, I., Pieniak, Z., Van Camp, J. & De Henauw, S. (2005) 'Consumer perception versus scientific evidence about health benefits and safety risks from fish consumption'. *Public health nutrition*, 8 (04). pp 422-429.
- Verlis, K. M., Campbell, M. L. & Wilson, S. P. (2013) 'Ingestion of marine debris plastic by the wedge-tailed shearwater *Ardenna pacifica* in the Great Barrier Reef, Australia'. *Marine Pollution Bulletin*, 72 (1). pp 244-249.
- Vianello, A., Boldrin, A., Guerriero, P., Moschino, V., Rella, R., Sturaro, A. & Da Ros, L. (2013) 'Microplastic particles in sediments of Lagoon of Venice, Italy: First observations on occurrence, spatial patterns and identification'. *Estuarine, Coastal and Shelf Science*, 130 pp 54-61.

- Vignola, R., Klinsky, S., Tam, J. & McDaniels, T. (2013) 'Public perception, knowledge and policy support for mitigation and adaption to Climate Change in Costa Rica: Comparisons with North American and European studies'. *Mitigation and Adaptation Strategies for Global Change*, 18 (3). pp 303-323.
- Vlietstra, L. S. & Parga, J. A. (2002) 'Long-term changes in the type, but not amount, of ingested plastic particles in short-tailed shearwaters in the southeastern Bering Sea'. *Marine Pollution Bulletin*, 44 (9). pp 945-955.
- Volckaert, A., Veiga, J. M. & Van Acoleyen, M. (2012) 'Sources and pathways for marine litter: Bottom up approach starting from local case-studies for the four European regional seas'. *VLIZ Special Publication*, 61
- von Moos, N. (2010) *Histopathological and cytochemical analysis of ingested polyethylene powder in the digestive gland of the blue mussel, Mytilus edulis (L.)*. Faculty of Science, University of Basel, Switzerland.
- von Moos, N., Burkhardt-Holm, P. & Koehler, A. (2012) 'Uptake and Effects of Microplastics on Cells and Tissue of the Blue Mussel *Mytilus edulis* L. after an Experimental Exposure'. *Environmental Science & Technology*, 46 (20). pp 11327-11335.
- Ward, J. E. and S. E. Shumway (2004). 'Separating the grain from the chaff: particle selection in suspension- and deposit-feeding bivalves'. *Journal of Experimental Marine Biology and Ecology* 300(1-2): 83-130.
- Watson, R., Albon, S., Aspinall, R., Austen, M., Bardgett, B., Bateman, I., Berry, P., Bird, W., Bradbury, R., Brown, C., Bullock, J., Burgess, J., Church, A., Christie, C., Crute, I., Davies, L., Edwards-Jones, G., Emmett, B., Firbank, L., Fitter, A., Gibson, A., Hails, R., Haines-Young, R. L., A., H., Heathwaite, L., Hopkins, J., Jenkins, M., Jones, L., Mace, G., Malcolm, S., Maltby, E., Maskell, L., Norris, K., Ormerod, S., Osborne, J., Pretty, J., Quine, C., Russell, S., Simpson, L., Smith, P., Tierney, M. K., Turner, Van der Wal, R., Vira, B., Walpole, M., Watkinson, A., Weighell, A., Winn, J. & Winter, M. (2011) *UK National Ecosystem Assessment : understanding nature's value to society. Synthesis of key findings*. Oxford: Information Press. Available.
- Watts, A. J. R., Lewis, C., Goodhead, R. M., Beckett, S. J., Moger, J., Tyler, C. R. & Galloway, T. S. (2014) 'Uptake and Retention of Microplastics by the Shore Crab *Carcinus maenas*'. *Environmental Science & Technology*, 48 (15). pp 8823-8830.
- Weatherell, C., Tregear, A. & Allinson, J. (2003) 'In search of the concerned consumer: UK public perceptions of food, farming and buying local'. *Journal of Rural Studies*, 19 (2). pp 233-244.
- Wegner, A., Besseling, E., Foekema, E. M., Kamermans, P. & Koelmans, A. A. (2012) 'Effects of nanopolystyrene on the feeding behavior of the blue mussel (*Mytilus edulis* L.)'. *Environmental Toxicology and Chemistry*, 31 (11). pp 2490-2497.
- Wesselinova, D. (2011). 'Current Major Cancer Targets for Nanoparticle Systems'. *Current Cancer Drug Targets*, 11(2): 164-183.
- White, M., Smith, A., Humphries, K., Pahl, S., Snelling, D. & Depledge, M. (2010) 'Blue space: The importance of water for preference, affect, and restorativeness ratings of natural and built scenes'. *Journal of Environmental Psychology*, 30 (4). pp 482-493.
- White, M. P. & Eiser, J. R. (2006) 'Marginal trust in risk managers: Building and losing trust following decisions under uncertainty'. *Risk analysis*, 26 (5). pp 1187-1203.
- White, M. P. & Johnson, B. B. (2010) 'The intuitive detection theorist (IDT) model of trust in hazard managers'. *Risk analysis*, 30 (8). pp 1196-1209.
- White, M. P., Pahl, S., Ashbullby, K., Herbert, S. & Depledge, M. H. (2013) 'Feelings of restoration from recent nature visits'. *Journal of Environmental Psychology*, 35. pp 40-51.
- Wick, P., M. J. D. Clift, M. Roeslein and B. Rothen-Rutishauser (2011). 'A brief summary of carbon nanotubes science and technology: a health and safety perspective'. *Chemsuschem* 4(7): 905-911.
- Widmer, W. M. & Reis, R. A. (2010) 'An experimental evaluation of the effectiveness of beach ashtrays in preventing marine contamination'. *Brazilian Archives of Biology and Technology*, 53 (5). pp 1205-1216.
- Wilber, R. J. (1987) 'Plastic in the North Atlantic'. *Oceanus*, 30 (3). pp 61-68.
- Wilcox, C., B. D. Hardesty, R. Sharples, D. A. Griffin, T. J. Lawson and R. Gunn (2013). 'Ghostnet impacts on globally threatened turtles, a spatial risk analysis for northern Australia'. *Conservation Letters*, 6(4): 247-254.
- Wilkinson, C., Allan, S., Anderson, A. & Petersen, A. (2007) 'From uncertainty to risk?: Scientific and news media portrayals of nanoparticle safety'. *Health, risk & society*, 9 (2). pp 145-157.
- Williams, P. R. & Hammitt, J. K. (2000) 'A comparison of organic and conventional fresh produce buyers in the Boston area'. *Risk analysis*, 20 (5). pp 735-746.
- Wilson, D. S. (1973) 'Food Size Selection Among Copepods'. *Ecology*, 54 (4). pp 909-914.

- Wilson, M. I., Robertson, L. D., Daly, M. & Walton, S. A. (1995) 'Effects of visual cues on assessment of water quality'. *Journal of Environmental Psychology*, 15 (1). pp 53-63.
- Wong, C. S., D. R. Green and W. J. Cretney (1974). 'Quantitative tar and plastic waste distributions in Pacific Ocean'. *Nature*, 247(5435): 30-32.
- Woodall, L. C., A. Sanchez-Vidal, M. Canals, G. L. J. Paterson, R. Coppock, V. Sleight, A. Calafat, A. D. Rogers, B. E. Narayanaswamy and R. C. Thompson (2014). The deep sea is a major sink for microplastic debris. *R. Soc. open sci.* 1: 140317. <http://dx.doi.org/10.1098/rsos.140317>
- World Health Organization [WHO] (2003) *Guidelines for safe recreational water environments. Volume 1: Coastal and fresh waters*. Geneva, Switzerland: WHO:
- World Tourism Organization (2013) *Annual Report 2012*. UNWTO, Madrid. Available.
- Wright, S. L., Rowe, D., Thompson, R. C. & Galloway, T. S. (2013a) 'Microplastic ingestion decreases energy reserves in marine worms'. *Current Biology*, 23 (23). pp R1031-R1033.
- Wright, S. L., Thompson, R. C. & Galloway, T. S. (2013b) 'The physical impacts of microplastics on marine organisms: A review'. *Environmental Pollution*, 178 (0). pp 483-492.
- Wyles, K. J., Pahl, S. & Thompson, R. C. (2014) 'Perceived risks and benefits of recreational visits to the marine environment: Integrating impacts on the environment and impacts on the visitor'. *Ocean & Coastal Management*, 88. pp 53-63.
- Wyles, K. J., Pahl, S., Thompson, R. C. & Thomas, K. (Under review) 'The detrimental effects of litter: Examining the impacts of different types of litter on psychological benefits and preferences'.
- Yakimets, I., D. W. Lai and M. Guigon (2004). 'Effect of photo-oxidation cracks on behaviour of thick polypropylene samples'. *Polymer Degradation and Stability* 86 (1): 59-67
- Yamashita, R., Takada, H., Fukuwaka, M.-a. & Watanuki, Y. (2011) 'Physical and chemical effects of ingested plastic debris on short-tailed shearwaters, *Puffinus tenuirostris*, in the North Pacific Ocean'. *Marine Pollution Bulletin*, 62 (12). pp 2845-2849.
- Yamashita, R. & Tanimura, A. (2007) 'Floating plastic in the Kuroshio Current area, western North Pacific Ocean'. *Marine Pollution Bulletin*, 54 (4). pp 485-488.
- Ye, S. & Andrady, A. L. (1991) 'Fouling of floating plastic debris under Biscayne Bay exposure conditions'. *Marine Pollution Bulletin*, 22 (12). pp 608-613.
- Zarfl, C., Fleet, D., Fries, E., Galgani, F., Gerdt, G., Hanke, G. & Matthies, M. (2011) 'Microplastics in oceans'. *Marine Pollution Bulletin*, 62 (8). pp 1589-1591.
- Zarfl, C. & Matthies, M. (2010) 'Are marine plastic particles transport vectors for organic pollutants to the Arctic?'. *Marine Pollution Bulletin*, 60 (10). pp 1810-1814.
- Zettler, E. R., Mincer, T. J. & Amaral-Zettler, L. A. (2013) 'Life in the "Plastisphere": Microbial Communities on Plastic Marine Debris'. *Environmental Science & Technology*, 47 (13). pp 7137- 7146.
- Zhao, S., Wang, H. & Jian, L. (2009) 'Influence of microplastics pollution on marine environment'. *Marine Sciences*, 33 (3). pp 84-86.
- Zimmerman, J. R., Ghosh, U., Millward, R. N., Bridges, T. S. & Luthy, R. G. (2004) 'Addition of carbon sorbents to reduce PCB and PAH bioavailability in marine sediments: Physicochemical tests'. *Environmental Science & Technology*, 38 (20). pp 5458-5464.
- Zlatev, M., Pahl, S. & White, M. (2010) 'Perceived risk and benefit for self and others as predictors of smokers' attitudes towards smoking restrictions'. *Psychology and Health*, 25 (2). pp 167-182.
- Zhou, P., C. Huang, H. Fang, W. Cai, D. Li, X. Li and H. Yu (2011). 'The abundance, composition and sources of marine debris in coastal seawaters or beaches around the northern South China Sea (China)'. *Marine Pollution Bulletin* 62(9). pp 1998-2007
- Zubris, K. A. V. & Richards, B. K. (2005) 'Synthetic fibers as an indicator of land application of sludge'. *Environmental Pollution*, 138. pp 201-211.

ANNEX I – MEMBERSHIP OF THE WORKING GROUP

Member	Affiliation
Peter Kershaw, chair	Independent consultant
Heather Leslie, co-chair	VU University, Amsterdam, Netherlands
Anthony Andrady	Independent consultant
Courtney Arthur	NOAA, United States
Joel Baker	Washington University, United States
Henk Bouwman	North West University, South Africa
Sarah Gall	Plymouth University, United Kingdom
Valeria Hidalgo-Ruz	Universidad Catolica del Norte, Chile
Angela Koehler	Alfred Wegener Institute, Germany
Kara Lavender Law	Sea Education Association, United States
Sabine Pahl	Plymouth University, United Kingdom
Jim Potemra	University of Hawaii, Hawaii
Peter Ryan	University of Cape Town, South Africa
Won Joon Shim	Korea Institute of Ocean Science & Technology, Republic of Korea
Hideshige Takada	Tokyo University of Agriculture & Technology, Japan
Richard Thompson	Plymouth University, United Kingdom
Alexander Turra	University of Sao Paulo, Brazil
Dick Vethaak	Deltares and VU University Amsterdam, Netherlands
Kayleigh Wyles	Plymouth Marine Institute, United Kingdom

Industry observers

Keith Christman, American Chemistry Council

Roberto Gomez, Plastics Europe

Ralph Schneider, Plastics Europe

Secretariat

Luis Valdes, IOC-UNESCO

Edward Kleverlaan, IMO

Fredrik Haag, IMO

Jennifer Rate, IMO

ANNEX II – LIST OF REPORTS AND STUDIES

The following reports and studies have been published so far. They are available from the GESAMP website: <http://gesamp.org>

1. Report of the seventh session, London, 24-30 April 1975. (1975). Rep. Stud. GESAMP, (1):pag.var. Available also in French, Spanish and Russian
2. Review of harmful substances. (1976). Rep. Stud. GESAMP, (2):80 p.
3. Scientific criteria for the selection of sites for dumping of wastes into the sea. (1975). Rep. Stud. GESAMP, (3):21 p. Available also in French, Spanish and Russian
4. Report of the eighth session, Rome, 21-27 April 1976. (1976). Rep. Stud. GESAMP, (4):pag.var. Available also in French and Russian
5. Principles for developing coastal water quality criteria. (1976). Rep. Stud. GESAMP, (5):23 p.
6. Impact of oil on the marine environment. (1977). Rep. Stud. GESAMP, (6):250 p.
7. Scientific aspects of pollution arising from the exploration and exploitation of the sea-bed. (1977). Rep. Stud. GESAMP, (7):37 p.
8. Report of the ninth session, New York, 7-11 March 1977. (1977). Rep. Stud. GESAMP, (8):33 p. Available also in French and Russian
9. Report of the tenth session, Paris, 29 May - 2 June 1978. (1978). Rep. Stud. GESAMP, (9):pag.var. Available also in French, Spanish and Russian
10. Report of the eleventh session, Dubrovnik, 25-29 February 1980. (1980). Rep. Stud. GESAMP, (10):pag.var. Available also in French and Spanish
11. Marine Pollution implications of coastal area development. (1980). Rep. Stud. GESAMP, (11):114 p.
12. Monitoring biological variables related to marine pollution. (1980). Rep. Stud. GESAMP, (12):22 p. Available also in Russian
13. Interchange of pollutants between the atmosphere and the oceans. (1980). Rep. Stud. GESAMP, (13):55 p.
14. Report of the twelfth session, Geneva, 22-29 October 1981. (1981). Rep. Stud. GESAMP, (14):pag.var. Available also in French, Spanish and Russian
15. The review of the health of the oceans. (1982). Rep. Stud. GESAMP, (15):108 p.
16. Scientific criteria for the selection of waste disposal sites at sea. (1982). Rep. Stud. GESAMP, (16):60 p.
17. The evaluation of the hazards of harmful substances carried by ships. (1982). Rep. Stud. GESAMP, (17):pag.var.
18. Report of the thirteenth session, Geneva, 28 February - 4 March 1983. (1983). Rep. Stud. GESAMP, (18):50 p. Available also in French, Spanish and Russian
19. An oceanographic model for the dispersion of wastes disposed of in the deep sea. (1983). Rep. Stud. GESAMP, (19):182 p.
20. Marine pollution implications of ocean energy development. (1984). Rep. Stud. GESAMP, (20):44 p.
21. Report of the fourteenth session, Vienna, 26-30 March 1984. (1984). Rep. Stud. GESAMP, (21):42 p. Available also in French, Spanish and Russian
22. Review of potentially harmful substances. Cadmium, lead and tin. (1985). Rep. Stud. GESAMP, (22):114 p.
23. Interchange of pollutants between the atmosphere and the oceans (part II). (1985). Rep. Stud. GESAMP, (23):55 p.
24. Thermal discharges in the marine environment. (1984). Rep. Stud. GESAMP, (24):44 p.
25. Report of the fifteenth session, New York, 25-29 March 1985. (1985). Rep. Stud. GESAMP, (25):49 p. Available also in French, Spanish and Russian.

26. Atmospheric transport of contaminants into the Mediterranean region. (1985). Rep. Stud. GESAMP, (26):53 p.
27. Report of the sixteenth session, London, 17-21 March 1986. (1986). Rep. Stud. GESAMP, (27):74 p. Available also in French, Spanish and Russian
28. Review of potentially harmful substances. Arsenic, mercury and selenium. (1986). Rep. Stud. GESAMP, (28):172 p.
29. Review of potentially harmful substances. Organosilicon compounds (silanes and siloxanes). (1986). Published as UNEP Reg. Seas Rep. Stud., (78):24 p.
30. Environmental capacity. An approach to marine pollution prevention. (1986). Rep. Stud. GESAMP, (30):49 p.
31. Report of the seventeenth session, Rome, 30 March - 3 April 1987. (1987). Rep. Stud. GESAMP, (31):36 p. Available also in French, Spanish and Russian
32. Land-sea boundary flux of contaminants: contributions from rivers. (1987). Rep. Stud. GESAMP, (32):172 p.
33. Report on the eighteenth session, Paris, 11-15 April 1988. (1988). Rep. Stud. GESAMP, (33):56 p. Available also in French, Spanish and Russian
34. Review of potentially harmful substances. Nutrients. (1990). Rep. Stud. GESAMP, (34):40 p.
35. The evaluation of the hazards of harmful substances carried by ships: Revision of GESAMP Reports and Studies No. 17. (1989). Rep. Stud. GESAMP, (35):pag.var.
36. Pollutant modification of atmospheric and oceanic processes and climate: some aspects of the problem. (1989). Rep. Stud. GESAMP, (36):35 p.
37. Report of the nineteenth session, Athens, 8-12 May 1989. (1989). Rep. Stud. GESAMP, (37):47 p. Available also in French, Spanish and Russian
38. Atmospheric input of trace species to the world ocean. (1989). Rep. Stud. GESAMP, (38):111 p.
39. The state of the marine environment. (1990). Rep. Stud. GESAMP, (39):111 p. Available also in Spanish as Inf.Estud.Progr.Mar.Reg.PNUMA, (115):87 p.
40. Long-term consequences of low-level marine contamination: An analytical approach. (1989). Rep. Stud. GESAMP, (40):14 p.
41. Report of the twentieth session, Geneva, 7-11 May 1990. (1990). Rep. Stud. GESAMP, (41):32 p. Available also in French, Spanish and Russian
42. Review of potentially harmful substances. Choosing priority organochlorines for marine hazard assessment. (1990). Rep. Stud. GESAMP, (42):10 p.
43. Coastal modelling. (1991). Rep. Stud. GESAMP, (43):187 p.
44. Report of the twenty-first session, London, 18-22 February 1991. (1991). Rep. Stud. GESAMP, (44):53 p. Available also in French, Spanish and Russian
45. Global strategies for marine environmental protection. (1991). Rep. Stud. GESAMP, (45):34 p.
46. Review of potentially harmful substances. Carcinogens: their significance as marine pollutants. (1991). Rep. Stud. GESAMP, (46):56 p.
47. Reducing environmental impacts of coastal aquaculture. (1991). Rep. Stud. GESAMP, (47):35 p.
48. Global changes and the air-sea exchange of chemicals. (1991). Rep. Stud. GESAMP, (48):69 p.
49. Report of the twenty-second session, Vienna, 9-13 February 1992. (1992). Rep. Stud. GESAMP, (49):56 p. Available also in French, Spanish and Russian
50. Impact of oil, individual hydrocarbons and related chemicals on the marine environment, including used lubricant oils, oil spill control agents and chemicals used offshore. (1993). Rep. Stud. GESAMP, (50):178 p.
51. Report of the twenty-third session, London, 19-23 April 1993. (1993). Rep. Stud. GESAMP, (51):41 p. Available also in French, Spanish and Russian

52. Anthropogenic influences on sediment discharge to the coastal zone and environmental consequences. (1994). Rep. Stud. GESAMP, (52):67 p.
53. Report of the twenty-fourth session, New York, 21-25 March 1994. (1994). Rep. Stud. GESAMP, (53):56 p. Available also in French, Spanish and Russian
54. Guidelines for marine environmental assessment. (1994). Rep. Stud. GESAMP, (54):28 p.
55. Biological indicators and their use in the measurement of the condition of the marine environment. (1995). Rep. Stud. GESAMP, (55):56 p. Available also in Russian
56. Report of the twenty-fifth session, Rome, 24-28 April 1995. (1995). Rep. Stud. GESAMP, (56):54 p. Available also in French, Spanish and Russian
57. Monitoring of ecological effects of coastal aquaculture wastes. (1996). Rep. Stud. GESAMP, (57):45 p.
58. The invasion of the ctenophore *Mnemiopsis leidyi* in the Black Sea. (1997). Rep. Stud. GESAMP, (58):84 p.
59. The sea-surface microlayer and its role in global change. (1995). Rep. Stud. GESAMP, (59):76 p.
60. Report of the twenty-sixth session, Paris, 25-29 March 1996. (1996). Rep. Stud. GESAMP, (60):29 p. Available also in French, Spanish and Russian
61. The contributions of science to integrated coastal management. (1996). Rep. Stud. GESAMP, (61):66 p.
62. Marine biodiversity: patterns, threats and development of a strategy for conservation. (1997). Rep. Stud. GESAMP, (62):24 p.
63. Report of the twenty-seventh session, Nairobi, 14-18 April 1997. (1997). Rep. Stud. GESAMP, (63):45 p. Available also in French, Spanish and Russian
64. The revised GESAMP hazard evaluation procedure for chemical substances carried by ships. (2002). Rep. Stud. GESAMP, (64):121 p.
65. Towards safe and effective use of chemicals in coastal aquaculture. (1997). Rep. Stud. GESAMP, (65):40 p.
66. Report of the twenty-eighth session, Geneva, 20-24 April 1998. (1998). Rep. Stud. GESAMP, (66):44 p.
67. Report of the twenty-ninth session, London, 23-26 August 1999. (1999). Rep. Stud. GESAMP, (67):44 p.
68. Planning and management for sustainable coastal aquaculture development. (2001). Rep. Stud. GESAMP, (68):90 p.
69. Report of the thirtieth session, Monaco, 22-26 May 2000. (2000). Rep. Stud. GESAMP, (69):52 p.
70. A sea of troubles. (2001). Rep. Stud. GESAMP, (70):35 p.
71. Protecting the oceans from land-based activities - Land-based sources and activities affecting the quality and uses of the marine, coastal and associated freshwater environment.(2001). Rep. Stud. GESAMP, (71):162p.
72. Report of the thirty-first session, New York, 13-17 August 2001. (2002). Rep. Stud. GESAMP, (72):41 p.
73. Report of the thirty-second session, London, 6-10 May 2002. (in preparation). Rep. Stud. GESAMP, (73)
74. Report of the thirty-third session, Rome, 5-9 May 2003 (2003) Rep. Stud. GESAMP, (74):36 p.
75. Estimations of oil entering the marine environment from sea-based activities (2007), Rep. Stud. GESAMP, (75):96 p.
76. Assessment and communication of risks in coastal aquaculture (2008). Rep. Stud. GESAMP, (76):198 p.
77. Report of the thirty-fourth session, Paris, 8-11 May 2007 (2008), Rep. Stud. GESAMP, (77):83 p.
78. Report of the thirty-fifth session, Accra, 13-16 May 2008 (2009), Rep. Stud. GESAMP, (78):73 p.
79. Pollution in the open oceans: a review of assessments and related studies (2009). Rep. Stud. GESAMP, (79):64 p.
80. Report of the thirty-sixth session, Geneva, 28 April - 1 May 2009 (2011), Rep. Stud. GESAMP, (80):83 p.
81. Report of the thirty-seventh session, Bangkok, 15 - 19 February 2010 (2010), Rep. Stud. GESAMP, (81):74 p.

82. Proceedings of the GESAMP International Workshop on Micro-plastic Particles as a Vector in Transporting Persistent, Bio-accumulating and Toxic Substances in the Oceans (2010). Rep. Stud. GESAMP, (82):36 p.
83. Establishing Equivalency in the Performance Testing and Compliance Monitoring of Emerging Alternative Ballast Water Management Systems (EABWMS). A Technical Review. Rep. Stud. GESAMP, (83):63 p, GloBallast Monographs No. 20.
84. The Atmospheric Input of Chemicals to the Ocean (2012). Rep. Stud. GESAMP, (84) GAW Report No. 203.
85. Report of the 38th Session, Monaco, 9 to 13 May 2011 (pre-publication copy), Rep. Stud. GESAMP, (85): 118 p.
86. Report of Working Group 37: Mercury in the Marine Environment (in prep.). Rep. Stud. GESAMP, (86).
87. Report of the 39th Session, New York, 15 to 20 April 2012 (pre-publication copy), Rep. Stud. GESAMP, (87):92 p.
88. Report of the 40th Session, Vienna, 9 to 13 September 2013, Rep. Stud. GESAMP, (88):86 p.
89. Report of the 41st Session, Malmö, 1 to 4 September 2014, Rep. Stud. GESAMP, (89): in prep.
90. Report of Working Group 40: Sources, fate and effects of microplastics in the marine environment: a global assessment. Rep. Stud. GESAMP, (90):96 p.



Science for Sustainable Oceans

ISSN 1020-4873