

## Chapter 5

# Effects on Terrestrial Ecosystems

John Derome, Sirkku Manninen, Julian Aherne, Paavo Hellstedt, Jean-Paul Hettelingh, Kevin Hicks, Satu Huttunen, Juha Kämäri, Galina Kashulina, Mikhail Kozlov, Annamari Markkola, Maximilian Posch, Anna-Liisa Ruotsalainen, Reijo Salminen, and Elena Zvereva

The first AMAP assessment of acidification (AMAP, 1998) addressed the processes involved in the acidification of arctic soils, and the direct effects of sulfur dioxide ( $\text{SO}_2$ ) and nitrogen oxides ( $\text{NO}_x$ ), as well as acidifying deposition, on the biotic components of terrestrial ecosystems. At that time, however, there was little empirical evidence to suggest that soil acidification was anything more than a local problem in very limited parts of, for example, the Kola Peninsula. The serious damage to flora and fauna reported in the area was mainly attributed to the direct toxic effects of  $\text{SO}_2$ , combined with the accumulation of toxic heavy metals in the arctic environment. It is extremely difficult to distinguish between the direct effects of  $\text{SO}_2$ , the indirect effects of the deposition of acidifying compounds ( $\text{SO}_2$  and sulfate ( $\text{SO}_4$ )), and the direct toxic effects of heavy metals on terrestrial ecosystems in the areas around the smelters. In the Arctic, the cumulative effects of acidifying emissions and the deposition of toxic heavy metals can be disastrous for ecosystems which are already subject to extreme climatic conditions. In this, the latest acidification assessment, a large amount of new empirical data are presented and discussed.

### 5.1. Effects on soils

Kuylenstierna *et al.* (2001) have mapped the relative sensitivity of terrestrial ecosystems to acidic deposition at the global scale. An overview of the sensitivity of arctic ecosystems is provided in a circumpolar map extracted from the global sensitivity map. The sensitivity classes are based on base saturation and cation exchange capacity data applied to FAO (Food and Agriculture Organization) soil types. The soils most sensitive to acidic deposition have a low base saturation (<40%) and a low cation exchange capacity (<10 meq/100 g); the implication is that these soils have low rates of long-term mineral weathering and a limited base cation content and may therefore be subject to rapid changes in base saturation and pH. The most sensitive areas occur in Fennoscandia, in parts of Russia, and in parts of Canada and Alaska. Much of the Canadian Arctic seems not to be sensitive to acidic deposition (Figure 5.1).

AMAP has identified three regions in the Arctic that may be susceptible to acidification caused by the deposition of acidifying compounds (sulfur and nitrogen): the Kola Peninsula in northwestern Russia, the Taymir Peninsula in northern Russia, and the Chukotka region in eastern Siberia. The Kola Peninsula and the Taymir Peninsula both receive acidifying compounds from major local point sources; deposition in these areas has been, and is still, high. There is a large amount of information about the effects of acidic deposition on soils on the Kola Peninsula but much less about effects on soils in the Norilsk area. Owing to the very high  $\text{SO}_2$  emissions from the Norilsk smelter complex, the acidifying effects are potentially many times higher than on the Kola Peninsula. Although the Chukotka region may receive large amounts of acidifying pollutants




Figure 5.1. Sensitivity of arctic ecosystems to acid deposition (Kuylenstierna *et al.*, 2001).

via long-range transport from industrial sources in China, India, and other parts of the Far-East, reliable deposition data and information about the soil acidification status of the region are extremely difficult to obtain. Other parts of the Arctic receive acidifying compounds via long-range transport from North America, Europe and the Far-East but, according to deposition measurements made in the Arctic, the levels are not likely to cause widespread soil acidification in background areas in the near future.

#### 5.1.1. Acidity status of soils on the Kola Peninsula

The results of two regional studies – the Kola Ecogeochemistry Project ([www.ngu.no/Kola](http://www.ngu.no/Kola)) and the Barents Ecogeochemistry Project ([www.gsf.fi/Barents](http://www.gsf.fi/Barents)) – have recently become available. These cover the distribution area of  $\text{SO}_2$  emissions from the Cu-Ni smelters in Nikel, Zapolyarnyy and Monchegorsk on the Kola Peninsula, and almost pristine areas of northern Finland (Reimann *et al.*, 1998a; Salminen *et al.*, 2004). The survey carried out as part of the Kola Ecogeochemistry Project had a sampling density of approximately 1 site per 300 km<sup>2</sup> and generated information about soil acidity status and the most important natural and anthropogenic factors contributing to soil acidity in this region (Reimann *et al.*, 1998a, 2000a; Kashulina *et al.*, 1998a,b, 2003; Kashulina and Reimann,

2001; Kashulina, 2002). Figure 5.2 shows the area discussed here. The results of the Barents Ecogeochemistry Project, undertaken five years later and with a lower sampling density, confirmed the overall conclusions of the Kola Ecogeochemistry Project, and in addition, provided information about temporal trends in soil acidity in the region.

According to the Kola Ecogeochemistry Project, the pH(H<sub>2</sub>O) of the organic (O) horizon of podzolic soils in 1995 ranged from 3.2 to 5.6. Although the region has received high levels of acidifying deposition for about 60 years, this wide range in pH cannot be related to the effects of two of the world's largest SO<sub>2</sub> emission sources, which are located on the Kola Peninsula, since an even greater pH range (3 to 6) has been reported as typical for the O-horizon of podzols in background areas (Targulian, 1971).

#### 5.1.1.1. Natural factors affecting soil acidity

##### *Marine input of base cations*

There is a relatively strong decreasing trend (0.6 pH units) in the acidity of the O-horizon along a south–north gradient running through the background area (from the Arctic Circle to the coast of the Barents Sea) of northern Finland (Figure 5.3) (Reimann *et al.*, 2000b; Kashulina *et al.*, 2003). The sharpest increase in pH (0.4 pH units) occurs on mov-

ing north through the 100 km wide coastal zone. There is a 10-fold increase in sodium (Na) and a 2-fold increase in magnesium (Mg) in ground mosses and the O-horizon over the same coastal zone (Figure 5.3). This indicates that marine aerosols are probably the main factor explaining the pH increase near the coast. The figure also indicates that natural precipitation chemistry can have a significant effect on acidity parameters in the O-horizon.

##### *Bioclimatic factors*

The distribution of Na concentrations (Na is a major element in marine aerosols, but not an important plant nutrient) in mosses and the O-horizon (Figure 5.3) suggests, however, that the influence of marine aerosols is only responsible for the south–north pH gradient in the O-horizon within a distance of 200 km from the coast. The minor increase in pH between 200 and 500 km from the coast coincides with a slight increase in Mg and calcium (Ca) concentrations (Mg and Ca are both important plant nutrients) (Figure 5.3). The southward decrease in exchangeable Ca and especially Mg concentrations in the O-horizon is probably due to increasing uptake and utilization by plants. Thus, the change in bioclimatic factors (climate + vegetation) on moving from south to north may also affect the distribution of acidity in the O-horizon in the region (Kashulina *et al.*, 2003).



Figure 5.2. The Kola Ecogeochemistry survey area and major industrial centers. The lines A–B, C–D, and E–F mark the transects shown in Figures 5.3, 5.4, 5.5, 5.7, and 5.9.

### Site-specific and temporal variation

All the soil acidity parameters showed very high site-specific variation (Figure 5.3). For instance, pH at two adjacent sites in background areas differed by more than one pH unit. Acidity in the O-horizon can vary within the same site by more than one pH unit over the year (Levina, 1969). Thus, spatial and temporal variation in pH in the O-horizon may be greater than any natural trends at the regional level. Site-specific characteristics and/or processes can have a stronger deterministic effect on soil acidity status at an individual site than any other regional factors (Kashulina *et al.*, 2003).

#### 5.1.1.2. Sulfur dioxide emissions and soil acidity

The O-horizon of podzols is directly influenced by the input of ions and cations in precipitation. In the study area the organic material in the O-horizon has a lifetime of around 20 to 50 years. As a result, it 'integrates' the effects of pollutant deposition over a relatively long period, and the cumulative effect of acidifying pollutants can be considerable. According to Kashulina *et al.* (2003), precipitation acidity is affected by SO<sub>2</sub> emissions only within a radius of 30 km around the smelters. In other parts of the Kola Peninsula, the emissions of base cations (derived from the smelters, marine aerosols, and basal and alkaline dust

from other industrial sources) supply enough base cations to maintain precipitation at a less acidic level than that recorded in northern Finland. Current emissions of nitrogen dioxide on the Kola Peninsula are low and have little or no effect on precipitation acidity; nitrate concentrations in precipitation are low and relatively uniform throughout the area (Tikkanen and Niemelä, 1995; Kashulina *et al.*, 1998a). Thus, the area where the soil can be affected by acidified rain on the Kola Peninsula is limited to restricted zones around the smelters. The contribution played by the direct adsorption of SO<sub>2</sub> on the soil surface on the Kola Peninsula needs to be investigated, however.

#### Monchegorsk area

The west-east transect of pH in the O-horizon, running through the area occupied by the Monchegorsk smelter (Figure 5.4), indicates that emissions from industrial activities at Apatity (mining and processing of alkaline rocks mined at Kirovsk near Apatity), located 40 km south-east of Monchegorsk, have increased the pH by 0.3 units. These emissions also affect the pH of the surface soil close to the Monchegorsk smelter, and are still detectable at the Russian/Finnish border to the west. The atmospheric origin of this increase in pH is supported by the distribution of base cations in moss (Figure 5.4).




Figure 5.3. Transect south-north across the survey area near the western boundary in 1995 (C–D on Figure 5.2), showing: (a) pH in a water extract of the O-horizon; (b) ammonium acetate extractable (pH 4.5) base cation concentrations in the O-horizon; and (c) total base cation concentrations in mosses (Kola Ecogeochemistry Project, Kashulina *et al.*, 2003).




Figure 5.4. Transect west-east across the survey area and through the industrial zone centered on Monchegorsk and Apatity in 1995 (A–B on Figure 5.2), showing: (a) pH in a water extract of the O-horizon of podzol soils; (b) exchangeable base cation concentrations in the O-horizon of podzol soils; and (c) total base cation concentrations in mosses (Kola Ecogeochemistry Project, Kashulina *et al.*, 2003).



The distribution of exchangeable Ca (the dominant base cation) in the O-horizon along the transect follows the same pattern as for pH (Figure 5.4). However, there is no corresponding trend for potassium (K) or Mg. This could be partly due to the increased leaching into stream water, observed within 30 km of Monchegorsk (Kashulina *et al.*, 2003). However, the increase in total Mg in mosses (Figure 5.4) and the O-horizon near Monchegorsk (Reimann *et al.*, 1998a) is in agreement with the increased deposition of Mg in the area. The decrease in the exchangeable Mg concentration may be associated with low solubility of Mg in anthropogenic particulate material deposited on the O-horizon.

A low pH (although within the natural range of pH variation) and extremely low base cation concentrations (Figure 5.4) are characteristic of soils at a number of sampling sites in the immediate vicinity of the Monchegorsk smelter where the vegetation cover has been completely destroyed. Low pH values were also reported in case or gradient studies within a distance of 5 km from the smelters on the Kola Peninsula (Chertov *et al.*, 1993; Koptsik and Muchina, 1995; Kashulina *et al.*, 2003). No changes in soil acidity were found in a gradient study within a distance of 10 to 80 km from Monchegorsk (Tikkanen and Niemelä, 1995).

#### Nikel/Zapolyarnyy area

Further north, the variation in pH in the O-horizon along an east–west transect through Nickel and Zapolyarnyy is less than 0.2 pH units (Figure 5.5). There is a very slight (<0.1 pH units) decreasing trend near Nickel, due to a small number of low pH values near Nickel. Correspondingly, the slightly increasing trend near Zapolyarnyy is due to a number of relatively high pH values. The eastern end of the transect has slightly higher pH values than the western end. The distribution of marine-derived cations (Na and Mg) in mosses and the O-horizon indicates an increasing maritime influence towards the east, starting near Nickel (Figure 5.5). The western section of the transect is more inland, while the eastern section ends at the Barents Sea coast. In addition to marine-derived deposition, the Zapolyarnyy area receives alkaline dust from opencast mining (Reimann *et al.*, 1997).

Generally, it appears that the sources of base cations are sufficient to prevent acidification of the O-horizon, and to maintain relatively constant pH and base cation concentrations in the vicinity of Nickel and Zapolyarnyy. As is the case for Monchegorsk, only a small number of low values were observed in the immediate vicinity of Nickel/Zapolyarnyy, especially at sites where the vegetation cover has been severely damaged.

#### 5.1.1.3. The role of overburden and bedrock chemistry

The simultaneous emission of fly ash by the smelters and associated power plants, and the alkaline nature of the overburden and bedrock in some areas near the emission sources, are frequently used to explain the lack of widespread soil acidification on the Kola Peninsula, despite the very high SO<sub>2</sub> emissions (Koptsik and Muchina, 1995; Moiseenko, 1997; Tikkanen and Niemelä, 1995). This raises the question of whether the relatively high base cation content of the parent material in some areas near both emission sources (Monchegorsk and Nickel/Zapolyarnyy) can influence the chemistry of the O-horizon and counteract the acidifying effect of the emissions. Relatively poor correlation has been found between the element concentrations in the parent material (C-horizon) and the O-horizon immediately around the Nickel/Zapolyarnyy smelters (Reimann *et al.*, 1998b). Base cation concentrations in the O-horizon at all the sampling points show a much closer relationship with the corresponding concentrations in mosses (the chemistry of which is primarily determined by atmospheric inputs) (Figures 5.3 to 5.6), than with those in the C-horizon (Kashulina *et al.*, 2003). Thus, the distribution of acidity in the O-horizon in the region is mainly determined by the atmospheric input, and the geological influence is hardly detectable in the relatively high atmospheric base cation deposition gradient on the Kola Peninsula. Fly ash from the smelter complexes is not the most important source of base




Figure 5.5. Transect west–east across the survey area and through the industrial zone centered on Nickel and Zapolyarnyy in 1995 (E–F on Figure 5.2), showing (a) pH in a water extract of the O-horizon of podzol soils; (b) exchangeable base cation concentrations in the O-horizon of podzol soils; and (c) total base cation concentrations in moss (Kola Ecogeochemistry project, Kashulina *et al.*, 2003).




Figure 5.6. Ammonium acetate extractable Ca and Mg in the O-horizon of soils (x-axis) vs. their total concentrations (aqua regia extractable) in the C-horizon of soils and in moss in 1995 (Kola Ecogeochemistry project, Kashulina *et al.*, 2003).

cations in deposition on the Kola Peninsula (Kashulina *et al.*, 2003). Alkaline dust from the apatite fertilizer plant, marine aerosols, and alkaline dust from open-cast mining appear to be the major sources of base cation deposition on forest soil in the western part of the Kola Peninsula. The current rate of base cation emissions is sufficient to maintain soil acidity parameters at levels that are even less acidic than, for example, in northern Finland. The only areas with increased soil acidity and exchangeable base cation depletion are in the severely damaged ecosystems immediately around the smelters.

#### Exchangeable aluminum

The major factor governing the distribution of exchangeable aluminum (Al) in the O-horizon on the Kola Peninsula appears to be the bedrock geology (Kashulina *et al.*, 2003). Aluminum concentrations in the C-horizon along the transect running through the major SO<sub>2</sub> emission source areas (Figure 5.7) are more similar to the concentrations in the O-horizon than to the moss concentrations (Figure 5.4). The Al concentrations in mosses along the transect

do, however, suggest that there is elevated deposition of Al into the O-horizon near Nikel/Zapolyarnyy (alkaline rock dust from the opencast mine) and near Apatity (apatite-nepheline dust). The Al concentrations in precipitation increase to such an extent immediately around the smelters (Reimann *et al.*, 1997; de Caritat *et al.*, 1998), that Al is a major cation (in equivalent terms) in rain and snow (Kashulina *et al.*, 2003). Thus, the relatively high exchangeable Al concentrations in the soil and percolation water reported near Monchegorsk (Motova and Nikonov, 1993; Lukina and Nikonov, 1996) are more likely to be associated with basal lithology and increased deposition than to the impacts of SO<sub>2</sub> emissions.

The aluminum:base cation (Al:BC) ratio in the O-horizon (Figure 5.7) shows extremely high spatial variation. The highest values occur around Apatity (apatite-nepheline, syenite) and in soils overlying the meta-sedimentary rocks of northern Norway. The Al:BC ratio in the O-horizon shows a good correlation with the total Al concentration in the C-horizon ( $r = 0.73$ ), and so the geology is therefore the main factor determining the Al:BC ratio in the O-horizon




Figure 5.7. A comparison of the two west-east transects across the survey area in 1995 (A–B and E–F on Figure 5.2), showing changes from east to west in (a) total Al (nitric acid) and exchangeable Al concentrations in the O-horizon; (b) total Al concentrations in moss and Al (aqua regia extracts) concentrations in the C-horizon; and (c) the exchangeable Al:(Ca+Mg+K+Na) ratio in the O-horizon (Kola Ecogeochemistry Project, Kashulina *et al.*, 2003).

(Kashulina *et al.*, 2003). Emissions affect the Al:BC ratio in the soil only within the immediate vicinity of the smelters. Furthermore, this appears to be an indirect effect of pollution via damage to the ecosystem. Elevated Al:BC ratios also occur in the completely destroyed ecosystems (Figure 5.8).

#### 5.1.1.4. Connections between soil condition and ecosystem quality

##### Soil acidity

The distribution of the major acidity parameters in the various classes of ecosystem damage (Kashulina *et al.*, 2003)




Figure 5.8. Cumulative distribution functions (CDF, %) of pH (water extraction), exchangeable Ca, Mg, K, and Al concentrations and the exchangeable Al:(Ca+Mg+K+Na) ratio in the O-horizon in the 'no visual damage' zone in Finnish Lapland and for various classes of ecosystem damage in the western part of the Kola Peninsula in 1995 (Kola Ecogeochemistry Project, Kashulina *et al.*, 2003).

shows that the onset of ecosystem damage (and even of serious ecosystem damage) on the Kola Peninsula occurs where there is a more favorable (higher pH and base cation concentrations) acidity status in the O-horizon compared with background areas in northern Finland (Figure 5.8). In contrast, severe ecosystem damage appears to be a prerequisite for soil acidification to appear on the Kola Peninsula.

#### Plant nutrients

In addition to base cations (Ca, Mg, K) and other important nutrients (e.g., nitrogen and phosphorus), depletion of some other nutrients (e.g., manganese (Mn) and zinc (Zn)) is regarded as one of the detrimental consequences of acidifying deposition on ecosystems (Galloway, 1995). A sharp decrease in some nutrients (Mn, Zn, Mg) has been reported in severely damaged ecosystems around the smelters on the Kola Peninsula (Lobersli and Venn, 1995; Tikkanen and

Niemelä, 1995). A deterioration in the nutrient status of the soil on the Kola Peninsula has occurred only in the completely destroyed ecosystems around the smelters. There is also a decrease in phosphorus concentrations in the seriously damaged ecosystems within a radius of 30 km around the smelters. Manganese concentrations below the deficiency level only occurred at sites with serious or complete destruction of the ecosystems (Kashulina, 2002). Furthermore, the depletion of major nutrients is not a specific reaction of the ecosystem to acidifying deposition.

Thus, the indirect effect of acidifying pollutants on the condition of ecosystems via soil acidification and nutrient depletion does not appear to be the main factor affecting forest ecosystem condition on the Kola Peninsula. The direct effects of SO<sub>2</sub> on the vegetation (Aamlid *et al.*, 1995; Tikkanen and Niemelä, 1995; Kashulina *et al.*, 2003) is a much more likely explanation for the widespread damage to ecosystems on the Kola Peninsula. The significant



increases in the concentrations of some other pollutants (e.g., nickel, cobalt, cadmium, silver, arsenic, lead, copper and others) of up to a few orders of magnitude in the soil near the major emission sources (Räisänen *et al.*, 1997; Reimann *et al.*, 1998a; Åyräs and Kashulina, 2000) is obviously contributing to the widespread ecosystem damage.

#### 5.1.1.5. Temporal trends in soil acidity

A comparison of soil acidity parameters near the Monchegorsk smelter in 1995 (Reimann *et al.*, 1997; Kashulina *et al.*, 2003) and 2000–2001 (Salminen *et al.*, 2004) showed no clear changes in the exchangeable Ca concentration in the O-horizon (Figure 5.9). The values obtained in the sparser sampling network in 2000 are within the range of spatial variability reported for 1995.

#### 5.1.2. Acidification and the acidity status of soils in the Norilsk area

The Norilsk mining area is one of the largest point sources of sulfur and certain heavy metals in the northern Arctic (AMAP, 1998). Annual sulfur emissions were around 1 million tonnes between 1985 and 2000 (Ekimov *et al.*, 2001), while annual nickel and copper emissions were up to 1300 and 2800 tonnes, respectively. These emissions are more than ten times higher than emissions reported for the Monchegorsk smelter on the Kola Peninsula (MRCENR, 1995; Ekimov *et al.*, 2001). Since 2000, there have been no reported changes in the volume of metal production or any investments to decrease emissions. SO<sub>2</sub> emissions for Norilsk were 2118 kt for 1992 and 1847 kt for 2003.

The effects of emissions on the soil are partly determined by climate. Owing to the presence of permafrost in the region, chemical processes can only occur in the uppermost part of the soil, and because of the long winter (9 to 10 months) the period favorable for chemical changes is short. However, the uppermost part of the soil profile is strongly affected by the emissions of sulfur and heavy metals.

Basalts with a basic chemical composition characterize the bedrock in the Norilsk area and in the neighboring Putorana Mountains. According to Naldrett *et al.* (1992), the sulfide ore deposits in the Norilsk area are hosted by a large flood basalt formation that erupted in the Devonian and Carboniferous sedimentary rocks. The age of the basalt is 251 million years. It consists of picritic, basaltic, and tholeiitic lavas. Today the basalts form the mountains (the Putorana Mountains) and sedimentary rocks (which are not metamorphosed but eroded deeper) fill the valleys (e.g., in the Norilsk area). These sedimentary rocks include

Devonian calcareous and dolomitic marls, and sulfate-rich evaporates and lower Carboniferous shallow water limestones. Westward from Norilsk the bedrock changes and mainly includes intermediate and more acidic types of rock.

In 2001, water and soil samples were collected from ten sites in the Norilsk industrial area and in the uncontaminated area to the east of the industrial zone. Stream water samples were collected from each site and complete soil profiles were collected from two sites – one in the industrial area and the other in the clean background area. Organic layer samples were also taken from two sites. The sampling and analytical procedures are as reported by Gregorauskiene *et al.* (2000) and Salminen and Gregorauskiene (2002).

The minerogenic soils that have developed on the weathering crust of basalt (Lake Lama) have pH values ranging from 5.1 to 6.2, and the soils that have developed on calcareous sedimentary rocks (Norilsk) from 7.1 to 8.6. These values are considerably higher than the respective values reported for Monchegorsk (Table 5.1). The buffering capacity, as indicated by Ca and Mg concentrations, is much higher in Norilsk than in Monchegorsk. The soils in the Norilsk area cannot be considered sensitive to acidification; the pH of the organic layer in the Norilsk industrial area was as high as 6.4. However, the Cu and Ni values of the organic layer are at the same level as in Monchegorsk (1370–2820 mg/kg), due to the smelter emissions. The corresponding concentrations in the Lake Lama area are much lower (72–82 mg/kg). However, even the concentrations in the Lake Lama area are higher than the median values (5.9–7.9 mg/kg) reported for large areas of the Kola Peninsula (Salminen *et al.*, 2004).

#### 5.1.3. Effects on soil micro-organisms

Soil acidification and the deposition of sulfur and heavy metals can influence soil micro-organisms via several routes. First, a decline in pH and the accumulation of pollutants in the soil may directly reduce microbial growth and activity (Bååth, 1989). Second, pollutants may indirectly harm the symbiotic and rhizosphere micro-organisms by decreasing the amount of photosynthesizing foliage of their hosts. This, in turn, reduces carbon flow to the roots and mycorrhizal fungi (which are dependent on carbon received from the host) and to rhizosphere microbes (which are negatively affected both by a reduction in the availability of suitable environment due to reduced root biomass and by a decline in the amount of carbohydrate exudates produced by the roots). Direct and indirect impacts of acidification




Figure 5.9. Transect west–east across the survey area (A–B on Figure 5.2) showing exchangeable Ca in the O-horizon of podzol soils in 1995 (Kola Ecogeochemistry Project) and from 2000 to 2001 (Barents Ecogeochemistry Project).



Table 5.1. Element concentrations in soil horizons in the Monchegorsk (Kashulina and Gregorauskiene, 2000), Norilsk, and Lake Lama areas.

Sites/ horizon	Lowest depth (cm)	pH	LOI (%)	C (%)	Ca (mg/kg)		Mg (mg/kg)		Cu (mg/kg)	Ni (mg/kg)	S (mg/kg)
					Total	AR	Total	AR	AR	AR	Total
Monchegorsk											
Oer	0	3.90	20.50	11.60		792		532	1370	2030	446
OEer	3	4.39	5.79	3.11	39800	614	37300	443	366	541	122
Bs	20				42400						
BC1	40	4.78	3.72	0.92	49900	3310	48200	4850	52.0	72.9	175
2BC2	82	5.42	2.10	0.47	54900	3110	43100	4540	62.2	60.3	95.4
Norilsk											
O	5-6	6.40	42.90	22.90		23600	7340		2180	2820	3580
B1	13	7.10	7.96	1.61	43100	12800	24200	8390	64.5	45.0	274
B2	25	7.60	9.36	2.32	44200	13100	27400	8290	50.3	37.0	173
BC	49	8.60	5.35	0.81	62900	25600	29200	9410	50.6	31.0	122
BCK2	70	8.60	5.96	1.07	73900	32900	30900	11100	53.5	32.5	<100
Lake Lama											
O	4	4.00	65.20	32.30		3900	2980		81.9	72.4	1080
E	3	4.30	25.10	12.80		4140		3560	30.9	18.4	
B	7	5.10	20.30	5.31	31600	7980	22800	8650	47.9	41.6	198
BC	19	5.40	13.70	2.91	37500	9750	23000	10100	58.1	42.3	198
BC2	41	6.00	7.68	1.10	43800	13400	25800	10900	66.5	39.7	124
C	70	6.20	6.00	0.70	50700	15800	29000	10800	56.1	36.6	157

LOI: loss in weight on ignition; AR: aqua regia extractable concentration.

and heavy metals include species-specific effects, leading to shifts in fungal species composition. Third, pollutants often reduce the amount of litter produced by vegetation, and also change its composition. In addition, the direct effects of acidifying pollutants on soil invertebrates, resulting for example in the disappearance of earthworms in polluted areas (section 5.3.2), can affect the mechanical degradation of litter. Both the amount and composition of the litter, as well as changes in the degradation processes of soil organic matter, may have several impacts on soil decomposer activity and diversity.

In the Arctic, relatively few studies have been carried out on the impacts of acidifying pollutants on soil micro-organisms. The adverse effects on soil microbial biomass and activity seem to be concentrated in relatively restricted areas around the nickel-copper smelters where there are high levels of heavy metal deposition. For instance, Ohtonen and Väre (1996) reported no detectable changes in soil microbial activity and biomass parameters at distances of no more than 14 km from the smelter complex at Nikel. On the other hand, soil microbial activity and biomass were reported to be extremely reduced near Monchegorsk. For example, basal respiration of micro-organisms at sites 8 and 10 km from the Monchegorsk smelter was 5 and 8 µg carbon dioxide (CO<sub>2</sub>) per gramme of organic matter (OM) per hour, respectively, while at sites 36 km or more the basal respiration was 16 to 40 µg CO<sub>2</sub>/g OM/hr. Microbial biomass was 2.2 and 2.4 C/g OM at nearby sites, and 2.4-5.6 C/g OM at more distant sites (Ohtonen and Väre, 1996). Also, the diversity of microfungi (Lebedeva, 1993; Evdokimova, 2000), algae, and prokaryotes (non-spore forming gram negative bacteria, cyanobacteria and streptomycetes) (Evdokimova, 2000) in forest soils decreased moving towards the pollution source. In another study (Nikonov *et al.*, 2001), populations of prokaryotes in the soils of Norway spruce stands were found to increase, and populations of eukaryotes to decrease, moving towards Monchegorsk. Deterioration of the forest soil, leading to losses in soil organic matter and nutrients, was assumed to be connected to the adverse effects of heavy metals on soil fungal populations in drier Scots pine forests (Polyanskaya *et al.*, 2001). Changes in the composition of soil microbial

communities have also been found in taiga forests around the Kostamuksha iron pellet plant (Zaguralskaya and Ziabchenko, 1994) and in subtundra forests of the Taymir Peninsula affected by the Norilsk nickel-copper smelter (Kirtsideli *et al.*, 1995). Intriguingly, rhizosphere of severely damaged larches near Norilsk contained 10 to 100 times more micro-organisms than healthy larches in unpolluted regions; this phenomenon presumably resulted from the development of saprotrophic microbiota that benefit from root decline (Raguotis, 1989).

Fluoride deposition originating from the Kandalaksha aluminum smelter caused slight alkalization of the soil and altered the community composition of soil fungi, although no effects on the biomass of soil bacteria and fungi were found (Evdokimova, 2001). However, a more detailed investigation revealed a significant (by a factor of eight) decline in fungal biomass near Kandalaksha (Evdokimova *et al.*, 2004).

Acidification may not necessarily affect the basidiomycete and ascomycete ectomycorrhizal fungal symbionts of forest trees, as they are well adapted to naturally acidic conditions in forest soil (Smith and Read, 1997). However, a decrease in ectomycorrhizal colonization in the roots of forest trees has been found in response to soil acidification (Danielson and Visser, 1989; Holopainen *et al.*, 1996; Brunner, 2001). Emissions from the Monchegorsk smelter have been reported to negatively affect the diversity of ectomycorrhizal fungi (Isaeva, 2004).

Studies on the effects of acidifying pollution on arbuscular mycorrhizal (AM) colonization in herbs and grasses in the Arctic are even fewer than those dealing with ectomycorrhizal symbionts of trees. There appears to have been only one survey (Ruotsalainen *et al.*, 2006) on root colonisation in *Deschampsia flexuosa*, a common perennial grass that is tolerant of both acidification and heavy metals. There was a decrease in hyphal colonization of both AM and DSE (dark septate) fungi with increasing pollution levels. In contrast, DSE microsclerotia were more abundant in highly polluted sites, possibly indicating the strategy of these fungi to survive and disperse from unfavorable environments. Severe damage to field layer vegetation creates a patchy environment, such as the industrial barrens around

Monchegorsk, and under these conditions AM fungi may have difficulties dispersing from one plant host to another. This may have further impacts on fungal communities in the soil.

The cover of *Cladina* lichens, which is assumed to be especially effective at filtering out heavy metals, has deteriorated and even disappeared in industrial barrens and declining forests (Tømmervik *et al.*, 1995), and this may accelerate the adverse impacts of acidifying pollutants on the activity of soil micro-organisms. The structure of the organic layer has been damaged severely in these areas (Rigina and Kozlov, 1999). This is partly due to the loss of living fungal hyphae which, together with the fine roots, normally play an important role in maintaining the structure of this layer.

Thus, changes in ground and field layer vegetation due to acidifying pollution in the Arctic are linked to soil deterioration and changes in soil microbiology. On the basis of sparse information on soil micro-organisms in the Arctic, the adverse effects of acidifying pollutants seem to be concentrated in restricted areas around the nickel-copper smelters and to be more strongly linked to the excessive deposition of heavy metals rather than to acidification. The diversity of soil micro-organisms may be affected to a greater extent or at lower levels of pollution, than the biomass and activity of the soil microbial communities.

## 5.2. Effects on vegetation in the European Arctic

The damage zones around Nikel and Monchegorsk on the Kola Peninsula reported in the previous AMAP assessment (AMAP, 1998) ranged from forest death to a zone with changes in the microscopic structure of epiphytic lichens only. The corresponding range in annual average atmospheric SO<sub>2</sub> concentrations was from >40 to 2–4 µg/m<sup>3</sup> (Tikkanen and Niemelä, 1995; Tømmervik *et al.*, 1995). Nikel, although having a much smaller smelter than Monchegorsk, has had (for technical reasons) considerably higher SO<sub>2</sub> emissions (but lower metal emissions) than Monchegorsk. It is also further north and so, having a harsher climate, the vegetation is dominated by subarctic birch forest rather than the boreal forests around Monchegorsk (Kashulina *et al.*, 1997). Satellite data showed that the total area affected by air pollution (average SO<sub>2</sub> concentrations >10 µg/m<sup>3</sup>) around Nikel increased from 400 km<sup>2</sup> in 1973 to more than 3900 km<sup>2</sup> in 1988, and remained at this level during the early 1990s (Høgda *et al.*, 1995; Tømmervik *et al.*, 1995; see also Aamlid *et al.*, 1995; Gytarsky *et al.*, 1997). Furthermore, episodes of high SO<sub>2</sub> emissions had caused changes in the lichen and dwarf shrub communities over an additional 1100 km<sup>2</sup> in 1988, the area affected by severe air pollution near Nikel thus increasing to more than 5000 km<sup>2</sup>.

The size of the forest-death area was over 400 km<sup>2</sup> in the Nikel-Zapolyarnyy and Varanger regions; the area near the border between Russia and Norway studied in 1988 (Tømmervik *et al.*, 1995). At Monchegorsk, the forest-death area covered between 400 and 500 km<sup>2</sup> and extended more than 10 km to the south and more than 15 km to the north of the smelter complex (Mikkola, 1996). The average modeled SO<sub>2</sub> concentrations exceeded 40 µg/m<sup>3</sup> in this area (Tuovinen *et al.*, 1993). Marked defoliation of conifers and an absence of epiphytic lichens occurred in the area with

annual average modeled SO<sub>2</sub> concentrations of 15 to 40 µg/m<sup>3</sup>. External damage on the needles of Scots pine (e.g., tip necrosis) and changes in species composition of the lichen communities occurred in areas with annual average modeled SO<sub>2</sub> concentrations of 8 to 15 µg/m<sup>3</sup>, and the damage area extended to the eastern part of Inari in Finland. The total area covered by visible damage zones surrounding Monchegorsk, Nikel, and Zapolyarnyy was estimated to be around 39 000 km<sup>2</sup>. Around Monchegorsk, chlorosis of pine needle stomata occurred in areas with annual average modeled SO<sub>2</sub> concentrations of 4 to 8 µg/m<sup>3</sup>, and there were changes in the microscopic structure of pine needles and epiphytic lichens where the annual average modeled SO<sub>2</sub> concentrations were 2 to 4 µg/m<sup>3</sup> (Tikkanen and Niemelä, 1995).

In the mid-1990s, the proportion of damaged ecosystems in the European Arctic increased steadily from the northern boreal forest zone to the tundra. Furthermore, the same stages of ecosystem disturbance (severe, damaged, depressed, no damage) coincided with lower concentrations of the major pollutants in humus in tundra and in subarctic birch forest ecosystems than in forested areas (Kashulina *et al.*, 1997). On the other hand, a vegetation study from South-Varanger in eastern Finnmark, Norway (Aarrestad and Aamlid, 1999) showed that the 'pollution variables' Cu and Ni in humus, Ni in *Cladina*, and the modeled atmospheric SO<sub>2</sub> concentration, together only explained around 9% of the species variation after taking into account variation due to natural environmental variables. The comparisons between sulfur concentrations in moss and Ni concentrations in humus at various levels of ecosystem damage demonstrate that the state of the ecosystems in the mid-1990s in Norway and Finland had, with few exceptions, no correlation with the deposition of pollutants from the Russian smelters. In Russia, however, pollution strongly influenced ecosystem degradation. There, even the 'no damage' class had a higher median value for Ni and sulfur than for all damage classes in Finland or Norway (Kashulina *et al.*, 1997).

As there have been very few studies on acidification effects on vegetation in the Arctic since the last AMAP assessment, this section reviews the few new studies together with the results of studies based on data collected in the early 1990s, but mainly published in the late 1990s or early 2000s. The studies include data on direct and indirect effects of air pollutants, especially SO<sub>2</sub>, and both critical levels and critical loads are addressed.

### 5.2.1. Lichen-dominated and mountain birch (tundra) ecosystems

The main vegetation types in the northernmost and high altitude areas of northern Norway and the Kola Peninsula consist of lichen and dwarf shrub communities. Changes in the area with reindeer lichen (*Cladina* spp.)-dominated vegetation (heaths and forests with dwarf birch and mountain birch, *Betula nana* and *B. pubescens* ssp. *czerepanovii*) around the Cu-Ni smelters in Nikel and Zapolyarnyy correlated with changes in emissions of SO<sub>2</sub> between the 1970s and late 1990s (Høgda *et al.*, 1995; Tømmervik *et al.*, 1995, 1998, 2003). Thus, there was a reduction in the areas of lichen-dominated mountain heaths and forests from 37% in 1973 to 10% in 1992, followed by a slight increase to 12% in 1999 in the border areas of Norway and Russia (Tømmervik *et al.*, 2003). Reindeer lichens had completely disap-

peared from around two Ni-Cu smelter complexes on the Kola Peninsula by the 1990s (Kalabin, 1991; Tømmervik *et al.*, 1995), their growth rates reaching normal levels at a distance of 50 to 60 km from the smelters (Helle and Kojola, 1992). In addition to *Cladonia stellaris*, the arctic lichen *Cladonia rangiferina* is also sensitive to SO<sub>2</sub> and/or metals (Koptsik *et al.*, 2003). Kapitsa and Golubeva (1997) ranked the lichens *Cladonia mitis* and *Cladonia cornuta* among the most sensitive species to anthropogenic pressure, while the lichen *Cetraria nivalis* was ranked as less sensitive, and the low bushes *Phyllodoce caerulea* and *Salix glauca* were among the most tolerant species. Gaseous uptake measurements on lichens are limited, but according to Winner *et al.* (1988, cited in Nash and Gries, 1995), SO<sub>2</sub> uptake by *C. rangiferina* is at least an order of magnitude greater than in a typical vascular plant. The lichen-dominated vegetation types were, however, not just changed into barrens or partly damaged vegetation entities, but were also changed into heath (and woodland) vegetation with sparse lichen cover and were dominated by dwarf shrubs such as bilberry (*Vaccinium myrtillus*) in the Nikel–Zapolyarnyy and South-Varanger areas in the early 1990s (Tømmervik *et al.*, 1995, 1998).

The gradual change of lichen-dominated vegetation between 1973 and 1999 into barrens or sparsely vegetated areas or into other vegetation formations: partly damaged heather-dominated vegetation, bilberry forests/heaths, and meadow forests (on better soils in lowland areas) occurred mainly between 5 and 30–40 km from the smelters at Nikel and Zapolyarnyy, depending on prevailing wind directions during the growing season. For example, the previous sparsely vegetated coastal mountains have become more barren due to large emissions of SO<sub>2</sub> and their location within the path of the prevailing winds. On the other hand, the slight increase from around 10 to 12% in the lichen-dominated communities (*Betula nana*-lichen heath and *Empetrum*-lichen types) during the 1990s indicates an improvement in air quality and environmental conditions for these vegetation formations (Tømmervik *et al.*, 2003); for example, a decrease from 20 to 10 µg/m<sup>3</sup> in the summer SO<sub>2</sub> concentrations at Svanvik in the Norwegian–Russian border area (SFT, 2002). The significant positive relationship between lichen (*Cladonia* sp.)-dominated forests (*Empetrum*-lichen type with birches;  $r=0.84$ ,  $p=0.018$ ) and the significant negative relationship between changes in the area of industrial barrens ( $r=-0.95$ ,  $p=0.001$ ), and the changes in SO<sub>2</sub> emissions between 1973 and 1999 may be explained by the response times of the different land cover classes to SO<sub>2</sub> emissions. There is always a lag in the vegetation response to changes in SO<sub>2</sub> emissions and in this case the area of lichen-dominated vegetation types decreased and the barrens increased at the same time as

emissions decreased (but not reaching 1973 levels before the mid-1990s) (Tømmervik *et al.*, 2003).

Average SO<sub>2</sub> concentrations for the lichen-dominated vegetation (*Betula nana*-lichen heath and *Empetrum*-lichen types) were 3.5 and 9.2 µg/m<sup>3</sup>, respectively, and were 13.6 µg/m<sup>3</sup> for heather woodland/partly damaged heather (*Empetrum-Vaccinium* type) with a sparse/reduced lichen cover (Tømmervik *et al.* 1995, 1998). There were associations between the damaged land cover, the industrial barren land, and the SO<sub>2</sub> concentration in air at ground level and Ni and sulfur concentrations in reindeer lichens. Average sulfur concentrations in reindeer lichens (*C. stellaris* and *C. mitis*) from *Empetrum*-lichen type forests and *Empetrum-Vaccinium* type heather woodland/partly damaged heather were 489 and 765 mg/kg, respectively, in the border areas between Norway (Southern-Varanger) and Russia (Nikel–Zapolyarnyy) in the late 1980s to early 1990s. In the bilberry forests/heaths (*V. myrtillus*-*D. flexuosa* and *Cornus suecica* types), the average SO<sub>2</sub> concentration was 14.9 µg/m<sup>3</sup> and the lichen sulfur concentration was 644 mg/kg (Tømmervik *et al.*, 1998). The vegetation damage observed and the slightly elevated sulfur concentrations in lichen thalli may be attributed to both high short-term SO<sub>2</sub> concentrations as well as long-term exposure to elevated sulfur deposition based on the proposed critical level of 10 µg SO<sub>2</sub> / m<sup>3</sup> (annual mean) to be adopted for cyanobacterial lichens (UNECE, 1993, 2004a).

One of the nine catchments where Reimann *et al.* (2001b,c,d, 2003) studied total sulfur concentrations in the leaves of various species was an arctic tundra catchment in Berlevåg on the Barents Sea coast of northern Norway. They did not report air quality and/or deposition data and so could not calculate correlations between air quality and/or deposition and plant and soil sulfur concentrations. At any rate, their results did not show any effects of changes in SO<sub>2</sub> emissions from the Kola Peninsula, and/or SO<sub>4</sub> in marine aerosols, on the sulfur concentrations of plant samples collected in summer 1999. For example, the total sulfur concentrations of *Hylocomium splendens* and *Pleurozium schreberi* ranged from 716 to 863 mg/kg and for *V. myrtillus* from 1720 to 2230 mg/kg. They concluded that total sulfur in moss is largely governed by the input of particulate material (dust), such as metal sulfides (Gregurek *et al.*, 1999), and not by atmospheric SO<sub>2</sub> concentrations. The total sulfur concentrations of plant and soil samples in Berlevåg were similar to those in the Russian catchments of Vorkuta and Naryan-Mar, with all three areas showing large variations in the total sulfur concentrations in leaves both between species within each area as well as between areas (Figure 5.10, Table 5.2).

Particulate emissions that contain heavy metals are deposited closer to the emission sources than SO<sub>2</sub> and

Table 5.2. Median sulfur concentrations (mg/kg dry weight) in plant leaves collected from four northern European catchments in summer 1999 (Reimann *et al.*, 2001c, 2003).

Catchment	Berlevåg (arctic tundra)	Monchegorsk (northern taiga)	Naryan-Mar (southern tundra with birch and shrub)	Vorkuta (moss-lichen shrub tundra with birch and willow)
Moss ( <i>Hylocomium splendens</i> )	754	1090	889	824
Blueberry ( <i>Vaccinium myrtillus</i> )	2025	1710		
Cowberry ( <i>Vaccinium vitis-idaea</i> )	1350	2095	1300	1480
Crowberry ( <i>Empetrum nigrum</i> )	1220	1260	1130	1360
Birch ( <i>Betula pubescens</i> )		1775	1890	
Willow ( <i>Salix</i> spp.)	3765	2335	2350	3180
Pine ( <i>Pinus sylvestris</i> )		1190		
Spruce ( <i>Picea abies</i> )		1985	547	





Figure 5.10. Location of the catchments included in the Barents Project (Reimann *et al.*, 2003; reprinted from Reimann *et al.*, 2001d).

SO<sub>4</sub> (e.g., Gytarsky *et al.*, 1995) and therefore the relative importance of the direct and indirect harmful effects of sulfur compounds increases with increasing distance from the emissions sources. The occurrence of visible SO<sub>2</sub> injuries in the leaves (*B. pubescens*, *B. nana*, *V. myrtillus*, *V. uliginosum*) and Scots pine needles (*Pinus sylvestris*) in northeastern Norway close to the Russian border (Aamlid, 1993; Aarrestad and Aamlid, 1999) (Figure 5.11) has mainly been attributed to episodes with high SO<sub>2</sub> concentrations that occur under specific meteorological conditions (Jerre, 1994). Hilmo and Larsen (1994), in turn, reported morphological differences (growth form, fertility, apothecial morphology, discoloration) in the epiphytic lichens *Hypogymnia physodes* and *Melaniella olivacea* at sites exposed to high concentrations of SO<sub>2</sub> in Sør-Varanger compared to lichens at an unpolluted control site. As *M. olivacea* had a larger mean cover on birch stems than *H. physodes* in the border area between Norway and Russia, and was more frequently found at the plots in the 1991 and 1993 studies (Aamlid and Skogheim, 2001), it may even be considered more tolerant to the direct and indirect effects of sulfur

deposition than *H. physodes*. The zero zone of lichen cover coincided approximately with the areas where the modeled annual mean SO<sub>2</sub> concentrations reached 25 µg/m<sup>3</sup>. However, sulfur compounds are not the only acidifying compounds; Hilmo and Larsen (1994) found morphological changes (discoloration, growth form) in *Parmelia sulcata* at an industrial site in Glomfjord that may be related to elevated nitrogen deposition.

The results of Tømmervik *et al.* (1998, 2003) on vegetation changes in the area of Pasvik and Zapolyarnyy/Nikel at the Norwegian–Russian border show an increase from about 7% in 1973 to about 20% in 1979, with a stabilization at approximately 30% by 1999 in bilberry and low herb vegetation formation and meadow vegetation (bilberry forests/heaths and meadow forests with mountain birch). This change coincided with the decrease in the areas of lichen-dominated vegetation and was mainly due to the transformation of dwarf shrub (*Empetrum hermaphroditum*) and lichen-dominated vegetation cover types into bilberry and *Deschampsia flexuosa*-dominated formations which appear up to 5 to 40 km from the smelters (Deyeva and




Figure 5.11. Visible (acute) SO<sub>2</sub> injuries on (a) birch (*Betula pubescens*) leaves and (b) Scots pine (*Pinus sylvestris*) needles at Svanvik, Norway, approximately 10 km northwest of Nikel (Photos: Dan Aamlid).





Figure 5.12. Land cover maps for the Pasvik–Nikel area for (a) 1973, (b) 1994, and (c) 1999 (Tømmervik *et al.*, 2003).



Figure 5.13. Location of joint Norwegian/Russian sites for intensive terrestrial monitoring (SFT, 2002).

Maznaja, 1993; Tømmervik *et al.*, 1995, 1998, 2003). With regard to pollution resistance of mountain birch, studies in the vicinity of the smelters at Monchegorsk may suggest phenotypic acclimatization of mature trees to a gradual increase in pollution as well as selection for pollution-resistant genotypes that may have occurred due to the exceptionally high emissions pressure over the past few decades (Valkama and Kozlov, 2001; Kozlov and Zvereva, 2004; Kozlov, 2005).

### 5.2.2. Coniferous forest ecosystems

According to Tømmervik *et al.* (2003), the vegetation cover that had the highest association ( $r=-0.94$ ,  $p=0.001$ ) with changes in  $\text{SO}_2$  emissions in the Pasvik and Nikel–Zapolyarnyy areas was the mixed birch-pine forests with lichen content (Figure 5.12). The increase in the area of mixed pine-birch forests, especially from 1988 to 1999, meant that the area affected by air pollution was reduced. The most important differences in species composition in the Nikel–Pasvik area along the Norwegian–Russian border (see Figure 5.13 for the location of the Russian–Norwegian vegetation monitoring sites) were related to the loss of species richness, i.e., a lower abundance of bryophytes (e.g., *H. splendens*, *P. schreberi*, *Dicranum* species) and lichens (several *Cladonia* species) (Aamlid *et al.*, 2000), and a concomitant increase in the dominance of air pollution resistant species such as *D. flexuosa* and *E. hermaphroditum* (SFT, 2002; Tømmervik *et al.*, 2003; see also Vassilieva *et al.*, 1995) (Table 5.3). These differences are explained on the basis of both the direct impacts of pollution and the indirect soil-mediated processes leading to, for example, soil




Table 5.3. Average number of species within 1 m<sup>2</sup> plots in pine forest along a western gradient from Nikel (SFT, 2002). The highest values for each species group (bold font) indicate the change from lichen- and moss-dominated vegetation to small shrub- and grass-dominated vegetation near the emissions sources.

Species group	Sample plots and their distance from Nikel						
	45 km south	25 km west	16 km west	12 km west	9 km northwest	6 km north	6 km west
Small shrubs	3.5	2.9	3.1	3.1	3.4	<b>4.2</b>	4.0
Herbs	1.9	1.9	1.7	0.8	1.9	<b>2.0</b>	1.1
Mosses	<b>5.1</b>	4.6	4.7	4.7	4.6	1.2	2.1
Liverworts	1.3	1.5	<b>2.0</b>	1.5	1.6	0.7	1.2
Lichens	3.7	4.1	7.5	<b>8.2</b>	3.9	2.3	3.1
All species <sup>a</sup>	15.4 (24)	14.9 (30)	<b>19.2 (32)</b>	18.7 (31)	15.6 (26)	10.9 (23)	11.5 (17)

<sup>a</sup> maximum number of species in brackets

acidification and nutrient imbalances (Tømmervik *et al.*, 1995; Aarrestad and Aamlid, 1999; Koptsik *et al.*, 1999a,b, 2003; SFT, 2002). Furthermore, for example, *Pohlia nutans* is a moss species that has increased its cover in the most polluted area. This species, characterized by wide ecological amplitude, colonized the emptying litter in industrial barren land in the absence of competition from other (moss) species more sensitive to pollution (Koptsik *et al.*, 2003).

According to Kashulina *et al.* (1997, 2003), direct exposure to SO<sub>2</sub> is the most likely, although not the only mechanism of vegetation damage in the eight catchments located on the Kola Peninsula and in northeastern Finland and Norway. The highest sulfur concentrations in plant samples in the most polluted sites were usually 2- to 4-fold higher than at control sites (Gytarsky *et al.*, 1995; Äyräs *et al.*, 1997; Aamlid *et al.*, 2000; Steinnes *et al.*, 2000; Reimann *et al.*, 2001a,c). Steinnes *et al.* (2000) reported a decrease from 4270 to 2000 mg/kg and from 2820 to 1420 mg/kg in the mean total sulfur concentration of *B. pubescens* and *E. hermaphroditum* leaves, respectively, on moving from the most polluted study sites around Nikel to the southernmost sites more close to background conditions in 1991. A decrease from around 3500 to 1000 mg/kg, and from around 3400 to 1500 mg/kg in the foliar sulfur concentration of *Betula* sp. and *V. myrtillus*, respectively, between the areas closest to the Nikel smelters (<5 km) and background areas (about 30 km) was, in turn, reported by Gytarsky *et al.* (1995). Sulfur was also the major element in moss (*H. splendens* and *P. schreberi* 2090–543 mg/kg) and the O and B soil horizons in the 1995 survey in northern Finland, Norway, and Russia (Äyräs *et al.*, 1997; Kashulina *et al.*, 1997, 2003; Kashulina and Reimann, 2001, 2002; Reimann *et al.*, 2001a).

Reimann *et al.* (2001b,c,d, 2003) measured total sulfur concentrations in the leaves of several plant species collected from nine catchments in northern Europe (Finland, Norway and Russia) in summer 1999. Only Norway spruce (*Picea abies*) needles showed significantly higher sulfur concentrations in all samples collected near Monchegorsk. Correlation between 'available' sulfur in surface soil and observed foliar sulfur concentrations for all plants growing in the soils was in general very low. Furthermore, Reimann *et al.* (2003) concluded that macroscopic and microscopic evidence of leaf damage are more reliable indicators of pollution impact than foliar sulfur concentration. In a study carried out in the early 1990s on Scots pine, Rautio and Huttunen (2003) also found a weak relationship between element concentrations in the soil and internal foliar concentrations. A factor representing high foliar levels of Ni, Cu, and sulfur, and low levels of Zn and Mn, was found to explain most of the variation in the number of needle age classes and tip necrosis. The macroscopic injury variables (including stomatal chlorosis and other discolorations) correlated clearly with the modeled SO<sub>2</sub> concentration in air.

However, the cell injuries studied did not show any distinct relationships with element concentrations in either foliage or deposited particles (Rautio *et al.*, 1998c).

High variation in the total sulfur concentrations in plant samples collected during the growing season may be explained (in addition to species-specific characteristics and differences in gaseous and particulate sulfur deposition) by nutrient status, pH, and other soil characteristics, as well as by climate, composition of the vegetation layer and altitude (Reimann *et al.*, 2001b, 2003). For example, despite higher SO<sub>2</sub> emissions from the Nikel smelter compared to the Monchegorsk smelter, plants accumulate more sulfur near the latter because of the specific combination of geomorphological and meteorological conditions (Koptsik *et al.*, 1999a). It is also well known that plant nutrient concentrations (including sulfur) vary considerably during and between growing seasons. For example, the average sulfur concentrations of current-year Scots pine needles collected in August–October between 1996 and 1999 were 755 to 901 mg/kg at Värriö in the eastern part of Finnish Lapland and 806 to 988 mg/kg at Sammaltunturi (i.e., Pallas) in the western part of Finnish Lapland (Sirkkä Manninen, University of Helsinki, unpubl. data). This is why needle samples for bioindicator studies mapping the dispersal of sulfur emissions, as well as the area possibly affected by sulfur deposition, are usually collected in winter, i.e., when emissions are at their highest and the trees are physiologically relatively inactive (see e.g., Manninen *et al.*, 1997a, 1998; Rautio *et al.*, 1998a,b).

The latitude-related decrease in the total sulfur concentration on moving northward (Reimann *et al.*, 2003) may be related to the length of the periods during which the plants have stomatal uptake of SO<sub>2</sub> versus the time when they are exposed to non-stomatal deposition only, as the proportion of (winter) non-stomatal deposition increases towards the north. The dry deposition of sulfur dominates in northern areas due to the low precipitation, and dry deposition accounts for about 80% of the total deposition (Tuovinen *et al.*, 1993). On the other hand, as most of the plants have a snow-cover for six to seven months of the year, more than 50% of the total annual deposition will deposit onto snow (Kashulina and Reimann, 2002). This means that the sulfur concentrations of mosses and lichens which are thallophytes (i.e., they have no roots but obtain all their nutrients through their above-ground parts as ions dissolved in water – rain and melting snow), may give a different picture about the rate and distribution of sulfur emissions compared to that given by foliar sulfur analyses of evergreen conifers. Steinnes (1995) pointed out that it may, however, be feasible to achieve representative results (at least for heavy metals) by carefully selecting the sampling site for moss in order to avoid interference from wind erosion or surface water flow. On the other hand,

the lower growth rate of mosses in arctic regions may be expected to lead to greater uptake of airborne trace elements per unit weight than in mosses at more southerly latitudes. Furthermore, the different growth structure of the plant may make it more difficult to accurately define the exposure period.

In addition to the total sulfur concentration of Scots pine needles, statistically significant correlations ( $p < 0.001$ ) were found between the annual mean  $\text{SO}_2$  concentration and needle sulfate concentration and S:N ratio at the sites close to Monchegorsk in the early 1990s (Manninen *et al.*, 1998) (Table 5.4). Koptsik *et al.* (1999a, 2001) also found a marked gradient in the needle total sulfur concentration and S:N ratio of Scots pine with increasing distance from the Nikel smelters in the early 1990s. The increase in the S:N ratios towards the Nikel smelters is not just explained by the increasing needle total sulfur concentrations, but by lower nitrogen supply from the soil observed as decreasing concentrations of nitrogen in soil organic horizons toward the smelter. In addition to nitrogen concentrations, those of ammonium acetate-extractable Mg and K in organic horizons tended to be lower within several kilometers of the Nikel smelters compared to remote sites (Koptsik *et al.*, 1999b, 2001, 2003) as they also were around the Monchegorsk smelters (Lukina and Nikonov, 1999, cited in Zvereva and Kozlov, 2005). With regard to temporal trends related to the decrease in  $\text{SO}_2$  emissions and, consequently, in the ambient  $\text{SO}_2$  concentrations in the border areas of Norway and Russia (Hagen *et al.*, 2005), the results of Koptsik *et al.* (G. Koptsik, Moscow State University, pers. comm.) suggested a decrease in the total sulfur concentration of current-year Scots pine needles from August 1991–1994 to August 2002. At any rate, as there are differences in foliar responses to elevated sulfur and nitrogen deposition between coniferous tree species (Manninen and Huttunen, 2000; Luyssaert *et al.*, 2005), more research is needed on foliar sulfur and nitrogen fractions and ratios, as well as on Mg:N and Ca:Al ratios, in the northernmost areas for use as indicators of acidification.

### 5.2.3. Reindeer grazing, climate change, nitrogen deposition, and other factors

Reindeer grazing is one factor that modifies or interacts with the effects of sulfur deposition on vegetation, especially in northern Norway. According to Gaare and Tømmervik (2000), reindeer have grazed down the new 'reindeer' lichens in the period 1992 to 1999, i.e., after the emissions

reductions at the smelters on the Kola Peninsula. There has also been extensive forestry in the southern part of the Pasvik area and this has led to reductions in lichen cover (succession) and pine forests. Severe desiccation of the forest vegetation close to the smelters at Nikel and Zapolyarnyy has led to increasing amounts of dead needles and woody debris, which in turn has led to more frequent wild forest fires (Tømmervik *et al.*, 2003). Moreover, outbreaks by caterpillars such as *Epirrita autumnata* have been followed by successions from lichen-dominated vegetation to more dwarf shrub- and grass-dominated vegetation (Lehtonen, 1987; Tenow and Bylund, 2000).

Over most of northernmost Europe, the trend towards a shift in lichen-dwarf shrub dominated forests to more grass- and herb-dominated forests, may reflect a situation where slow-growing species such as lichens, crowberry (*E. hermaphroditum*), and cowberry (*V. vitis-idaea*) are declining due to intensive reindeer grazing (Olofsson *et al.*, 2001). Tømmervik *et al.* (2004) also gave climate change (increased precipitation) and long-range transported nitrogen pollution (increasing nitrogen deposition is also linked to increasing precipitation) as possible accentuators for the increase of birch forests and vegetation types dominated by bilberry, wavy hair-grass (*D. flexuosa*), the dwarf cornel (*C. suecica*), and mosses (*P. schreberi*, *H. splendens*) in northern Norway over the last 40 years.

Increased nitrogen deposition through precipitation and/or deposition of manure (feces) from the reindeer may create unfavorable living conditions for slow-growing species. Bobbink *et al.* (2002) considered an empirical critical load of 5 to 10 kg N/ha/yr, based on field manipulations in which exceedance was indicated by changes in biomass and in the moss layer, reliable for tundra ecosystems. They gave the same critical load for arctic, alpine, and sub-alpine shrub habitats (with exceedance indicated by a decline in lichens, mosses, and evergreen dwarf-shrub), but its reliability was referred to as 'expert judgment' (see also UNECE, 2004a). Recent results from field experiments in an area of northern Sweden with low background nitrogen deposition and from a large-scale monitoring study, show that important vegetational changes start to occur when adding low nitrogen doses and that recovery of the vegetation after nitrogen inputs stop is a very slow process. Based on data from the Swedish research program Abatement Strategies for Transboundary Air Pollution, Nordin *et al.* (2005) suggested that the critical load should be lowered from the current 10 to 15 kg N/ha/yr to a level of 6 kg N/ha/yr to protect the biodiversity of the boreal forest understorey vegetation. The impact of elevated nitrogen

Table 5.4. Concentrations (mg/kg) and ratios of sulfur and nitrogen in current-year Scots pine needles collected at different distances from the smelters on the Kola Peninsula showing change along the pollution gradients and seasonal variation in needle chemistry.

	Sampling	Total S <sup>a</sup>	Organic S	SO <sub>4</sub> -S	SO <sub>4</sub> -S/organic S <sup>a</sup>	Total N	Total S/Total N <sup>a</sup>	Reference
Finnish Lapland <sup>b</sup>	Sep-Oct 1990, Sep 1992	720-1153	479-862	141-361	0.20-0.53	9200-13000	0.032-0.045	Manninen <i>et al.</i> , 1997b
Monchegorsk- Jena <sup>c</sup>	Apr 1991 Jul 1992	856-2548 794-1474	672-1599 619-971	105-1297 168-656	0.14-1.04 0.21-0.80	8100-13800 8400-14200	0.032-0.089 0.034-0.055	Manninen <i>et al.</i> , 1998
Norwegian-Russian border <sup>b</sup>	Aug 1991-1994 Aug 2002	499-1917 896-1248				13400-21300	0.031-0.046	Koptsik <i>et al.</i> , 1999a; 2001; pers. comm.
Monchegorsk- Finnish border <sup>d</sup>	Sep 1990 Apr 1991	1112±471 1321±499						Rautio <i>et al.</i> , 1998b

<sup>a</sup> The background total sulfur concentration is around 600 mg/kg,  $\text{SO}_4$ -S concentration 100 to 200 mg/kg, and total sulfur:total nitrogen ratio 0.028 (on a gramme-atom basis) (Manninen, 1995 and references therein); <sup>b</sup> min-max site means, <sup>c</sup> min-max values for individual tree, <sup>d</sup> mean±SD for all sites.



deposition was seen as decreased abundance of *V. myrtillus* and *V. vitis-idaea* and increased abundance of *D. flexuosa* in particular, as well as decreased abundance of *P. schreberi* and *H. splendens*.

Care must be taken when drawing conclusions about nitrogen deposition on the basis of plant nitrogen analyses. For example, Walker *et al.* (2003a) found marked regional gradients in terricolous lichen nitrogen concentrations in the Usa river basin, northeast European Russia, suggesting that there were concomitant gradients in nitrogen deposition. On the 240 km south-to-north transects through the town Inta, values of  $[N]_{\text{apex}}$  (the apical 5 mm part) in *Cladonia stellaris* decreased from  $0.57 \pm 0.01$  mmol/g to  $0.43 \pm 0.01$  mmol/g. Moreover, elevated  $[N]_{\text{apex}}$  in *C. arbuscula*, and to a lesser extent in *Flavocetraria cucullata*, in the Vorkuta area suggested elevated nitrogen deposition due to local industrial pollution. Variation in  $[N]_{\text{apex}}$  in *C. arbuscula* correlated well with other pollutant signals in snow ( $[SO_4^{2-}]_{\text{snow}}$ ,  $[Ca^{2+}]_{\text{snow}}$ ,  $[K^+]_{\text{snow}}$ , pH) (Walker *et al.*, 2003b) as well as thalli, but as on the other transects, it was not correlated with  $[NO_3^-]_{\text{snow}}$ . Molar concentrations of organic nitrogen were broadly similar to those of nitrate, except at sites close to Vorkuta where  $[\text{organic N}]_{\text{snow}}$  values were greater than  $[NO_3^-]_{\text{snow}}$  by a factor of 2 or 3, varying spatially in a manner similar to that of  $[SO_4^{2-}]_{\text{snow}}$ . Thus Walker *et al.* (2003a) concluded that a higher deposition rate of organic nitrogen at the sites close to Vorkuta might have contributed to elevated  $[N]_{\text{apex}}$  in this area. In addition, they did not exclude that impaired lichen growth at sites close to Vorkuta, perhaps due to phytotoxic air pollutants, might have resulted in higher tissue nitrogen concentrations. They emphasized that data for snow pack chemistry were for a single winter period, whereas  $[N]_{\text{apex}}$  values probably gave an indication of nitrogen deposition integrated over a period of at least one to two years. Furthermore, that these interpretations were made in the absence of commensurate data on  $[NH_4^+]_{\text{snow}}$ .

According to Walker *et al.* (2003a), mat-forming lichen cover was generally poor in the Vorkuta region due to heavy grazing and trampling by reindeer; for example, *C. stellaris* was largely absent at most sites. Virtanen *et al.* (2002), in turn, identified two impact zones around Vorkuta on the basis of remote sensing data: (1) a pollution zone (150–200 km<sup>2</sup>), where most of the lichen species are absent and changes in vegetation communities' species composition in all main plant groups are obvious (willows especially being more dominant than in unpolluted sites), and (2) a slight pollution/disturbance zone (600–900 km<sup>2</sup>), where vegetation changes are mainly similar but less so than the changes in the first zone – in particular the amount of herbs and grasses is increased relative to unpolluted areas. The upper soil layers (15–30 cm) in Vorkuta, where atmospheric pollution is mainly caused by dust from open coalmines, emissions from coal combusting power plants and a cement factory, and burning of waste rock near the coalmines, have become chemically modified with pH ranging from 6.7 to 8.9 indicating strong alkalization, and concentrations of exchangeable Ca and total nitrogen are 10 to 20 and 2 to 4 times higher, respectively, than at comparable non-polluted sites. Heavy metal levels in soils are also increased in the area (Getsen *et al.*, 1994, cited in Virtanen *et al.*, 2002). Emissions data for the 1990s suggested that SO<sub>2</sub> emissions averaged around 40 000 tonnes per year from the Vorkuta industrial complex in the Komi Republic (Virtanen *et al.*, 2002). The only information found on the effects of emissions on terrestrial vegetation in the

Norilsk area was that by Tutubalina and Rees (2001) who reported a loss of vegetation from the city of Norilsk between 1961 and 1995 and the growth of landscape damage over a larger area between 1972 and 1995. Several studies on the spatial and temporal dynamics of forest decline in this area were published in Russian.

#### 5.2.4. Needs and recommendations for future research and monitoring

In the multi-medium ecogeochemical mapping carried out in the European Arctic (Russia, Finland and Norway) in the mid-1990s, the only sample material in which the highest bulk sulfur deposition figures were directly reflected in a subarctic birch forest catchment close to Zapolyarnyy was vegetation (Äyräs *et al.*, 1997; Kashulina *et al.*, 1997, 2003; Kashulina and Reimann, 2001, 2002; Reimann *et al.*, 2001a). However, the almost 10-fold increase in the bulk sulfur deposition on moving from a coniferous forest catchment (0.17 g S/m<sup>2</sup>/yr) at Pallas in the western part of Finnish Lapland to the subarctic birch forest catchment near Zapolyarnyy (1.5 g S/m<sup>2</sup>/yr) resulted in only a 2-fold increase in the sulfur concentration of moss samples (from a median of 800 to 1500 mg/kg).

Gytarsky *et al.* (1997) calculated that the annual dry deposition of sulfur is about 2-fold higher in deciduous forests compared to coniferous forests on the Kola Peninsula. Taking into account the high proportion of dry-deposited sulfur, they expected that the values of total sulfur deposition over the slightly damaged area (based on the classification by Boltneva *et al.*, 1982; Karaban *et al.*, 1988), where the modeled growing season mean SO<sub>2</sub> concentration was 21 µg/m<sup>3</sup>, would increase to 0.6 to 1.0 g/m<sup>2</sup>/yr in coniferous forests and 1.2 g/m<sup>2</sup>/yr in deciduous forests (Gytarsky *et al.*, 1995, 1997). In general, dry deposition to coniferous forest is considered to be higher than that in deciduous forest. The dry deposition velocity of 1.1 cm/s for SO<sub>2</sub> on deciduous forests during the vegetation period (McMahon and Denison, 1979) versus that of 0.3 cm/s on coniferous forests (Garland, 1977) used by Gytarsky *et al.* (1997) in their calculations may explain the high sulfur deposition in deciduous forest. At any rate, the values given by Gytarsky *et al.* (1995, 1997) are 4-fold higher than the proposed SO<sub>2</sub> critical level of 5 µg/m<sup>3</sup> as a growing season mean (Manninen, 1995; Manninen and Huttunen, 2000; see also Tsvetkov, 1993), as well as the critical load of 0.32 g/m<sup>2</sup>/yr derived for the highly sensitive forest ecosystems of northern Europe (Downing *et al.*, 1993). The summertime (April–September) mean SO<sub>2</sub> concentrations have been around 5 to 10 µg/m<sup>3</sup> at Svanvik (about 10 km west of Nikel) and reached 150 µg/m<sup>3</sup> at Nikel in the late 1990s (Hagen *et al.*, 2001, cited in SFT 2002). Thus acute and chronic impacts of SO<sub>2</sub> and sulfur deposition still occur in the area. There is a need for measurements of dry deposition velocities on different vegetation types.

According to Alexeyev (1995), the least studied but most important stage of air pollution impacts on arctic and subarctic terrestrial ecosystems is the initial damage, when the negative irreversible processes of ecosystem destruction are still preventable. Simultaneous determination of several signals can provide an early warning of damage to terrestrial vegetation by low-level airborne pollutants. It is recommended that studies be carried out at permanent plots (in cooperation with air quality, deposition, soil and faunal monitoring) in order to assess the state and



functioning of the tundra, mountain birch and coniferous forest ecosystems. These studies should focus on the biomass (growth) and species composition and coverage of field and ground layer vegetation, the condition and frequency of epiphytic lichens, visible foliar injuries, foliar sulfur and nitrogen concentrations, and ratios of S:N, Mg:N and Ca:Al using standardized methods (including different sampling dates for the coniferous, deciduous and thallophytic key species). In particular, more attention should be paid to nitrogen-related issues and interactions. Lichens (e.g., *Lobaria*, *Peltigera* and *Stereocaulon* species and *Nephroma arcticum*) containing cyanobacteria are abundant in arctic ecosystems and may account for a major part of the nitrogen fixed in these systems. Their nitrogen fixation may be even more sensitive to acidity than photosynthesis and may thus be decreased by decreasing pH (Nash and Gries, 1995). Climate change and elevated nitrogen deposition are expected to affect negatively the slow-growing lichen vegetation in particular (e.g., Cornelissen *et al.*, 2001; Gordon *et al.*, 2001; Fremstad *et al.*, 2005) and higher CO<sub>2</sub> concentrations and UV-B radiation may reduce nutrient cycling and so may potentially reduce ecosystem productivity (Callaghan, 2005 and references therein).

On the other hand, if the most sensitive tundra vegetation is at least partly replaced by more tolerant forest vegetation over the next 100 years (Kaplan *et al.*, 2003) and SO<sub>2</sub> emissions do not increase, the state of the vegetation may (continue to) improve – also taking into account that plant compensatory responses as well as allocation to reproduction are species specific (e.g., Zvereva and Kozlov, 2001, 2005). Alekseev and Soroka (2002) reported that a higher growth rate occurred in Scots pine on the northwestern Kola Peninsula despite significant air pollution. Generally, the younger the trees, the greater the growth increase. The most probable reasons for the marked increase in radial increment growth of Scots pine in the region are climate warming (including an increase in growth season length) and higher levels of CO<sub>2</sub> as well as changes in forest fire frequencies (increased frequency and, consequently, a shift towards younger forests). The lack of significant differences in diameter growth between trees in different damage classes is explained by Alekseev and Soroka (2002) by the low number of samples on severely damaged trees, which are most susceptible to growth suppression by pollution.

### 5.3. Effects on fauna

#### 5.3.1. Effects on birds and mammals

Acidification and acidic deposition typically affect terrestrial vertebrates indirectly. There are only a few studies concerning the direct effects of acidification on mammals and birds, but indirect effects mediated through exposure to metals in food or water have been reported (Dudley and Stolton, 1996). On the Kola Peninsula, all the vertebrate species studied, for example, root vole (*Microtus oeconomus*), grey-sided vole (*Clethrionomys rufocanus*), common shrew (*Sorex araneus*), willow grouse (*Lagopus lagopus*) and common frog (*Rana temporaria*), had elevated heavy metal concentrations in liver, bone, muscle, and skin. The accumulation of heavy metals seems to depend on trophic level (AMAP, 2005).

#### 5.3.1.1. Mammals

Mitotic activity of spleen and cornea cells in root voles of different age and sex was higher 4 km from the Monchegorsk smelter than 28 km from the smelter. Mature root vole males had a lower hemoglobin content at 4 km than at 28 km. The mean frequency of chromosome aberrations in spleen, cornea, and marrow cells of root voles ranged from 34 to 50%. In spleen and cornea cells there were no differences in aberration frequency between 4 and 28 km, but in marrow cells the frequency of aberrations was significantly higher at 4 km than at 28 km (Kataev and Popova, 1993).

Males of the field vole (*Microtus agrestis*) collected 4 km south of the Monchegorsk smelter had lighter livers and heavier spleens than males collected 28 km away (Kataev and Popova, 1993). However, correlation analysis based on data from eight sites located 3 to 52 km south of Monchegorsk did not show any relationship between body mass of adult grey-sided voles and distance from the smelter (Kataev *et al.*, 1994). The proportion of pregnant females was higher in unpolluted sites, but the number of fetuses per female was the same along the pollution gradient (Kataev *et al.*, 1994). However, resorption of embryos was much higher at the most polluted site (36.8%) than at the control site (3.8%) (Kataev and Makarova, 1984).

The sex ratio of grey-sided voles near the Monchegorsk smelter was around 1:1, while males made up 65% of the population at an unpolluted forest site. The age structure of the populations was also different: no overwintered individuals were found in the polluted sites compared to 5.9 to 7.7% of the population overwintering in unpolluted forests (Kataev and Makarova, 1984). The proportion of adult voles in catches from the moderately polluted sites around the Monchegorsk smelter was higher than that at unpolluted sites (Kataev *et al.*, 1994), suggesting that juvenile mortality is higher at moderately polluted sites than unpolluted sites.

The density of grey-sided voles, the most abundant vole species, was lowest close to the Monchegorsk smelter and increased with distance to the farthest, less-polluted trapping sites. The bank vole, red vole (*Clethrionomys rutilus*) and Norway lemming (*Lemmus lemmus*) were absent from the most severely damaged area, and were also scarce at the moderately polluted area 28 km south of the smelter (Kataev *et al.*, 1994). Snap trapping in the surroundings of Monchegorsk revealed six species of small mammals; the common shrew and grey-sided vole were the most common in unpolluted forests. Only the grey-sided vole and root vole were captured in industrial barrens (Kozlov *et al.*, 2005a).

A long-term study by Kataev (1995) revealed a monotonous decline in the population of small mammals on approaching the Monchegorsk smelter: the biomass decreased from 87 kg/km<sup>2</sup> at the unpolluted forest site to 69 kg/km<sup>2</sup> at the moderately polluted site to 15 kg/km<sup>2</sup> at the most polluted site. However, the pattern in 2001 (Kozlov *et al.*, 2005a) differed from earlier data (Kataev, 1995): the density decline in the most polluted sites was much smaller (51.8% of the background value) and, most importantly, the highest densities were associated with slightly to moderately polluted forest. This result may be partially explained by the presumably higher plant quality in moderately polluted sites: grey-sided voles preferred bilberry shoots at distances of 20 to 40 km from Monchegorsk compared to shoots from sites both closer and further away (Suomela and Palokangas, 1993).

Although pollution may have suppressed the cyclic density fluctuations of the bank vole (Figure 5.14), this and many other results obtained near Monchegorsk should be viewed as tentative due to the non-replicated experimental design (one polluted vs. one or two unpolluted sites).




Figure 5.14. Effects of pollution on density of bank vole (*Clethrionomys glareolus*) in Lapland Biosphere reserve (the break in observations is due to the Second World War). Site 1 is located 30 km south of Monchegorsk; it was considered unpolluted until 1958–1960, when the first signs of forest damage by emissions were detected. Site 2 is located 30 km west of the smelter and represents an unpolluted control site (protected from emissions by the Monche-tundra mountain range) (after Kozlov and Barcan, 2000).

Hörnfeldt (2004) listed several hypotheses to explain the long-term decline in vole numbers and changes in vole dynamics. One hypothesis attributed the decline to the decrease in quantity or quality of food, or to the decrease in availability of natural shelter for voles. Lichens are an important winter food for bank voles (*Clethrionomys glareolus*), and another hypothesis explained the changes in regular 3- to 5-year, synchronous, high amplitude northern Fennoscandian vole dynamics by the decline in pendulous lichens (Hansson, 1999). Thus, acidification might indirectly affect vole dynamics. Voles, especially *Microtus* voles, the grey-sided vole, and the bank vole, are key species in northern vertebrate communities, and changes in their dynamics may affect many of the avian and mammalian predators dependent on voles as prey (e.g., Hanski *et al.*, 2001). The supply of bilberry (*Vaccinium myrtillus*) may also affect vole dynamics (Hörnfeldt, 2004). Bilberry is sensitive to the direct effects of  $\text{SO}_2$  (SFT, 2002) and, together with birch, is regarded as one of the most suitable bio-indicators for monitoring pollutants (Steinnes *et al.*, 2000). Bilberry is an important food item for the grey-sided vole (Kalela, 1957) and bilberry stands provide shelter for grey-sided voles against predators (Löfgren, 1995).

The densities of large mammals also declined on approaching the Monchegorsk smelter. Although this pattern seems trivial, no data on the effects of pollution on the abundance of large mammals had ever been published prior to the study by Kozlov *et al.* (2005a). During the winter the tracks of eight species of large mammal were recorded. Mountain hare (*Lepus timidus*) and red fox (*Vulpes vulpes*) were the most common, and only these two species were recorded in industrial barrens near Monchegorsk (Kozlov *et al.*, 2005a).

#### 5.3.1.2. Birds

Changes in reproduction have been documented for three abundant hole-nesting bird species in the impact zone of

the Monchegorsk smelter: redstart (*Phoenicurus phoenicurus*), pied flycatcher (*Ficedula hypoleuca*), and Siberian tit (*Parus cinctus*). Up to 80% of breeding attempts by these species failed in the most polluted habitats; as a whole, reproductive success in industrial barrens was five times lower than in background areas. In severely polluted sites, 46% of nests were abandoned in the nest building phase (9% in an unpolluted site), with 36% of nests abandoned before completing the clutch (3% in an unpolluted site) (Gilyazov, 1993).

In the 1980s, the densities of breeding birds in moderately and severely polluted habitats along the Monchegorsk pollution gradient were estimated to be 60 and 13%, respectively, of the density in the unpolluted forests (Gilyazov and Kataev, 1990; Gilyazov, 1993). The results of recent counts are in good agreement with these earlier data: bird biomass declined monotonously on approaching the pollution source and was reduced to 12.7% in the industrial barrens relative to unpolluted forests (Kozlov *et al.*, 2005a).

In the zone of declining forests (about 15 km south of Monchegorsk), bird abundance was reduced to around two-thirds of that in unpolluted forests. In particular, the densities of typical northern taiga species like tree pipit (*Anthus trivialis*), redstart, redwing (*Turdus iliacus*), song thrush (*T. philomelos*), willow warble, pied flycatcher, and Siberian tit were lower than in control sites. In this zone, the abundances of capercaillie (*Tetrao urogallus*), hazel grouse (*Bonasia bonasia*), willow grouse, and black grouse (*Tetrao tetrix*) had been reduced by 1988 to approximately half the numbers observed in 1963 (2.5 versus 4.9 ind/km<sup>2</sup>, respectively), while no reduction was recorded in unpolluted forests (7.4 versus 7.3 ind/km<sup>2</sup>) (Gilyazov, 1993). Since the 1970s, hazel grouse, *Strix* owls, eagle owl (*Bubo bubo*), Tengmalm's owl (*Aegolius funereus*), and treecreeper (*Certhia familiaris*) have not been recorded closer than 40 km from Monchegorsk and birds of prey did not nest in this area (Gilyazov, 1993).

Counts of 16 bird species around the Monchegorsk smelter demonstrated that, in spring, birds migrate to heavily polluted areas, possibly because of the earlier disappearance of the continuous snow cover, which is reduced to a third of the background depth (Kozlov, 2001). Later in the season, however, they return to the surrounding (moderately contaminated) forest habitats. As a result, during the spring migration the density of birds in heavily polluted habitats was, on an average, 1.3 times higher than in moderately polluted sites. In contrast, during the nesting period bird density in heavily polluted sites was 2.5 times lower, and during the post-nesting period 15 to 20 times lower than in moderately polluted sites (Gilyazov, 1993).

Species richness of birds in an industrial barren south of Monchegorsk, estimated by observations repeated during 1977 to 1990, was reduced by two-thirds relative to two unpolluted sites (Gilyazov, 1993). Similarly, the number of bird species recorded in 2001 by standard surveys (three 1 km-long transects per site) decreased from 11 to 15 in unpolluted forests to 3 to 4 in industrial barrens near Monchegorsk (Kozlov *et al.*, 2005a).

Species characteristic of tundra and forest-tundra, such as yellow wagtail (*Motacilla flava*), meadow pipit (*Anthus pratensis*), bluethroat (*Luscinia svecica*), wheatear (*Oenanthe oenanthe*), little bunting (*Emberiza pusilla*), and redpoll (*Carduelis flammea*), all of which are migratory species, constituted about 40% of the records at a barren site. Willow warble represented 50 to 80% of the northern taiga species complex (redstart, redwing, willow warble, pied

flycatcher, Siberian tit and brambling (*Fringilla montifringilla*) in undisturbed conditions, but at a barren site its share did not rise above 15%. Both willow warble and meadow pipit were relatively abundant, possibly because of habitat changes (replacement of coniferous forests by birch and willow bushes, along with replacement of ericaceous dwarf shrubs by grasses) that favor these species. Also, the abundance of crows (*Corvus corone*) increased in polluted habitats relative to unpolluted forests (Gilyazov, 1993).

#### 5.3.1.3. Concluding comments on birds and mammals

The most likely explanation for the observed changes in vertebrate populations near Monchegorsk is habitat deterioration. Forest decline results in a decrease in the number of nesting sites for birds; however, the increase in secondary open habitats favors certain bird species. Changes in the abundance of invertebrates (see section 5.3.2) may influence food availability for insectivorous birds – birds in the polluted area were reported to be suffering from nutritional stress (Framstad, 2002; Eeva *et al.*, 2003). The low number of microtine rodents in the damaged and moderately polluted areas is due to a decrease in the quantity of important food plants: epiphytic lichens for bank vole and possibly also for red vole, mosses for Norway lemming, and vascular plants, especially bilberry, for grey-sided vole (Kalela, 1957; Kataev *et al.*, 1994; Löfgren, 1995; Hansson, 1999; Hörnfeldt, 2004).

Although there is almost no direct evidence of pollution-induced mortality of vertebrates in the Arctic, data on the population densities of a carrion-feeding beetle, *Nicrophorus vespilloides*, indicate an increase in the mortality rates of vertebrates in moderately polluted forest habitats around Monchegorsk (Kozlov *et al.*, 2005a). Even if pollutants do not kill the birds and mammals, mortality may increase due to a reduced ability to sustain environmental stresses such as disease, low winter temperatures, and shortage of food (Heinz, 1989; Hörnfeldt, 2004). Winter mortality (e.g., Novikov, 1981) may be especially high in the contaminated habitats.

The increase in mortality, in the absence of a compensatory increase in reproduction (as reported in earlier studies), suggests that populations in the polluted (low quality) habitats are supported by migration from the surrounding unpolluted territories, and that migrants are not able to assess habitat quality in terms of toxic contamination before breeding. Moderately polluted habitats may be attractive for dispersing birds and mammals not only because of the permanent vacation of territories due to high mortality, but also because some resources are more abundant there than in the surrounding unpolluted forests. The contaminated area of about 1000 km<sup>2</sup> around the Monchegorsk smelter complex acts like a death trap for dispersing birds and mammals, and the moderately contaminated forested part of this area can possibly be classified as an attractive sink habitat (Kozlov *et al.*, 2005a).

#### 5.3.2. Effects on invertebrates

The effects of acidifying pollutants on invertebrates have been explored to a much lesser extent than many of the other components of terrestrial ecosystems, and the previous AMAP assessment (AMAP, 1998) contained no information on invertebrates. However, recent studies conducted

mainly in the vicinity of the Monchegorsk nickel-copper smelter on the central part of the Kola Peninsula provide some information on the responses of several groups of invertebrates, primarily insect herbivores, to pollution.

Empirical data collected in the vicinity of the point sources, which emit an extremely wide range of metals and other compounds, can not be used to unequivocally attribute observed effects to a specific pollutant. Nevertheless, many pollutants are considered to have a minor impact on biota around point sources. In particular, there has been no correlation found between larval weight for the moth *Eriocrania* and nickel concentrations in the larval body (Kozlov *et al.*, 2000), which agrees with the reported resistance of many insects to heavy metals (Heliövaara and Väisänen, 1993; Boyd and Martens, 1994). In contrast, both direct and indirect effects of SO<sub>2</sub> on Lapland leaf beetle (*Chrysomela lapponica*) and autumnal moth (*Epirrita autumnata*) have been confirmed experimentally (Kozlov *et al.*, 1996c; Kozlov and Selikhovkin, 1997). Therefore in this assessment changes observed in invertebrate communities are linked to both the direct and indirect effects of acidifying pollutants, primarily to changes in habitat structure following forest decline (Rigina and Kozlov, 1999). This supports the conclusion in the previous AMAP assessment (AMAP, 1998), that the deterioration of ecosystems around the smelter complexes on the Kola Peninsula is mainly due to pollution by high concentrations of SO<sub>2</sub>, leading to extreme cases of acidification.

##### 5.3.2.1. Size, individual performance, and population structure

Insect size frequently declines with an increase in pollution levels (Heliövaara and Väisänen, 1993). However, this trend has not been found in the Monchegorsk area: weight of the Lapland leaf beetle, *Chrysomela lapponica* (Zvereva *et al.*, 1995a), femur length in the autumnal moth (Ruohomäki *et al.*, 1996), and the length of the hind tibia in black fly (*Simulium pusillum*) (Kozlov *et al.*, 2005b), were independent of pollution load. The absence of a pollution effect in *S. pusillum*, together with an extremely low abundance of black fly near the Monchegorsk smelter, implies that at least some of the specimens from the most heavily polluted sites may represent occasional migrants rather than locally breeding populations. However, this explanation is not valid for *C. lapponica* or the autumnal moth as their larvae were collected in heavily polluted sites.

Despite some biochemical adaptations to pollution (Zvereva *et al.*, 2003), the individual performance of *C. lapponica* beetles collected in heavily polluted, industrial barrens was lower than in unpolluted forests; females laid less eggs and egg hatchability was significantly lower (Zvereva *et al.*, 1995a).

Although an increase in the rate of melanic specimens with an increase in pollution is well documented (Brakefield, 1987), this has not been reported for the Arctic. There were no melanic specimens among geometrid and noctuid moths collected around the Monchegorsk smelter by bait-traps during 1991 to 1994 (Kozlov, 1997). Similarly, frequencies of melanic morphs of the Lapland leaf beetle *C. lapponica* were independent of pollution load, despite pronounced variation in morph frequencies among study sites and differences between dark and light morphs in both size and performance (Zvereva *et al.*, 2002).

Changes in spatial structure of insect populations are limited to observations on their distribution among and



within host-plant individuals. Distribution of mines of two moths, *Eriocrania* spp. on birch and *Phyllonorycter strigulatella* on elder, is more aggregated near the pollution source (Kozlov, 1987, 2003). For *Eriocrania*, an increase in aggregation can be explained by higher wind speed (Kozlov, 2002) in industrial barrens, which forces females of tiny moths to walk rather than to fly between the oviposition sites. However, this explanation is not valid for *P. strigulatella*, due to much lower habitat disturbance near the power plant in Apatity than in the industrial barrens around the Monchegorsk smelter. Instead, the latter may simply concentrate on longer, presumably more vigorous, shoots, because the preference for longer shoots increased towards the emission source (Figure 5.15).




Figure 5.15. Distribution of *Phyllonorycter strigulatella* mines among shoots of its host, *Alnus incana*, with different number of leaves relative to distribution of shoots in a random sample in polluted and background sites around the power plant at Apatity. Sample sizes: polluted sites, all shoots: N=500, mine number: N=1250; background sites, all shoots: N=300, mine number: N=1000. Statistics ( $\chi^2$ ) reflect the difference between the distributions of all shoots and mined shoots (after Kozlov, 2003).

### 5.3.2.2. Changes in population densities

Some species clearly benefit from the impact of pollution. In particular, the willow feeding Lapland leaf beetle (*Chrysomela lapponica*) is generally infrequent in subarctic forests; however, outbreaks of this species, sometimes resulting in complete defoliation of willow bushes, have been detected in the vicinity of Monchegorsk and Nikel (nickel-copper smelters), and Vorkuta (a cement factory and a power plant) (Figure 5.16) (Zvereva *et al.*, 1997b, 2002; Virtanen *et al.*, 2002). Long-term studies on the Lapland leaf beetle suggest that pollution favors this species in a number of ways. First, the abundance of *Salix borealis*, which is the most favorable host plant of the leaf beetle, is much higher in the willow- and birch-dominated secondary communities around the Monchegorsk smelter than in virgin coniferous forests located at considerable distances from Monchegorsk (Zvereva *et al.*, 1995a,b). This may be especially important in relation to the findings concerning

a decrease in feeding niche breadth with increasing pollution levels: near the smelter, beetles concentrated on *S. borealis*, while in unpolluted habitats they used other willow species as well (Zvereva *et al.*, 1995b). Second, the death of large trees in polluted habitats makes the forest more open, well illuminated, and with higher summer temperatures (Kozlov and Haukioja, 1997); these microclimate changes may be beneficial for many insects in northern areas. In particular, larval survival of *C. lapponica* increased with an increase in mean June temperature (Zvereva, 1999). Third, the impact of moderate concentrations of  $\text{SO}_2$  improves host willow quality for *C. lapponica* (Kozlov *et al.*, 1996c), an effect that seems to be general for herbivores (Hughes, 1988; Riemer and Whittaker, 1989). Fourth, the overall im-




Figure 5.16. Population dynamics of Lapland leaf beetle (*Chrysomela lapponica*) at two relatively clean and two heavily polluted sites near the copper-nickel smelter in Monchegorsk. Density was assessed by the number of beetles found during 10 minute counts (conducted in three replicates at each site); bars indicate standard errors (after Zvereva *et al.*, 2002).

part of pollution creates enemy-free space for *C. lapponica* (Zvereva and Kozlov, 2000) by decreasing the abundance of some predators, especially syrphid flies, birds, and ants; in contrast, fly parasitoids are not sensitive to pollution (Figure 5.17). Thus, it seems that the release from predation pressure allows leaf beetles to increase their abundance, despite a decrease in individual performance and to utilize high-quality host plants up until the point where overgrazing causes a deterioration in host quality (termed Delayed Inducible Resistance), with a subsequent crash in the outbreak (Zvereva *et al.*, 1997a).

Pollution may interact with the natural density fluctuations of some insects and, in these situations, the effects of pollution become visible only in high-density years. This is the case with a tiny moth, *Phyllonorycter strigulatella*, the larvae of which develop in the leaves of *Alnus incana*. This was monitored around a coal-fired power plant (emitting 11 000 to 29 000 tonnes of  $\text{SO}_2$  each year) near Apatity during 1991 to 2001. The periodicity in density fluctuation was



not affected by pollution; peak densities of the leafminer were observed at both polluted and unpolluted sites in 1993 and 1999 (Figure 5.18). Densities of *P. strigulatella* showed no correlation with pollution levels between outbreaks, but increased strongly near the power plant during outbreaks. At polluted sites the density increased by a factor of 15 to 20, whereas at unpolluted sites it increased by a factor of 3 to 4, relative to the density between outbreaks (Kozlov, 2003).

Increase in density with increasing proximity to the Monchegorsk smelter complex has also been reported in leafmining moths (*Eriocrania* spp.) feeding on birches, in the moth *Argyresthia pygmaeella* the larvae of which live in willow buds (Kozlov, 1997), and in saprophagous flies (families Calliphoridae, Scatophagidae, Caloboridae, Otitidae, Sepsidae) (Zvereva, 1993a,b). Around Nikel, a density increase was found for the autumnal moth (Ruohomäki *et al.*, 1996).

The opposite – a decline in density near the pollution source – has been reported more frequently than a density increase; however, this may have resulted from the strong publication bias against the reporting of ‘positive’ results. This trend seems most common for soil mesofauna around all Kola Peninsula polluters (Koneva, 1993; Zenkova, 1999; Zenkova and Zainagutdinova, 2002). In particular, the density of spiders in the most polluted sites was around 10% of that observed in unpolluted forests around Monchegorsk (Gilyazova, 1993) and Nikel (Koneva and Koponen, 1993). Abundance of larvae of two families of soil-inhabiting beetles – click beetles (Elateridae) and soldier beetles (Cantharidae) – near the Kandalaksha aluminum smelter decreased by a factor of 10 to 20 (Zenkova and Zainagutdinova, 2002). The most pronounced decline – to 0.43%




Figure 5.17. Mean mortality of different stages of Lapland leaf beetle (*Chrysomela lapponica*) due to predators and parasitoids in two relatively clean and two heavily polluted sites near the copper-nickel smelter at Monchegorsk. Bars indicate standard errors (after Zvereva and Kozlov, 2000).




Figure 5.18. Density fluctuations for the leafmining moth *Phyllonorycter strigulatella* in polluted and background areas around the power plant at Apatity on the Kola Peninsula. Means are based on five (polluted sites) or four (background sites) site-specific values; bars indicate standard errors (after Kozlov, 2003).

of the background density – was reported for soil mites near Monchegorsk (Zenkova, 1999). Ants also decline near Monchegorsk, as revealed by direct censuses (Gilyazova, 1993) and estimates of their predatory activity against leaf beetle larvae (Zvereva and Kozlov, 2000). Among herbivores, this pattern was evident in the birch-feeding leafroller (*Eulia ministrana*) (Kozlov, 1997) and several noctuid moths (*Xestia rhaetica*, *X. speciosa*, *Eurois occultus*) (Kozlov *et al.*, 1996a). A decrease in the abundance of the autumnal moth was reported near Monchegorsk (Ruohomäki *et al.*, 1996); this, however, contradicts the density increase of the autumnal moth reported near Nikel (Ruohomäki *et al.*, 1996). Catches of human-biting mosquitoes (Culicidae) and black flies (Simuliidae) decreased near the Monchegorsk smelter by a factor of 10 to 100, presumably due to the combined action of pollutant toxicity, pollution-induced forest degradation, and a decline in vertebrate density (Kozlov *et al.*, 2005b). Among other fly species, the density decline was most pronounced in predatory groups such as Empididae (Zvereva, 1993a,b).

Although detection of a non-linear (e.g., dome-shaped) response to pollution is more difficult than a linear response, this kind of density pattern has been found in several species, such as noctuid moths (*Acronicta auricoma*, *Hyppa rectilinea*, *Diarsia mendica*, *Xestia alpicola*, *Sympistis heliophila*), geometrid moths (*Rheumaptera subhastata*, *Ematurga atomaria*), and butterflies (*Clossiana euphrosyne*, *Vacciniina optilete*) (Kozlov *et al.*, 1996a,b). The density of frit flies (Chloropidae) peaked at 7 to 17 km from Monchegorsk, being equally low in industrial barrens and in unpolluted forests (Zvereva, 1993b). Population density of the burying beetle (*Nicrophorus vespilloides*) determined by bait-trapping, peaked at 15 to 30 km from Monchegorsk. Since the abundance of vertebrates decreases towards the pollution source, this pattern indicates an increase in the beetle's food supply (carcasses of vertebrates), i.e. an increase in the mortality rates of vertebrates even in moderately polluted forest habitats (Kozlov *et al.*, 2005a).

Finally, densities of some groups of invertebrates do not seem to change along pollution gradients. This pattern was detected for the willow-feeding leaf beetle (*Phratora vitellinae*) (Kozlov, 1997), springtails (Collembola), and soil-dwelling beetle larvae (Zenkova, 1999).

### 5.3.2.3. Changes in species richness, diversity, and community structure

Species richness of different taxa or ecological groups, including moths and butterflies, ants, flies (including human-biting flies), birch-feeding insects and some other insects, does not decline with decreasing distance to the smelter at Monchegorsk (Kozlov, 1996, 1997; Kozlov *et al.* 1996a,b, 2005b; Kozlov and Whitworth, 2002). Moreover, some rare or endangered species of moths and butterflies, such as *Sympistis zetterstedti* and *Clossiana freja*, have been recorded in severely polluted habitats (Kozlov, 1996). The occurrence on the Kola Peninsula of the lunar hornet clearwing (*Sesia bembeciformis*) which had been considered extinct in Finland for decades, is only associated with industrial barrens adjacent to the Monchegorsk smelter (Kozlov and Jalava, 1994). Interestingly, both species of parasitic fly that develop in larvae of the Lapland leaf beetle, *C. lapponica*, were considered extremely rare prior to the mass occurrence reported in the impact zone around the Monchegorsk smelter (Richter and Zvereva, 1996; Disney *et al.*, 2001).

In contrast, the diversity (measured by the Shannon-Weaver index) of several groups of insect, for example moths and butterflies, or all birch-feeding insects, was reported to decrease with increasing levels of pollution, while the diversity of other groups, such as flies, did not change (Kozlov, 1997; Kozlov and Whitworth, 2002; Kozlov *et al.*, 2005b). These data suggest that acidification modifies insect communities primarily via changes in abundance, rather than through selective removal of some species.

The few studies that have been undertaken on soil mesofauna suggest that several taxa, such as earthworms (Lumbricidae) and millipedes (Myriapoda), may have completely vanished in severely polluted habitats (Koneva, 1993; Gilyazova, 1993; Zenkova, 1999). Only two species of spider, *Steatoda phalerata* and *Agyneta gulosa*, were caught in the industrial barren 2.5 km from Monchegorsk, compared to 18 species collected in forests 20 and 30 km from Monchegorsk (Koponen, 2005). However, the conclusion concerning an overall decline in the taxonomic diversity of soil mesofauna (Zenkova, 1999) near the Monchegorsk smelter must be viewed as tentative, due to deficient sampling design and because the diversity data were not corrected for sample size.

#### 5.3.2.4. Concluding comments on invertebrates

Much circumstantial evidence suggests that acidic precipitation is responsible for a decline in populations of some insects in Europe and North America (Heliövaara and Väisänen, 1993). However, densities of many insects significantly increased near point polluters, thus pollution impacts on invertebrates can not be seen as purely adverse. Although the causal reasons behind the reported changes in insect populations are far from clear, the effect is more likely to be linked to changes in host plant abundance and physiology, or with changes in habitat structure, rather than to toxicity (Kozlov, 1997). Data collected in heavily contaminated industrial barrens show that these habitats are relatively rich in insect fauna and can even serve as refugia for some rare and endangered species.

### 5.4. Critical loads of acidity and their exceedance

Several protocols under the Convention on Long-range Transboundary Air Pollution (LRTAP) require signatories to reduce sulfur and nitrogen emissions, the latest being the Protocol to Abate Acidification, Eutrophication and Ground-level Ozone which was signed in Gothenburg in 1999 and entered into force in May 2005. This is an effects-based protocol, meaning that ecosystem vulnerabilities were used in negotiations of emission reduction targets. The vulnerability of ecosystems to the deposition of sulfur and nitrogen is quantified by critical loads, 'the quantitative estimate of an exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge' (Nilsson and Grennfelt, 1988). Under the auspices of the LRTAP Convention, critical loads are calculated at national focal centers of many European countries following agreed methodologies (UBA, 2004). The data are collected, vetted, and collated by the Coordination Centre for Effects, which produces European maps

and makes the data available for integrated assessments, for example in support of emission reduction negotiations (Hettelingh *et al.*, 2001).

Parties to the LRTAP Convention mostly compute critical loads for terrestrial ecosystems (forests and semi-natural vegetation) following agreed procedures laid down in the Mapping Manual (UBA, 2004). Although the LRTAP Convention covers the region of the UN Economic Commission for Europe (UNECE) – i.e. Europe, the area of the former Soviet Union, the United States, and Canada – critical loads data for North America have not been available to date. Only recently a level-0 approach has been used to compute critical loads for terrestrial ecosystems in Canada.

The resulting critical load map for Canada represents the long-term buffering capacity (or weathering rate) in the top 60 cm (at most) of soil (Aherne and Watmough, 2005). The principal data set underlying the current map is the Soil Landscapes of Canada (SLC; scale of 1:1,000,000) database. The SLC map (and associated databases) was produced by generalizing detailed soil survey data. The level of mapping was designed to be used for broad, regional-scale assessments. Using semi-quantitative methods, soil buffering capacities (derived from percent clay and substrate type) have been allocated to soil types. Buffering capacities were vertically and spatially weighted for each soil type to derive a single average value for each mapping unit. This level-0 approach does not yet consider vegetation uptake, atmospheric inputs, or soil leaching, and represents an initial attempt at mapping critical loads for Canada. Figure 5.19 shows the acidity critical loads for terrestrial ecosystems above 60° N, both for Canada and northern Europe (note: terrestrial critical loads for Alaska and Greenland have not been mapped).




Figure 5.19. Critical loads of acidity for terrestrial ecosystems in northern Europe and Canada north of 60° N.

Exceedances of critical loads were calculated by combining the critical load maps with modeled deposition data, using the hemispheric Eulerian model DEHM. Three emission/deposition scenarios were used for the exceedance calculations (see section 2.3 for details): (i) 1990 emissions, (ii) the Current Legislation (CLE) scenario for 2010 (the CLE scenario reflects the current perspectives of individual countries on future economic development and takes into account the effects of presently agreed emission control legislation in the individual countries), and (iii) the Maximum technically Feasible Reduction (MFR) scenario for 2020 (the MFR scenario projects the scope for emission reductions offered by full implementation of presently available emis-

sion control technologies, while maintaining the projected levels of anthropogenic activities).

Estimated exceedances of the critical loads for the three scenarios are shown in Figure 5.20. The results indicate that for Canada there is no exceedance (north of 60° N) under the three deposition scenarios (i.e., not even in 1990). Although critical loads are comparable in Canada and northern Europe, deposition of sulfur and nitrogen is much smaller in Canada. The minimum critical load is about 84 eq/ha/yr and the maximum rates of sulfur and nitrogen deposition are about 30 to 40 eq/ha/yr each (not in the same place). Thus not even the combined sulfur and nitrogen deposition exceeds any critical load in this region.

In northern Europe large regions show critical load exceedances for 1990 (Figure 5.20). However, this area is much reduced assuming implementation of currently agreed emission reduction measures (CLE 2010), and would almost completely disappear assuming the MFR 2020 scenario.




Figure 5.20. Estimated exceedance of critical loads of acidity for soils for three emission/deposition scenarios: 1990 emissions, the CLE scenario for 2010 (CLE, 2010), and the MFR scenario for 2020 (MFR, 2020).



## Chapter 6

# Effects on Freshwater Ecosystems

*Brit Lisa Skjelkvåle, Julian Aherne, Tarja Bergman, Kevin Bishop, Martin Forsius, Laura Forsström, Natalia A. Gashkina, Jean-Paul Hettelingh, Dean Jeffries, Øyvind Kaste, Atte Korhola, Antti Lappalainen, Hjalmar Laudon, Jaakko Mannio, Tatyana Moiseenko, Marjut Nyman, Maximilian Posch, Ann Kristin Schartau, John Stoddard, Jouni Tammi, Jussi Vuorenmaa, Anders Wilander, and Valery Yakovlev*

### 6.1. Evidence from water quality monitoring

The previous AMAP assessment of acidification of surface water in the Arctic focused on northern Fennoscandia and the Kola Peninsula (AMAP, 1998). There were no reports of acidified lakes and rivers in Arctic Canada and Alaska. Although lakes in some areas of North America were reported to be acid sensitive there were no indications of acidification.

The geographical extent of surface water acidification was based on national lake surveys in Finland, Norway, and Sweden between 1986 and 1990 (Henriksen *et al.*, 1992). These surveys showed that the impacts from high levels of sulfur deposition were mostly limited to a distance of about 50 km from the large pollution sources. Nitrate concentrations were very low over the whole area, due to very low nitrogen deposition.

The border areas between Norway, Finland, and Russia were reported to be heavily polluted. Studies of lakes in the eastern parts of Finnmark in 1986 (Traaen, 1987) showed that the sulfate concentration had more than doubled since 1966 and was at the same stage of acidification as the most acidified lakes in southern Norway. Numerous small mountain lakes in the area were chronically acidic (pH<5). Even large lakes had little of their original buffering capacity left. Some small lakes, in particular at Jarfjorðfjellet, were too acidic to support fish. Studies in 1987 to 1989 (Traaen, 1991) indicated that large areas in Sør-Varanger would become increasingly damaged, with a loss of fish stocks, if the acid deposition increased further.

On the Kola Peninsula acidified lakes were reported around the industrial centers, and along the northern and eastern parts of the peninsula. However, at Kola, pollution from nickel and copper was a bigger problem in lakes than acidification (Moiseenko *et al.*, 1995).

An extensive inventory of small lakes in northern Finland between 1987 and 1991 showed that acidic and poorly buffered lakes were widely found in northeastern Lapland near the Norwegian and Russian borders. Sulfate concentrations here were generally higher than in other parts of Lapland (Kähkönen, 1996). A survey of small mountain lakes and brooks in northeastern Lapland in 1991 to 1993 found that the alkalinity of surface waters was lowest (mostly <50 µeq/L) in the Vätsäri area 40 to 50 km west of the Nikel smelter (Lappalainen *et al.*, 1995).

Areas sensitive to acidification and areas possibly acidified by acid deposition were quantified by calculating the critical loads for acidification of surface waters (see Box). Sulfur was regarded as the only acidifying agent as nitrate concentrations in lakes and streams in this area were very low. Critical loads for northern Fennoscandia, the Kola Peninsula, and the Spitsbergen Archipelago (Henriksen *et al.*, 1994; Lien *et al.*, 1995) were very variable, ranging from <300 to >1300 mg S/m<sup>2</sup>/yr. Most areas of Finland

### Definitions, terms, and calculations

#### Water chemistry

*The chemical composition of surface waters in an undisturbed ecosystem is mainly determined by the contribution of ions from weathering and ion exchange in the catchment and from atmospheric deposition. Atmospheric deposition is influenced by sea salts, soil dust, and long-range transported air pollutants. The chemical composition of precipitation depends on distance from the sea and anthropogenic pollution sources. It is modified by a number of processes when passing through a catchment. Such processes are biological (microbial activity, uptake by plants, release of ions through decomposition, etc.) and chemical (weathering, ion exchange, adsorption and desorption, redox processes, precipitation etc.). In general, bedrock composed of gneisses and granitic gneisses has low weathering rates, yielding waters with a low content of base cations and bicarbonate that consequently have a low buffering capacity toward acidification. Bedrock with a more basic character such as gabbro, greenstone, and schist has higher weathering rates and releases more base cations and bicarbonate, yielding surface waters with a higher buffering capacity. Biological processes usually result in modification of ion composition or removal of ions, while weathering gives a net contribution. The total sum of ions from deposition and weathering together with all the various processes occurring within a catchment determine the chemistry of the runoff water. Additional, but often similar, in-lake processes can further modify the ion composition.*

#### Acidification and naturally acidified lakes

*The pH of most natural, mineral-bearing waters falls within the range 6 to 9. pH 6 is a threshold value below which lake biology is detrimentally affected by acidification (Baker *et al.*, 1990; Doka *et al.*, 2003; Holt and Yan, 2003). pH in a lake is affected by several natural processes within the catchment, such as the rate of weathering and production of bicarbonate, mobilization and leaching of organic acids derived from humic substances (organic anions, typically indicated by dissolved organic carbon/total organic carbon levels), and input of sulfate from natural sulfide-bearing minerals. Low weathering rates, high input of organic anions, and high input of natural sulfate help to depress pH in the lake or river. Large inputs of anthropogenically derived sulfur and nitrogen that exceed the critical loads for the lake can also depress the pH to biologically harmful levels. It is not obvious, by measuring pH only, if a low pH value in a water body is due to anthropogenic or natural processes.*

#### Alkalinity

*Alkalinity is a measure of the buffering capacity of water, or the capacity of bases to neutralize acids. Alkalinity is a measure of the water's ability to resist change in pH and to neutralize*



acid inputs. Alkalinity is a more integrative indicator of lake acidification than pH. The most important buffering materials in natural waters in the Arctic are primarily bicarbonate ( $\text{HCO}_3^-$ ) and organic acids. Waters with low alkalinity ( $<20 \mu\text{eq/L}$ ) are very susceptible to changes in pH. Waters with high alkalinity ( $>200 \mu\text{eq/L}$ ) are able to resist major shifts in pH. As increasing amounts of acid are added to a water body, the buffering capacity of the water is consumed, and the pH of the water decreases (acidification). At pH 5.5, only very weak buffering ability remains, and at pH levels below 4.5 there is no alkalinity left. Alkalinity is measured by titration. An acid of known strength (the titrant) is added to a volume of a treated sample of water. The volume of acid required to bring the sample to a specific pH level reflects the alkalinity of the sample. Alkalinity can be measured in the laboratory in many different ways, making it difficult to compare results from different investigations.

### Acid neutralizing capacity

Calculated Acid Neutralizing Capacity (ANC) is an equivalent to measured alkalinity. ANC is an even more integrative and robust parameter than alkalinity in establishing good dose/response relationships between water chemistry and damage to the biological community. ANC is the parameter used as the critical chemical criterion for sensitive indicator organisms in surface waters within the international critical loads work. ANC is defined as (Reuss and Johnson, 1986): equivalent sum of base cations minus the equivalent sum of strong acid anions  $\text{ANC} = ([\text{Ca}^{2+}] + [\text{Mg}^{2+}] + [\text{Na}^+] + [\text{K}^+] + [\text{NH}_4^+]) - ([\text{Cl}^-] + [\text{SO}_4^{2-}] + [\text{NO}_3^-])$ . Waters with low ANC ( $<50 \mu\text{eq/L}$ ) indicate possible damage to biota.

### Biologically relevant chemistry

The ultimate goal of emissions control programs is biological recovery, or the return of sensitive species that have been eliminated during the course of acidification. An assessment of biologically-relevant chemical trends can only suggest that biological recovery is possible (or expected), not that it has occurred. When surface water trends are shown to be moving in the correct direction (e.g., decreases in sulfate, or increases in pH), they indicate improvement in the acid-base chemistry of lakes and streams. It is important to note that these improvements do not necessarily equate to recovery. The term 'recovery' implies that the chemistry has returned to some pre-acidified status, such as pre-industrial levels of sulfate or alkalinity; trends indicate only that surface waters are moving toward this recovered status, not that they have reached it. In the absence of good data on biological recovery, it is common to assume that biological recovery will eventually occur, after a sufficient time lag, when key chemical variables have recovered their pre-acidification levels. These key chemical variables are those that have direct toxic effects on biota (primarily hydrogen ion and aluminum) and those that ameliorate some of the toxic effects (primarily base cations like calcium). For these reasons, evaluations of chemical recovery are often focused on acidity (pH and alkalinity), aluminum, and base cations (calcium).

### Non-marine concentrations

All calculations and presentations of sulfate and base cations (sum of Ca + Mg) in this work are non-marine fractions, i.e.

the seasalt contributions have been subtracted from the total levels measured. For sulfate, what is left represents the natural background input from weathering (which is normally very low) and the anthropogenic contribution from deposition. For base cations the remaining fraction is derived from weathering. Non-marine fractions (denoted by an asterisk) of sulfate\* and base cations\* in lake and river water are calculated (see below) under the assumptions that all chloride (Cl) is of marine origin (cyclic sea salts) and is accompanied by other ions in the same proportions as in seawater. Base cations\* are in this assessment taken as the sum of non-marine Ca + Mg. All units are in  $\mu\text{eq/L}$ .

$[\text{Ca}^*]$	=	$[\text{Ca}]$	-	0.037	·	$[\text{Cl}]$
$[\text{Mg}^*]$	=	$[\text{Mg}]$	-	0.198	·	$[\text{Cl}]$
$[\text{SO}_4^*]$	=	$[\text{SO}_4]$	-	0.103	·	$[\text{Cl}]$

### Acid sensitive lakes and critical loads

Lake water chemistry gives information on sensitivity to acidification. An extremely sensitive lake typically has an alkalinity of  $<20 \mu\text{eq/L}$ , and a less sensitive lake from 20 to  $50 \mu\text{eq/L}$ . A lake with an alkalinity of  $>200 \mu\text{eq/L}$  is considered to be insensitive to acidification. The concentration of base cations can also give an indication of the acid sensitivity of the water, as the base cation concentrations directly reflect the weathering rate and bicarbonate production rate within the catchment. Surface water with low concentrations of base cations ( $\text{BC}^* < 100 \mu\text{eq/L}$ ) indicates sensitivity to acidic atmospheric inputs. Concentrations of base cations from 100 to  $400 \mu\text{eq/L}$  indicate moderate sensitivity and values  $>400 \mu\text{eq/L}$  indicate general insensitivity.

To evaluate anthropogenic acidification of lakes, the water chemistry must be evaluated together with deposition input. This is done by calculating the critical load of acidity ( $\text{CL}_{\text{Ac}}$ ) and the exceedance of the critical load based on atmospheric inputs of sulfur and nitrogen. The critical load concept is a method of estimating ecosystem sensitivity to acidic inputs (i.e., sulfur and nitrogen), and was used to prepare the two protocols to the LRTAP Convention for reducing emissions of sulfur and nitrogen in Europe: the Oslo Protocol in 1994 (UNECE, 1994) and the Gothenburg Protocol in 1999 (UNECE, 1999). Similarly, critical loads were used to guide the distribution of sulfur dioxide emissions reductions in south-eastern Canada during the 1980s and 1990s (Jeffries, 1997). The  $\text{CL}_{\text{Ac}}$  is a property of the lake and its catchment and is primarily based on weathering rates in the catchment. Since weathering is a function of bedrock geology, the sensitivity to acidification of surface waters can also be determined from geological maps. Exceedance of critical loads compares the critical load with deposition – actual or expected. When the deposition is greater than the critical load the aquatic ecosystem is expected to become damaged.

The geographical extent of surface water sensitivity to acidification in the AMAP region can also be determined from a map prepared by the Stockholm Environmental Institute based on soil type, land cover, and soil moisture. This is mainly assumed to hold for assessment of soil sensitivity, but will in general also hold for surface water sensitivity.




Figure 6.1. Distribution of (a) sulfate levels and (b) acid neutralizing capacity in arctic lakes. For densely sampled areas (Scandinavia and Spitsbergen), data were summarized into grid cells by averaging the data points (lakes).

and Norway near the smelters were quite sensitive, and critical loads for the Kola Peninsula were low. Exceedance of critical loads (based on 1990 sulfur deposition), which indicates possible surface water acidification, occurred in 70% of the county of Sør-Varanger in Norway. Critical loads were exceeded in 48% of lakes on the Kola Peninsula (Moiseenko, 1994). Small exceedances (based on 1990 sulfur deposition) were recorded in 5% of the ice-free area of Svalbard and Bear Island, but only in the northern parts (Lien *et al.*, 1995).


Long-term trends in surface water chemistry in Finland, Norway, Sweden, and on the Kola Peninsula, showed that between the mid-1980s and early 1990s, acidification had stabilized and may even have reduced slightly in some lakes. This was assumed to have resulted from decreased sulfur emissions in Europe. Some of the lakes and rivers had enough buffering capacity that they were not affected by high acid inputs.

#### 6.1.1. Current status

The current status of acidification of surface waters in the Arctic has been determined from a compilation of regional and sub-regional lake surveys. Most were undertaken in northern Fennoscandia and the Kola Peninsula, but ar-

eas of Arctic Canada, northern Russia, Alaska, Iceland, and the Spitsbergen Archipelago were also covered. Since the previous AMAP assessment on acidification (AMAP, 1998) there has been one large lake survey in the arctic part of the Fennoscandian region. This was undertaken in 1995 and included the Kola Peninsula, the northern part of Fennoscandia, and Iceland (Henriksen *et al.*, 1997a,b; Skjelkvåle *et al.*, 2001c).

This assessment compiles data from several arctic regions. Sources of data and median values for key chemical variables for different arctic regions are summarized in Table 6.1. The geographical distribution of sulfate and acid neutralizing capacity (ANC) in arctic lakes is shown in Figure 6.1. It should be remembered that compiling data in this way means that the sample population is not a statistical subset of the overall lake population. Chemical data from 605 lakes within the Canadian Arctic were compiled for this assessment. Most of the data were collected in the 1990s (Table 6.1). The variables include: pH, calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ), sodium ( $\text{Na}^{+}$ ), potassium ( $\text{K}^{+}$ ), alkalinity, sulfate ( $\text{SO}_4^{2-}$ ), chloride ( $\text{Cl}^{-}$ ), nitrate ( $\text{NO}_3^{-}$ ), plus dissolved organic carbon (DOC) and specific conductance. Some datasets contained dissolved inorganic carbon (DIC) values rather than alkalinity, and in such cases, the latter was estimated by assuming that the DIC exists entirely as bicarbonate ( $\text{HCO}_3^{-}$ ).

Table 6.1. Median values for a selection of key chemical variables including the calculated critical load of acidity  $CL_{Ac}$  and exceedance.

		n	% of total lake population	pH	BC*, $\mu\text{eq/L}$	Alkalinity, $\mu\text{eq/L}$	ANC, $\mu\text{eq/L}$	SO <sub>4</sub> *, $\mu\text{eq/L}$	DOC, mg C/L	NO <sub>3</sub> , $\mu\text{eq/L}$	CL <sub>Ac</sub> , meq/m <sup>2</sup> /yr	% of area exceeded by sulfur deposition
Northern Russia <sup>a</sup>	Taymir area	23		7.97	490		445	80	4.0			
Northern Russia <sup>a</sup>	Pechora River basin	29		6.75	196		195	53	7.7			
Northern Russia <sup>a</sup>	Lena River basin	31		7.40	393		312	37	4.0			
Northern Russia <sup>b</sup>	Kola Peninsula	460	2.3	6.45	172	79	144	36	7.6	0.1	66	14
Finland <sup>b</sup>	Lapland	184	2.1	6.81	196	116	175	35	5.3	0.2	64	8
Norway <sup>b</sup>	Nordland, Troms and Finnmark	205	1.4	6.80	145	88	106	27	1.4	0.2	101	12
Sweden <sup>b</sup>	Norbotten	641	3.5	6.84	188	114	165	30	4.0	0.4	64	3
Iceland <sup>c</sup>		39	2.1	7.30	369	441	477	10	1.0	>0.1	546	0
Svalbard and Bear Island <sup>c,d</sup>		167	30	7.54	541	492	429	82	0.5	1.9	193	5
US, Alaska <sup>e</sup>	Kenai	59		6.88	101	96	144	3	6.4	0		
US, Alaska <sup>f</sup>	CABAL	22		7.79	457	459	448	89	4.7	0		
Canada <sup>g</sup>	Yukon (YK)	96		8.09	972	692	756	55	5.6	0.6		
Canada <sup>g</sup>	Northwest Territories (NT)	167		7.89	1866	1525	1519	83	3.0	0.7		
Canada <sup>g</sup>	Nunavut (NU)	3		7.90	1577	1200	1098	275	10.6	0.8		

CL<sub>Ac</sub> is calculated by using the catchment dependent ANC<sub>limit</sub> (Henriksen and Posch, 2001) and the Norwegian background sulfate concentration for all countries, except Sweden (Wilander, 1994). Exceedance CL is calculated with S-deposition from different sources, for each country year of deposition and name of the institute providing the deposition numbers are given: Norway (1990, NILU Norwegian Institute for Air Research), Sweden (1994, SMHI Swedish Meteorological and Hydrological Institute), Finland (1990, SYKE), Kola and Iceland (1992, EMEP European Monitoring and Evaluation Programme).

Data sources: <sup>a</sup> Duff *et al.* (1998); <sup>b</sup> Henriksen *et al.* (1997b); <sup>c</sup> Skjelkvåle *et al.* (2001c); <sup>d</sup> Lien *et al.* (1995); <sup>e</sup> Newell and Mitch (1992), Eilers *et al.* (1993); <sup>f</sup> Allen-Gil *et al.* (1997); <sup>g</sup> Nahanni (NT), Tuktoyaktuk (NT), Yellowknife (NT) and Wager Bay (NU) from Environment Canada, unpub. data; region between Coronation Gulf and Great Slave Lake (both NT and NU) from Rühland and Smol (1998), Rühland *et al.* (2003); region between southern YK and Tuktoyaktuk (NT) from Pienitz *et al.* (1997a,b); Wood Buffalo National Park (NT) from Moser *et al.* (1998); Melville Island (both NT and NU) from Bronwyn Keatley (Queen's University, unpub. thesis data); Prince Patrick Island (NT), Ellef Ringnes Island (NU) and Ellesmere Island (NU) from Antoniadou *et al.* (2003a,b); Victoria Island and Axel Heiberg Island (NU) from Michelutti *et al.* (2002a,b); Baffin Island (NU) from Joynt and Wolfe (2001); Southampton Island (NU) from Mallory *et al.* (2006).



#### 6.1.1.1. Northern Fennoscandia and the Kola Peninsula

The Euro-Arctic Barents region consists of two major geological provinces: the northern part of the Precambrian shield and the Caledonian fold belt toward the north and west. Within the framework of European geology, the Norwegian, Swedish, Finnish, and Kola Precambrian is a part of the Fennoscandian or Baltic Shield. The Caledonian fold belt includes Precambrian and Paleozoic sediments. The Precambrian rocks are dominated by gneisses and granitic gneisses with low weathering rates, while the Caledonian and the Archean Karelian province also include rocks of more basic character with higher weathering rates (Lidmar-Bergström and Näslund, 2005).

The results from the 1995 Nordic Lake Survey were reported by Henriksen *et al.* (1997a,b), Henriksen (1998), and Skjelkvåle *et al.* (2001c) and form the basis of this section.

Lakes with relatively high concentrations of base cations are found in areas with basic/mafic rocks and metasediments; south-central Kola, central Finnmark, and central Lapland as well as scattered areas in the Swedish mountains. Low base cation concentrations, giving the greatest sensitivity to acidification, are found scattered all over the area and are most abundant in areas with bedrock composed of granite and granitic gneisses, i.e., the northern part of Kola, Norwegian coastal areas, the northeastern and southern parts of Lapland, and the western part of Norrbotten county. Northern Norway had the highest percentage of lakes with low concentrations of base cations; in Norway 40% of lakes had <100 µeq/L, while Finnish Lapland had only 10%. In Norrbotten and Kola respectively, 21 and 26% of lakes were acid sensitive. On the other hand, Norway also had the highest number of lakes with very high concentrations of base cations (11% of lakes had >500 µeq/L compared to 4 to 6% in the other countries).

Low alkalinity lakes (<20 µeq/L) were most abundant in the northern parts of the Russian Kola. Here, the higher content of organic matter in lakes in the eastern parts and the higher anthropogenic sulfate content in the western parts lowers the alkalinity. In the northern parts of the Nordic countries both the geological conditions and the precipitation amounts determine the alkalinity; depending on the weathering rate of the minerals in the soil and dilution by precipitation, respectively. In some regions with a high proportion of peatland, low alkalinity values in lakes are largely attributable to organic acidity. In general, low alkalinity lakes are rare in the northern Nordic countries and are scattered throughout the region as a whole.

Sulfate is considered to be the major acidifier of surface waters. High sulfate concentrations were most common in the western part of the Kola Peninsula, particularly around the smelters in Nikel and Monchegorsk. More than 10% of the lakes on the Kola Peninsula had >100 µeq/L non-marine sulfate (SO<sub>4</sub><sup>\*</sup>). High concentrations were also found scattered around the whole Euro-Arctic Barents region. Most of these lakes probably had a significant input of sulfate from geological sources, reflecting the diverse geology of the area. The eastern Kola Peninsula represented the largest area with consistently low (<20 µeq/L) sulfate levels, reflecting the low sulfur deposition in this area.

Nitrate concentrations in the lakes of the Euro-Arctic Barents region were generally very low. More than 75% of lakes had <1 µeq/L NO<sub>3</sub>. It was therefore concluded that nitrogen deposition was an insignificant acidifier of these lakes.

pH values of <5.5 were most abundant in the northern part of the Russian Kola (19% of lakes). Russian Kola

also had the highest frequency of lakes with pH levels of <6.0 (28% of lakes), followed by northern Norway (14%), Norrbotten (7%), and Lapland (7%). In general, the lakes of the Euro-Arctic Barents region are less acid than lakes further south in these countries.

There was a pronounced west to east gradient in total organic carbon (TOC) concentrations. The coastal areas of Norway usually had <2 mg C/L, while lakes with >8 mg C/L were abundant in Norrbotten, Lapland, and the eastern parts of the Kola Peninsula. The topographically flat areas with peatland accumulation on the eastern Kola Peninsula and around Bothnian Bay are reflected in high TOC levels. However, despite the high proportion of peatland areas in the north, TOC concentrations in lakes in northern regions were found to be lower than in the countries as a whole, due to the colder climate, longer soil frost period, and lower peat decomposition rate (Kortelainen, 1993). Low TOC values in mountain lakes in the border area between Norway and Sweden are due to less organic soils in the catchments and high precipitation leading to dilution.

The estimated critical loads of acidity were exceeded in all countries. The highest percentage of exceeded lakes occurred in the Russian Kola and Norway, while Sweden had the lowest percentage of exceeded lakes. The calculations were based on sulfur deposition for 1990 (Finland and Norway), 1992 (Kola), and 1994 (Sweden) (see Henriksen *et al.*, 1997a for full references of source data). In Norrbotten, the mean exceedance for 47 NILU-grids (95 percentile) using the FAB model with respect to deposition in 1990, 1997, and 2010 (in accordance with the Gothenburg Protocol) was 117, 38, and 31 eq/ha/yr, respectively.

#### 6.1.1.2. Iceland

Iceland was created by volcanic activity along the Mid-Atlantic Ridge during the last 20 million years. The volcanic rocks in Iceland are predominantly mafic. Owing to the vesicular nature of the lava with glassy crusts, they weather readily. The lake water chemistry clearly reflects this geology. Results from the 1995 lake survey (Skjelkvåle *et al.*, 2001c) showed that Icelandic lakes generally had high concentrations of base cations and high values of ANC and pH. Sulfate and nitrate concentrations were in general very low, due to low anthropogenic deposition of sulfur and nitrogen. High sulfate lakes were found in the active geological zones. Owing to the basic conditions of Icelandic lakes, they are not sensitive to acidification.

Owing to the high critical loads and the low sulfur deposition in Iceland, critical loads were not exceeded in any of the lakes studied. Hence, acidification of lakes is not a problem in Iceland.

#### 6.1.1.3. Svalbard and Bear Island

The sedimentary rocks at Svalbard and Bear Island have a diverse mineralogy that includes limestone, dolomite, gypsum/anhydrite and shales containing phosphatic nodules, and yields waters with highly variable ionic composition. The lakes were sampled in 1990 to 1992 (Lien *et al.*, 1995; Skjelkvåle *et al.*, 2001c). High sulfate lakes were related to gypsum/anhydrite in the soils and bedrock. In the northern parts of Svalbard some lakes were very acid sensitive, and in this area the critical loads were exceeded given the 1990 sulfur deposition (Lien *et al.*, 1995).

#### 6.1.1.4. New critical loads and exceedance calculations for the Euro-Arctic Barents region

Critical load calculations are used for assessment and policy work under the Convention on Long-range Transboundary Air Pollution (LRTAP) and the European Union and so are regularly updated (see section 5.4). This assessment focuses on data for northern Europe submitted to the Coordination Centre for Effects in 2004 (Hettelingh *et al.*, 2004) and new data for 104 lakes in the Kola region. The data on individual lake catchments are aggregated to the 'EMEP 50x50 km grid', which is used under the LRTAP Convention for deposition and exceedance calculations. Figure 6.2 shows the 5 percentile of the critical load of acidity on this grid system. Also shown are the critical loads for surface waters in Svalbard, which were computed and documented by Lien *et al.* (1995).

Exceedance of critical loads was calculated by combining the critical load maps with modeled deposition data, using the hemispheric Eulerian model DEHM (see section 3.7.4). Three emission/deposition scenarios were used in the calculations (see section 2.3): (i) 1990 emissions, (ii) the 'Current Legislation' scenario for 2010 (CLE 2010), and (iii) the 'Maximum technically Feasible Reduction' scenario for 2020 (MFR 2020). The same scenarios were used for calculating critical load exceedance for soils (see section 5.4).

Estimated exceedance of critical loads for the surface waters in northern Europe are shown in Figure 6.3 for the three emissions scenarios. The results indicate that implementation of presently agreed emissions reductions will reduce both the area and magnitude of exceedance substantially. However, the results also show that there is still clear exceedance of critical loads for surface waters in parts of the Kola region even after implementation of maximum technically feasible emissions reductions.




Figure 6.2. Critical loads of acidity for surface waters in northern Europe.




Figure 6.3. Estimated exceedance of critical loads of acidity for surface waters in northern Europe for three scenarios: 1990 emissions, Current Legislation for 2010 (CLE 2010), Maximum technically Feasible Reductions for 2020 (MFR 2020).

#### 6.1.1.5. Current status in Arctic Canada

The area north of 60° N in Canada (the southern boundary of the AMAP domain) contains three politically defined territories: Nunavut, the Northwest Territories and the Yukon, plus the northernmost part of the province of Quebec. The area (more than 4 million km<sup>2</sup>) comprises eight terrestrial ecozones spanning the boreal, taiga, arctic, and cordillera types that encompass the full spectrum of geological sensitivity to acidification. The largest expanses of acid-sensitive terrain (defined by a combination of bedrock and surficial geology) occur on Baffin Island and the continental mainland west of Hudson Bay. Most of the islands of the Arctic Archipelago have carbonate geology and so are not acid-insensitive.

Owing to the low solubility of the soils and bedrock characteristic of acid-sensitive terrain, associated surface waters are typically low in dissolved minerals and so the concentration of base cations indicates aquatic sensitivity. Fifteen percent of lakes investigated had base cation levels of <100 µeq/L, 20% of 100 to 400 µeq/L, and 65% of >400 µeq/L. The only extremely acid-sensitive lakes occur on Baffin Island and the central mainland straddling the border of Nunavut and the Northwest Territories. It should be noted, however, that the sample population excluded a large proportion of the acid-sensitive terrain.

Sulfur dioxide has long been acknowledged as the main air pollutant acidifying Canadian lakes (Jeffries, 1995, 1997) and is ultimately deposited on aquatic and terrestrial landscapes as SO<sub>4</sub>. Sulfate deposition in the Canadian Arctic is very low (~3 meq/m<sup>2</sup>/yr) so if atmospheric deposition is the main source of SO<sub>4</sub> in lakes, then lake concentrations should also be uniformly low. Figure 6.1 shows that lake SO<sub>4</sub> concentrations in Canadian arctic lakes are in fact spatially variable indicating the presence of geological sources in some areas, particularly Axel Heiberg Island.

Dissolved organic carbon levels in arctic lakes exhibit a broad range in concentration (Table 6.1). Approximately 25% of lakes in Arctic Canada had substantial levels of DOC (>10 mg/L), which may contribute to some acidification by naturally occurring organic acids.

Lakes having alkalinity concentrations of <200 µeq/L have traditionally been considered sensitive to acidic deposition and 26% of the samples fell into this category. The spatial distribution of low alkalinity lakes tends to confirm the sensitivity interpretation using base cations. Unfortunately, there are no alkalinity data for Baffin Island lakes.

Of the lakes sampled, 251 had sufficient data to permit a calculation of critical loads for acidity (CL<sub>Ac</sub>) using the Steady-State Water Chemistry Model (Henriksen and Posch, 2001) with an acid neutralizing capacity threshold value (ANC<sub>limit</sub>) of 40 µeq/L. The median CL<sub>Ac</sub> was 127 meq/m<sup>2</sup>/yr and 8% of lakes had values of ≤10 meq/m<sup>2</sup>/yr. The spatial distribution of three CL<sub>Ac</sub> classes confirmed that the sample lakes mostly likely to acidify occur on Baffin Island and the central mainland. Whether or not the Steady-State Water Chemistry Model should be applied in a desert environment (annual surface runoff for almost all lakes was estimated to be 150 mm or less) merits further consideration.

#### 6.1.1.6. Naturally acidic lakes in Arctic Canada

Most of the Canadian lakes fell within the pH range 6 to 9, which is assumed as the normal range for most natural, mineral-bearing waters. Only 3% of sample lakes had a pH

of <6. Furthermore, only three of 605 lakes were strongly acidic, having a pH of <5. Two of the three strongly acidic lakes are on Axel Heiberg Island (pH 3.6 and 3.8) and both had extremely high SO<sub>4</sub> concentrations (Michelutti *et al.*, 2002a). Schiff *et al.* (1991) used isotopic techniques to identify the source of the acid in the most acidic of these lakes (Colour Lake). They showed that oxidation of pyrite in the surrounding basin and oxidation of ferrous iron from an anoxic zone of the lake following spring ice-off and water column overturn caused the observed low pH and high SO<sub>4</sub> concentrations. It is not clear whether the arctic setting of these lakes is an important determinant in this acidification process. Certainly, lakes and rivers in temperate climates that are affected by acid mine drainage become acidic through similar processes. In natural temperate settings, however, acid generation by sulfide mineral oxidation tends to be self-limiting because the iron oxide weathering product coats the mineral surface rendering it less reactive. The occurrence of a naturally extremely acidic lake such as Colour Lake (and probably the other Axel Heiberg lake) is quite rare. Given the widespread occurrence of sulfide and to a lesser extent sulfate minerals, it seems reasonable to assume that some and perhaps even many lakes have been affected by acid-generating oxidation processes. The best evidence of this is probably the SO<sub>4</sub> data rather than the pH data. The third extremely acidic lake in the Canadian sample population was on Baffin Island (pH 4.66), but there are no SO<sub>4</sub> data to confirm oxidation of local sulfide minerals as the acid source.

A unique ecosystem acidification occurs at the Smoking Hills along the sea shore of Cape Bathurst in the Northwest Territories (70°14' N, 127°10' W; Havas and Hutchinson, 1983). Bituminous shales appear to have been burning for thousands of years thereby producing ground-level acidic fumigations that strongly influence the local tundra. Ponds in the area (not included in the data compilation described above) typically have pH levels of >8, but those within the fumigation zone have been acidified, sometimes to a pH of <2. The ponds exhibit elevated metal concentrations (aluminum, iron, zinc, nickel, manganese, and cadmium) and biota that are characteristic of acidic environments elsewhere. Biota in adjacent non-acidified ponds are typical of other arctic environments.

#### 6.1.1.7. Alaska

There are very few lake or stream chemistry data available from acid-sensitive parts of the Alaskan arctic. The Kenai Lakes Investigation Project (Newell and Mitch, 1992; Eilers *et al.*, 1993) characterized the major ion chemistry of over 800 lakes on the Kenai Peninsula from a statistical survey of 59 lakes in 1988. The results showed two groups of lakes: those with an alkalinity of <300 µeq/L (78% of lakes) and those with an alkalinity of >700 µeq/L. Low-alkalinity lakes had significantly lower concentrations of base cations and silica and significantly higher average concentrations of DOC than high-alkalinity lakes. Despite widespread acidic soils and bog vegetation, and resulting high DOC levels, none of the lakes sampled were acidic (minimum alkalinity 20 µeq/L).

The Chemistry and Biology of Arctic Lakes (CABAL; Allen-Gil *et al.*, 1997) project sampled 22 lakes on the North Slope of Alaska in 1992 (D. Landers, U.S. Environmental Protection Agency, unpubl. data). The results showed only two lakes with an alkalinity of <200 µeq/L (minimum 156 µeq/L); none were considered to be acidified.



### 6.1.1.8. Northern Russia, Siberia

Lakes were sampled in three regions of Siberia, northern Russia, in 1993 to 1995 (Duff *et al.*, 1998). The lakes were situated in the Pechora River basin, in the Yenisey River basin on the Taymir Peninsula, and in the Lena River basin. Only the lakes in the Taymir region were located in the same area as the large emissions source at Norilsk, but these were still around 200 km away. All the lakes (with the exception of a few in the Pechora River basin) were small, dilute, and oligotrophic with a neutral to slightly alkaline pH. Forested lakes near the mining center of Norilsk had higher concentrations of the major ions and metals; sediments in these lakes also had elevated metal concentrations (Blais *et al.*, 1999). However, all the sample lakes in this area had ANC levels of  $>200 \mu\text{eq/L}$  and so were well buffered towards acidification. Some acidic lakes in the Pechora River basin had very high levels of organic carbon or high sulfate concentrations, probably due to geological sources.

### 6.1.2. Temporal trends

Temporal trends in water chemistry can be identified using data from regular monitoring over many years or from a series of repeated lake surveys. This assessment is based on both types of data but is limited to the Euro-Arctic Barents region (i.e., the northern part of Finland, Norway, Sweden and the Kola Peninsula of Russia). Data for other areas are not available.

The Euro-Arctic Barents region is affected by long-range transboundary air pollution and by emissions from industrial centers on the Kola Peninsula. Although the pollution exhibits relatively large year-to-year variations, it has still been possible to identify a general decrease in sulfur deposition and consistently low nitrogen deposition (see section 3.3).

#### 6.1.2.1. Lakes in Finland, Norway, and Sweden

Acidification has been monitored in Finnish lakes since 1990. In northern Finland, there are estimated to be around 2100 small (4–100 hectare) headwater or seepage lakes susceptible to acidification (Forsius *et al.*, 2003). A subset of 35 small forest or mountain lakes has been monitored annually in Finnish Lapland. In Norway, 24 lakes have been monitored on an annual basis since the 1986 lake survey (Henriksen *et al.*, 1988). The monitoring is focused on areas of eastern Finnmark near the Russian border. The monitoring network includes six lakes in the Jarfjord area and 11 lakes in the county of Sør-Varanger. In addition there are six lakes in the counties of Nordland and Troms (SFT, 2005). In Sweden, eight ‘reference’ lakes have been sampled three to four times each year (Wilander, 1998).

The data from these 60 lakes for 1990 to 2004 were compiled to assess surface water trends in non-marine sulfate

( $\text{SO}_4^*$ ), non-marine base cations ( $\text{Ca}+\text{Mg}$ )<sup>\*</sup>, alkalinity, ANC, and pH. The lakes were sub-divided into three groups based on geographical location: (1) Lapland, Finland, (2) eastern Finnmark, Norway, (3) northern Norway and Sweden (Figure 6.4). Trends for each lake and each variable were analyzed using the non-parametric tests given by Hirsch *et al.* (1982). These are equivalent to a non-seasonal Mann-Kendall trend test and the Sen slope estimator (cf. Helsel and Hirsch 1995). Trend slopes for each region were calculated using the same test, but in this case each lake was treated as a ‘season’. The results give the direction of the trend from the trend slope. The detected trends are monotonic, i.e., proceed in only one direction.

#### Non-marine sulfate

Most sites showed a significant decrease in sulfate between 1990 and 2004 (Figures 6.4 and 6.5, Tables 6.2 and 6.3). The exception was a single site in northern Sweden (Abiskojaure) in which concentrations increased due to sulfur-containing minerals within the drainage basin. Lakes in eastern Finnmark near the emissions sources showed the most pronounced decrease with a trend slope of  $-1.4 \mu\text{eq/L/yr}$  (Table 6.3) based on concentration levels of  $78 \mu\text{eq/L}$  in 1990 to  $55 \mu\text{eq/L}$  in 2004 (representing a 30% decrease). Lakes in Finnish Lapland showed a trend slope of about  $-1.0 \mu\text{eq/L/yr}$  based on concentrations of  $32 \mu\text{eq/L}$  to  $18 \mu\text{eq/L}$  (a 43% decrease). Lakes in southern and central Lapland are more affected by long-range transported air pollutants from the south and so have benefited from the decrease in total European sulfur emissions. Lakes in northern parts of Norway and Sweden showed very slight decreases with an overall trend slope of  $-0.4 \mu\text{eq/L/yr}$  (Table 6.3) based on concentrations of about  $20 \mu\text{eq/L}$  to  $13 \mu\text{eq/L}$  (a 30% decrease).

#### Non-marine base cations

Base cations showed very small changes in concentration and the trends were both positive and negative (Figures 6.4 and 6.5, Table 6.3). Of the 60 lakes in this assessment, 25% had slightly significant increasing trends while 12% had significant decreasing trends. Trends for two of the three regions showed significant increases with average trend slopes of  $+0.5 \mu\text{eq/L/yr}$  for northern Norway and Sweden and  $+0.8 \mu\text{eq/L/yr}$  for Lapland. The increase in the first group was strongly dependant on the increase in a single lake (Lake Abiskojaure). Small changes and even increases in  $\text{BC}^*$  concentrations compared to  $\text{SO}_4^*$  con-

Table 6.2. Decrease in sulfate concentrations ( $\mu\text{eq/L}$ ) between 1990 and 2004 for the three sub-regions. Values are calculated from linear regressions of average values.

	$\text{SO}_4^*$ , $\mu\text{eq/L}$		Change (%)
	1990	2004	
Lapland, Finland	32	18	-43
Eastern Finnmark, Norway	78	55	-30
Northern Norway and Sweden	20	13	-30

Table 6.3. Regional trends for 1990 to 2004. Values are median slopes with significant results ( $p < 0.05$ ) in bold. Units are  $\mu\text{eq/L/yr}$ .

	n	Alkalinity		ANC		$\text{BC}^*$		$\text{H}^+$		$\text{SO}_4^*$	
		p	Theil slope	p	Theil slope	p	Theil slope	p	Theil slope	p	Theil slope
Lapland, Finland	35	<b>0.002</b>	<b>1.0</b>	<b>0.04</b>	<b>-0.3</b>	<b>0.006</b>	<b>0.8</b>	0.2	0.00	<b>0.0003</b>	<b>-1.0</b>
Eastern Finnmark, Norway	17	<b>0.004</b>	<b>0.3</b>	<b>0.0003</b>	<b>1.9</b>	0.05	-0.4	<b>0.0004</b>	<b>-0.04</b>	<b>0.0001</b>	<b>-1.4</b>
Northern Norway and Sweden	8	<b>0.007</b>	<b>0.9</b>	<b>0.001</b>	<b>1.6</b>	<b>0.007</b>	<b>0.5</b>	<b>0.002</b>	<b>-0.01</b>	<b>0.001</b>	<b>-0.4</b>




Figure 6.4. Trends in non-marine sulfate, non-marine base cations (Ca+Mg), alkalinity, acid neutralizing capacity, and pH across the Euro-Arctic Barents region for 1990 to 2004. Large circles denote the three sub-regions.




Figure 6.5. Trends in non-marine sulfate, non-marine base cations, alkalinity, acid neutralizing capacity, pH, and labile aluminum for lakes in the three sub-regions.

centrations enabled alkalinity to increase (see Figures 6.4 and 6.5). The median annual change in alkalinity ranged from +0.3 to +1.0  $\mu\text{eq/L/yr}$  for the three regions (Table 6.3). The signs of chemical recovery are not reflected in the same way in calculated ANC (Figures 6.4 and 6.5). For Lapland, the median annual change in ANC was a decrease of -0.3  $\mu\text{eq/L/yr}$  (Table 6.3). The small lakes in northern Finland are low ionic, dilute lakes (with  $\text{BC}^*$  typically  $<100 \mu\text{eq/L}$ ) that have had only modest impacts from air pollutants. Therefore, a high interannual variation in ion concentrations in these lakes reflects hydrological variations.

#### Alkalinity and pH

Increases in alkalinity are at an early stage and not really reflected in the pH values. In 37% of lakes, pH showed a significant increase (Figures 6.4 and 6.5, Table 6.3 (calculated as decrease in concentration of  $\text{H}^+$ )). This increase was most pronounced in eastern Finnmark. There is a large interannual variation in pH in lakes in eastern Lapland, due to variations in hydrology and natural organic acids. So a consistent and corresponding increase in pH cannot be expected. There is also pronounced seasonal variation, with the highest values in late summer and the lowest at the end of winter or at snowmelt.

#### Aluminum

There are few data on aluminum. In Finland, when sample pH was  $>6.2$ , the laboratories did not analyze aluminum. Moreover, at many sites inorganic bound aluminum was below the detection limit ( $<10 \mu\text{g/L}$ ). The 1995 lake survey showed that a median value for lakes in northern Finland was  $<10 \mu\text{g/L}$  (with a 90 percentile of  $10 \mu\text{g/L}$ ), and the median in northern Norway was also  $<10 \mu\text{g/L}$  (90 percentile of  $13 \mu\text{g/L}$ ). Lakes in eastern Finnmark show a clear decrease from around 20 to  $30 \mu\text{g/L}$  in the early 1990s to  $10 \mu\text{g/L}$  in 2004 (Figure 6.5).

#### 6.1.2.2. Lakes on the Kola Peninsula

A lake survey has been conducted every five years on the Kola Peninsula (1990, 1995, 1998, 2000, 2005). Six lakes with different atmospheric loads were chosen to illustrate long-term changes. Two are near (10 km) the large point sources at Nikel and Monchegorsk while the other four are over 80 km away (Figure 6.6).

Water quality in the two lakes near the point sources had improved, with decreased concentrations of sulfate (Figure 6.6), nickel, and copper. Base cation levels have also decreased at these sites. As the concentrations of sulfate, nickel and copper also decreased in the four lakes located






Figure 6.6. Trends in non-marine sulfate, alkalinity, non-marine base cations, and pH for lakes on the Kola Peninsula.

some distance from Nikel and Monchegorsk it appears likely that these decreases represent a decrease in regional sulfate and metal loads in the Kola North. There was no tendency for pH or alkalinity to increase despite the large decrease in sulfate.

#### 6.1.2.3. Swedish repeated lake survey

In each of the Swedish lake surveys (1990, 1995 and 2000) 315 lakes were sampled in Norrbotten county. An evaluation using Theil's slope showed that  $\text{SO}_4$  concentrations fell in 96% of lakes, which supports the results of other studies presented here. Alkalinity and ANC decreased in 91 and 69% of lakes, respectively. So this study shows no anticipated response of the lakes to the decreased sulfur deposition. Seven Norrbotten reference lakes studied sev-

eral times each year also had decreasing trends for alkalinity, but in all seven the ANC increased (median 0.0015 meq/L/yr). A comparison of these two studies shows that changes observed on just a few occasions may be more indicative of variation between years than a trend. Trends, as evident from more intense monitoring, are very small and so surveys with few observations over time may be insufficient to identify the trends found using more intensive studies (Wilander, 1998).

#### 6.1.2.4. Concluding comments on trends

Long-term monitoring indicates that lakes in the Euro-Arctic Barents region are recovering from acidification and that this is due to reduced sulfur deposition. However, even if the lakes have received acidic inputs and have consequently experienced a decrease in ANC, alkalinity, and pH, the lake water has not necessarily been acidified to a level at which visible damage to the biota can be expected. Lakes in eastern Finnmark near the non-ferrous metal smelters on the Kola Peninsula show the clearest signs of recovery due to reduced sulfur deposition. Other studies from the Finnish-Norwegian-Russian border area also show signs that lakes there are recovering from acidification. Surveys of small mountain lakes and brooks in northeastern Lapland in 1993 and 2000 found a significant increase in alkalinity and a decrease in sulfate concentrations. The increase in alkalinity was pronounced in the most acidic lakes (i.e., those with an alkalinity of  $<20 \mu\text{eq/L}$ ) in 1993 (Lappalainen *et al.*, 1995; Tammi *et al.*, 2003b). The recovery of lakes further from these major pollution sources (that experienced only modest impacts from air pollutants) are affected by a high interannual variation in ion concentrations and so trends are not easily detected. Although the changes seen in Swedish surface waters are thus less pronounced than in northern Norway, they are still significant evidence of improvement in the Barents region.

## 6.2. Effects of acidification on arctic biota

The first AMAP assessment (AMAP, 1998) reviewed data on the effects of acidification on the freshwater biota of arctic surface waters. One of the key conclusions of the assessment was that arctic biota live in an extreme environment – the limited temperature range, ice-free season, and productivity of the Arctic produce biotic assemblages that may require very little additional stress before they are affected. The assessment also described those characteristics of arctic biota that make them especially sensitive to acidification. The assessment concluded that, while the taxa found in arctic waters are also likely to occur in sub-arctic (and even temperate) waters, the number of different taxa in any one arctic lake or stream is relatively low – low taxa richness and low productivity create the potential for small changes in biotic assemblages to have relatively large ecosystem effects (Hobbie, 1984). Within the Arctic, the likelihood of low productivity and paucity of species increases with latitude. The assessment found that crustacean zooplankton may be absent from ultra-oligotrophic arctic lakes due to lack of food and that benthic invertebrate assemblages are generally dominated by chironomid larvae in lakes, and in flowing waters by the same sensitive insect taxa (e.g., mayflies) that dominate in northern temperate areas. The assessment also found that there are a small number of fish

species in most arctic waters – fish life spans may be very long (e.g., 25 to 40 years for Arctic char) due to slow growth and the slow accumulation of sufficient energy reserves for reproduction. Thus, arctic fauna is considered sensitive or vulnerable to anthropogenic alterations, including acidification (Hammar, 1989), principally because arctic assemblages occupy simple and labile ecosystems that undergo extreme climatic conditions and fluctuations.

Although there are relatively few (compared to temperate zones) published studies on arctic biota and acidification, research and monitoring prior to 1996 supports the characterization of arctic biota as susceptible to acidification (AMAP, 1998). The few documented effects include sensitive zooplankton (e.g., some species of *Daphnia*) being rare or absent from acidified lakes in the Jarfjord area of Norway (Nøst *et al.*, 1992); there are no other reports of acidification effects on plankton in the Arctic. Also in the Jarfjord area, Nøst *et al.* (1992) reported low abundance of acid-sensitive mayflies (Ephemeroptera) in acidified lakes; in the Dalelva catchment (Norway) the loss of sensitive benthic invertebrate taxa was attributed to low pH and elevated aluminum concentrations (Bækken and Aanes, 1990). In the Murmansk region of Russia, the abundance of zoobenthos in small acidified lakes was as low as that in lakes contaminated by heavy metals (Yakovlev, 1992). In northeastern Finland, no acidification effects were reported for the benthos, with sensitive taxa (e.g., *Baetis rhodani*, *B. lapponicus*) found throughout the area. Many researchers concluded that fish populations in the Arctic were relatively unaffected by acidification (e.g., Erkinaro *et al.*, 1992; Lappalainen *et al.*, 1995; Nøst *et al.*, 1992); deleterious effects of acidification on fish may only be found in the Jarfjord area of Norway (Hesthagen *et al.*, 1992).

### 6.2.1. Current status

More recent data support the conclusions of the first AMAP assessment (AMAP, 1998), namely that acidification effects are rare. These data are all for the areas immediately adjacent to the Kola Peninsula – there continues to be no biological data for any acid-sensitive areas of the North American Arctic.

#### 6.2.1.1. Phytoplankton and periphyton

Planktonic and epilithic diatoms are considered some of the best biotic indicators of acidification; the disappearance of acid-sensitive diatom species, and the dominance of acid-tolerant species, has long been used to quantify the long-term progression of acidification (see section 6.4). The current species composition of diatoms and chrysophytes can also be used to assess whether acidification has altered the current status of lakes and streams. There is currently no evidence of altered diatom assemblages due to acidification in the Arctic. Sorvari *et al.* (2002), for example, examined sediment cores from five lakes in Finnish Lapland and concluded that, although diatom assemblages have changed over the past two centuries, there was no evidence of changing diatom-inferred lake water pH. Similarly, Korhola *et al.* (1999) reported stable diatom assemblages from three lakes in northeastern Finnish Lapland, and concluded from their pH reconstructions that 'no substantial changes in the acidification status of lakes have occurred within the last century despite the very high local acidic deposition' (see also section 6.4).

Lakes in northeastern and northwestern Finnish Lapland have been studied for their contemporary phytoplankton fauna as a part of the European-wide project EMERGE (European mountain lake ecosystems: regionalization, diagnostics & socio-economic evaluation). One purpose of the project was to set the baseline of the ecological conditions of pristine mountain/arctic lakes in Europe for future monitoring. Even though the lakes do not appear to be anthropogenically acidified, variations in the species composition of phytoplankton indicate different responses and sensitivities to pH.

Phytoplankton were studied from 33 lakes in northern Finland as part of EMERGE. All the lakes had low biomass and low species numbers and are classified as oligotrophic or ultraoligotrophic from their chlorophyll a concentrations and phytoplankton biomass (Vollenweider and Kerekes, 1982). Most lakes were dominated by chrysophytes, which are characteristic of oligotrophic lakes with cool summer water temperatures, low alkalinity and conductivity, and neutral or slightly acid pH (Sandgren, 1988). Several lakes showed a pronounced dominance pattern, where only a few phytoplankton taxa contributed up to 80% of total phytoplankton biomass. This pattern is considered to be a consequence of stress, such as harsh climate (Willén, 2003). In the EMERGE study this pattern seemed to be more common in lakes with some additional stress (other than climate), such as high altitude or low pH. Although the studied lakes are widely distributed across Finnish Lapland, representing different environmental, altitudinal, and geomorphological settings, site-specific lake chemical properties (e.g., pH, conductivity, calcium, magnesium, sodium, alkalinity) seem to be the best predictors of species composition.

#### 6.2.1.2. Macroinvertebrates

Data on benthic invertebrates from lakes and streams suggest that, while demonstrable acidification effects are rare, sensitive taxa are common in the Arctic, and the potential for subtle (or future) effects is high.

Nearly 400 small lakes and their inlet/outlet streams have been studied in northeastern Fennoscandia (Finnish Lapland, northern Norway, and the Murmansk region). Data collected between 1990 and 1997 suggest that both anthropogenic (mineral) and natural (organic) acidification have effects on the structure of benthic assemblages (Yakovlev, 1999). These effects (e.g., declines in species diversity, relative abundance and biomass in acid-sensitive invertebrates such as *Gammarus*, as well as snails, mayflies, and stoneflies) also vary strongly with natural abiotic factors of the landscape, including lake size, hydrological type, and morphology. Impoverished benthic fauna, typical of acidified lakes, was found in roughly 25% of the acid and/or humic lakes in central and northern Lapland, and attributed to decreased pH and the toxic effects of elevated aluminum (Yakovlev, 1999). Yakovlev (2000) also reported increased dominance of the benthos by predators, and a decrease in primary consumers, with decreasing pH in Fennoscandia. Both Yakovlev (1999) and Erkinaro *et al.* (2001) reported no observable acidification effects on macroinvertebrate assemblages in northernmost Finland.

Benthic assemblages may also be strongly affected by acidic episodes in the Arctic. Hämäläinen and Huttunen (1998) used a set of 17 test streams in northeastern Finland to construct a weighted averaging (WA) model predicting minimum pH from invertebrate assemblage data, and then

applied the WA model to an additional 37 streams in the region. They found that the predicted pH values correlated most closely with the minimum pH values observed during spring snowmelt. They also found a strong correlation with longitude, suggesting stronger effects (and lower minimum pH values) near the Kola region.

Midge larvae (chironomids; Diptera: Chironomidae) have been sampled extensively in the EMERGE lakes in Finnish Lapland. The composition of benthic littoral chironomid assemblages in this region appears to be strongly influenced by pH, but the assemblages do not exhibit signs of acidification. Nyman *et al.* (2005) found that the variation in chironomid communities of 50 shallow lakes across western Finnish Lapland was explained by the factors that are likely to respond to future climate change (e.g., sediment organic content, total organic carbon, pH, and mean July air temperature). These subarctic lakes are not anthropogenically acidified, but are highly vulnerable to multiple stresses caused by climate change and ultraviolet radiation (ACIA, 2004; Rautio and Korhola, 2002) (see section 6.5).

### 6.2.1.3. Fish

Recent data from acid-sensitive regions of the Arctic suggest little evidence of widespread effects on fish assemblages, but significant effects in some highly affected areas. Hesthagen *et al.* (1998) analyzed questionnaire data from 401 lakes (236 with Arctic char, 293 with brown trout) in northern Norway near the Russian border. They concluded that only three populations of Arctic char had been lost due to acidification, while three populations of char and eight of brown trout were reduced at least to some degree. This is consistent with the larger context of acidification (and recovery) in the Nordic countries; results from the project 'Fish status of Nordic Lakes' indicate that 'fish population losses were most frequent in the most highly acidified region of southern Norway and least common in eastern Fennoscandia' (Rask *et al.*, 2000; Tammi *et al.*, 2003a). Similarly, a study of 13 rivers in northernmost Finland found no signs of acid-induced failure in salmonid reproduction and/or recruitment (Erkinaro *et al.*, 2001). Importantly, many studies of fish in acid-sensitive regions of the Arctic have focused on salmonids (e.g., Arctic char, brown trout) which may be relatively tolerant of low pH and elevated aluminum (Poléo and Bjerkely, 2000).

In a fish and water chemistry survey carried out in northeastern Finnish Lapland in 1991-1993, Lappalainen *et al.* (1995) found that the buffering capacities of small lakes and brooks were lowest in the Väsäri area. The surface waters of the area consist of an exceptionally high

number of small headwater lakes and brooks, which are sensitive to acidification due to the geochemistry of the local soil (Kähkönen, 1996). Owing to the relatively large differences in altitude over a short distance, natural obstacles to migration are common and fish populations in many lakes and brooks are isolated from each other. In some of the small lakes in the Väsäri area, the alkalinity values were critically low in 1993, and studies revealed the first signs of acid-induced fish population damage in Finnish Lapland in local minnow (*Phoxinus phoxinus*) populations. In continuing this research and monitoring, special attention has been focused on the minnow and its reproduction because of the sensitivity of the species to acidification and its frequent occurrence in the study area (see section 6.2.2.2).

## 6.2.2. Temporal trends

### 6.2.2.1. Invertebrates

Temporal trend data are available for 12 arctic lakes in Norway (included in the Norwegian national monitoring program on long-range transported air pollution (SFT, 2005)). The microcrustacean assemblages of six lakes situated in Nordland and Troms counties were surveyed in 1999, and the data indicate no or only minor impacts of acidification. One of these lakes has been followed with annual sampling and shows no signs of changing acidification status. Six lakes in the eastern part of Finnmark (Varanger Peninsula) were surveyed in 2000 and 2004 (additional data are available for 1990 to 1997 for most of the lakes); these data indicate minor to moderate impacts of acidification in the microcrustacean communities.

Lake Dalvatn is an acid-sensitive lake in Finnmark that has been monitored for zooplankton since 1990. Signs of improvement in the acidification status of Lake Dalvatn are given by the increased abundance of the acid-sensitive cladoceran *Daphnia longiremis* (Figure 6.7). This species, which is absent in acidified lakes (Keller *et al.*, 2002), was first recorded in Lake Dalvatn in 1996, and has been found each year since 1999 with increasing dominance. The presence of other acid-sensitive species of microcrustaceans has also increased in the last six years.

### 6.2.2.2. Fish

Tammi *et al.* (2003b) reported on fish data collected as part of a re-sampling of 20 lake and stream sites in the Väsäri area of northeastern Finland in 2000; these were the same sites as were sampled in 1993 by Lappalainen *et al.* (1995),





Figure 6.7. Presence of the acid-sensitive cladoceran *Daphnia longiremis* in Lake Dalvatn (Varanger Peninsula, Norway) (SFT, 2005).



who found evidence of acidification in the age-structure of minnow populations. Comparisons of the 1993 and 2000 data are shown for four sites in Figure 6.8. In all brooks and lakes, the alkalinity values were significantly higher, and the sulfate concentrations and conductivity values significantly lower, in 2000 than in 1993. The increased densities of minnow, and changes in the length distribution of the sampled fish, indicate reproductive success and recruitment of young fish at most sampling sites in the late 1990s. Although fish samples were taken in single years, the catches of electrofishing included several age-classes – the combined data on abundance and length distribution provided convincing evidence of the reproductive success of fish in the late 1990s. Despite the improvement in water




Figure 6.8. Length distributions of minnows caught by electrofishing in the Joulujärvi area (sites 6 and 7), and in the Äälisjärvi area (sites 16 and 20) in 1993 and 2000 (re-drawn from Tammi *et al.*, 2003b).

quality, sparse minnow populations at some of the study sites appear to have disappeared completely during the 1990s. At these sites, the catch for 1993 consisted of only a few, rather large (60–80 mm) individuals and no minnows were found at all by electrofishing in 2000.

Lake Otervatn in eastern Finnmark has been monitored for water chemistry and brown trout populations (using benthic gill nets) since 1986 (Figure 6.9). The lake has shown a dramatic recovery in both acid/base status and brown trout populations over the last 20 years. Brown trout catch per unit effort shows a significant increase over time, and is significantly correlated with the observed increase in alkalinity. These data are strongly suggestive of recovery from acidification.




Figure 6.9. Data for Lake Otervatn in eastern Finnmark, showing (a) alkalinity (µeq/L) and (b) brown trout catch per unit effort (CPUE, expressed in number of fish caught per 100 m<sup>2</sup> of gill net area) since 1986, and (c) brown trout populations in relation to changes in alkalinity.

### 6.3. Episodic acidification

Episodic acidification of surface streams during spring floods is a ubiquitous phenomenon all over the world, including arctic regions. Acid episodes are especially dangerous in the Arctic; contaminants deposited from the atmosphere accumulate in the snowpack during the long polar winter and are released rapidly into drainage basins during the short spring flood in acidified polluted snowmelt waters that cause a sharp decrease in pH and alkalinity. Simultaneous pulses of protons and metals may lead to episodes with extremely strong toxic effects in streams and rivers. pH in rivers often drops during spring or rain floods under natural conditions because of base-flow dilution by atmospheric precipitation, although in some cases this phenomenon can be due to the leaching of natural organic acids from soils in forested or wetland catchments (Laudon and Bishop, 1999). However, this natural pH decline can also be dramatic and may affect biota. The accumulation of anthropogenic acids ( $\text{SO}_4$ ,  $\text{NO}_3$ ) in catchments abruptly intensifies the episodic acidification in flood periods. Arctic ecosystems are especially vulnerable to acid pulses during spring floods. After polar night, the vulnerability of arctic biota to acid and toxic impacts is much higher. Situations in which two or more stressors occur simultaneously, thereby multiplying the risk for aquatic life, are dealt with in more detail in section 6.5. Episodic pH decline has been observed in several arctic regions, for example, Sweden, Finland, and Russia (Kinnunen, 1992; Moiseenko, 1999; Laudon *et al.*, 2000; Moiseenko *et al.*, 2001).

How pH episodes form depends on the conditions of water formation. Catchments in the Arctic are mostly of the tundra or forest type, although hilly and mountainous catchments also occur. The mechanism of pH depression in surface waters is determined, as a rule, by the interaction of several factors capable of causing acidification. Published reports emphasize five factors that contribute to depressed ANC during flood time: (1) dilution from increased discharge; (2)  $\text{H}_2\text{SO}_4$  and (3)  $\text{HNO}_3$  derived from precipitation or natural sources; (4) organic acids derived from watershed soils or wetlands; and (5) HCl derived from 'salt-effect', i.e., reactions within catchment soils (e.g., Jeffrey *et al.*, 1992).

#### 6.3.1. Acidic episodes in the Kola region

Acidic episodes have been studied in the Kola North, where a high level of anthropogenic sulfur deposition has been recorded. The decrease in pH during spring flood relative to the pre-episode period (i.e., the winter low-water period) was examined in 21 streams (Moiseenko, 1999; Moiseenko *et al.*, 2001). The decrease was most noticeable in streams where the pre-episode pH was high, while in streams that were chronically acidified (with pH values during the low-water period of  $<6$ ) the pH drop was insignificant owing to the similarity between the chemistry of water in acidified streams and atmospheric precipitation (Figure 6.10). This is in general agreement with data collected in Europe and North America (Wigington *et al.*, 1992). The period of pH depression during spring flood in this part of the Arctic is short and rarely exceeds five to seven days.

Acidic episodes facilitate the discharge of metals into streams and rivers. It is well known that water acidification causes an increase in the concentrations of labile metal forms (Dillon *et al.*, 1988; Nelson and Campbell, 1991; Jef-

fries, 1997). In arctic regions, an abrupt drop in water pH in a short flood period is accompanied by a pulse of metals, especially in their ionic forms, and it is this that generates the greatest hazard for aquatic organisms. The leaching of metals during flood (2–3 weeks) can account for up to 75% of their total annual load (Moiseenko, 1999; Moiseenko *et al.*, 2001). Data on streams of the Kola North showed that in the periods of pH depression during floods, the total metal concentration increases in all types of creek, notwithstanding dilution by snowmelt water. The total concentration of aluminum in water increased by 50–88%. In the most acidified tundra stream studied, an increase in the concentration of its labile form was accompanied by a drop in the form bound with organic complexes.

#### 6.3.2. Acidic episodes in the Dalelva catchment in eastern Finnmark, Norway

Dalelva is a small (3.2 km<sup>2</sup>) undisturbed catchment dominated by heathland and mountains, in Jarfjord in eastern Finnmark, Norway (69°45' N, 30°23' E). Vegetation comprises birch forest to an elevation of about 150 m with heath and moorland above. Small lakes cover about 15% of the catchment. The catchment is usually covered with snow for six to seven months of the year, approximately from mid-October to late May, and receives relatively little precipitation (500–600 mm/yr). At the adjacent meteorological station (Lanabukt) about 45% of the annual precipitation was accumulated as snow during 1990 to 2000 (Kaste and Skjelkvåle, 2002). The extensive snow accumulation has a large impact on the seasonal runoff pattern, which is characterized by very low flow during winter and a distinct snowmelt flood in May to June.

The large seasonal variations in streamwater flow also have big impacts on water chemistry (Kaste and Skjelkvåle, 2002; SFT, 2005). During the long cold winters, when precipitation is accumulated as snow, the stream is dominated by baseflow with relatively high solute concentrations




Figure 6.10. Water pH depression during spring flood versus pH during the low-flow period in streams of the Kola North. The upper line shows the relationship in streams where the minimum pH during high flow was  $<5$ , and the lower line where the minimum pH during high flow was  $>5$ .

(Figure 6.11). Concentrations of major ions and nitrogen compounds (not shown) tend to build up during winter and then decrease rapidly to initial levels at the start of the snowmelt flood. This dilution of base cations during snowmelt also causes a rapid decrease in streamwater pH, from the normal level of around 6.0 to 6.5, to values of around 5.5 (Figure 6.11). Sea-salts accumulated in the snowpack during winter are eluted from the snow in the early melting phase, resulting in an annual peak in chloride concentrations immediately before the onset of the main snowmelt flood (Figure 6.11).

After snowmelt, streamwater flow and solute concentrations return rapidly to normal summer levels. The Dalelva catchment usually experiences a short and intense growing season that lasts approximately four months. During summer, air temperatures can be relatively high and owing to the relatively low precipitation amounts, streamwater flow is usually very low at this time.

The overall water chemistry at Dalelva is characterized by relatively high concentrations of non-marine sulfate

owing to industrial  $\text{SO}_2$  emission sources on the Russian side of the border. During 1990 to 2000, however, streamwater concentrations of non-marine sulfate declined by 35% as a result of reduced  $\text{SO}_2$  emissions (SFT, 2005). Dalelva is moderately affected by humic substances; average concentrations of total organic carbon between 2001 and 2003 were in the range 3.7 to 4.4 mg C/L. The greatest influence of total organic carbon is often associated with the onset of snowmelt. Streamwater concentrations of nitrate are relatively low, with annual peaks up to 70 to 100  $\mu\text{g N/L}$  prior to snowmelt, and growing season values typically below 5  $\mu\text{g N/L}$ .

### 6.3.3. Acidic episodes in northern Sweden

A large, multi-investigator project in northern Sweden developed the Boreal Dilution Model (BDM) to quantitatively distinguish the natural and anthropogenic mechanisms that drive episodic decline of ANC and pH during hydrological events (Bishop *et al.*, 2000; Laudon *et al.*, 2000). The BDM identifies the anthropogenic component of episodic ANC decline ( $\Delta\text{ANC}_{\text{poll}}$ ) from relative differences in the runoff dynamics of base cations and anthropogenic acid anions during episodes. Snow chemistry data are not used in the BDM. However, comparison of snow chemistry with the results from over 50 applications of the BDM in catchments from the alpine zone to the coast of northern Sweden sampled between 1991 and 1999 revealed a strong relationship between  $\text{SO}_4$  in snow and the anthropogenic impact on the subsequent spring flood. This suggests an immediate and proportional response in spring flood acidification to changes in winter  $\text{SO}_4$  deposition. Nitrogen is not a factor in the spring flood of this nitrogen-poor region (Laudon *et al.*, 2000).

Trends in anthropogenically driven episodic acidification were analyzed in five streams from northernmost Sweden between 1990 and 1999 using the BDM (Laudon and Hemond, 2002). Although there was no significant change in annual average stream water chemistry, the anthropogenically driven episodic acidification associated with spring flood runoff decreased by 40 to 80%. A strong correlation between winter  $\text{SO}_4$  deposition and the anthropogenic component of episodic acidification in these five streams suggests that future reductions of acid deposition will further improve the spring flood acidification situation in northern Sweden. These results also indicate that reduced emissions of acid precursors have generated significant improvements in the surface water chemistry during episodes associated with spring runoff in northern Sweden.

While the data requirements of the BDM are too big for regional assessments, the correlation between snow  $\text{SO}_4$  and  $\Delta\text{ANC}_{\text{poll}}$ , together with a relatively consistent amount of snowmelt in the peak of spring flood creates the basis for the more empirical 'one point BDM' (pBDM, Laudon *et al.*, 2004). This model uses widely available lake chemistry measurements, and can thus provide a more synoptic view of how human impact on spring melt ANC in northern Sweden has responded to changed  $\text{SO}_4$  deposition. This is a region where many surface waters experience low ANC and pH in conjunction with hydrological episodes despite a relatively low annual acid deposition load. The largest episodic ANC declines and greatest biological effects in the region are associated with spring flood (Laudon *et al.*, 2000).




Figure 6.11. Surface water runoff from the Dalelva catchment, northeastern Norway, 2001–2003. (a) streamwater flow and pH, (b) non-marine concentrations of sulfate and base cations, and (c) chloride and non-marine sodium.






Figure 6.12. Intensity of anthropogenic episodic acidification in northern Sweden during the 1970 spring flood and the 1990 spring flood, and projected for the 2010 spring flood. Assessed using the acidification index of the Swedish Environmental Protection Agency and the pBDM model (Laudon and Bishop, 2002).

Sulfur deposition in northern Sweden peaked in 1970 (Mylona, 1996) and had declined by 65% by 1990 (S. Mylona, Norwegian Pollution Control Authority, unpubl. data). A further 55% reduction relative to the 1990 level is expected by 2010 in accordance with the 1999 Gothenburg Protocol to the LRTAP Convention. Winter chemistry data suitable for use in the pBDM was available for 1240 lakes sampled in conjunction with national monitoring (Figure 6.12). Application of the pBDM to these lakes using the 1990 and 2010 data projects a clear recovery (Figure 6.12, Table 6.4). In 1970, 54% of lakes in Sweden were subject to significant human impact (classes 3–5, Table 6.4). By 1990, only 14% of the lake population was more than moderately impacted by acid deposition during spring. By 2010 that number is projected to be 3%. Reductions in acid deposition have led to rapid and substantial chemical recovery from spring flood acidification in arctic waters, including 70 000 lakes and 1 000 000 km of watercourses in northern Sweden (Laudon and Bishop, 2002). These results are based on the strong correlation between  $\text{SO}_4$  concentration in snow and the anthropogenic component of spring flood ANC decline (Laudon *et al.*, 2004). Laudon *et al.* (2004) estimated that the 65% reduction in sulfur deposition between 1970 and 1990 reduced the area of acidified spring floods across 250 000 km<sup>2</sup> of northern Sweden by 75%.

A large and rapid reduction in the anthropogenic influence on spring flood acidity does not mean that spring ANC/pH decline disappears. A large spring flood ANC decline (equivalent to ca. 50%) is a natural feature of aquatic ecosystems in northern Sweden. The relative increase in organic acids during spring flood also contributes to a natural pH decline (Ivarsson and Jansson, 1995; Laudon *et al.*, 1999).

The difficulties in assessing episodic acidification have contributed to the focus of previous acidification recovery assessments on changes in average lake conditions. The failure to account for an episodic response may greatly underestimate the immediate benefits of reducing acid deposition. Furthermore, since spring flood responds directly to snow acidity in this region that is not chronically

Table 6.4. Extent of anthropogenic acidification in northern Sweden (Laudon and Bishop, 2002).

Acidification class	ANC <sub>p</sub> / ANC <sub>pi</sub>	Acidification index	Percentage cover		
			1970	1990	2010
1	1.0 - 0.75	not significant impact	20	63	91
2	0.75 - 0.50	moderate impact	26	23	6
3	0.50 - 0.25	large impact	18	6	1
4	0.25 - 0.10	very large impact	7	1	1
5	<0.10	extreme impact	29	7	1
1-2			46	86	97
3-5			54	14	3

ANC<sub>p</sub>: peak in ANC; ANC<sub>pi</sub>: pre-industrial peak in ANC

acidified, delays in deposition reduction translate directly into recovery delays. Important as these results are for northern Sweden where large investments are made each year to remediate acidification in spring flood by liming, it is very likely that extensive regions in the boreal zones of North America, Europe and Asia may be similarly sensitive to changes in winter  $\text{SO}_4$  deposition.

#### 6.3.4. Concluding comments on episodic acidification

In arctic regions, episodic acidification is a ubiquitous natural phenomenon during spring flood that has been intensified by acidic deposition. It develops swiftly due to the pollutants accumulated in the snowpack over winter being released rapidly into drainage basins during snow melt. The dominant factor in the acidification mechanism is the type of stream catchment. Replacement of hydrocarbonates by stronger acids and dilution are the most apparent and well-known factors and these affect acid episodes in all streams. In forest and wetland streams, organic acids also contribute strongly to pH decline. In remote coastal tundra areas, HCl is the dominating factor because of ion-exchange processes in catchments (adsorption of marine aerosol Na<sup>+</sup>).

A general pattern in the behavior of metals induced by acidification is an increase in their concentration, redistribution toward the most toxic ionic form, and pulses of metals during the period of episodic acidification. An abrupt increase in the ionic form, which is the most toxic form for biological systems, together with low pH causes a toxic stress for water inhabitants. In polar regions, the maximum stress for biota occurs during spring flood periods, when pH is at a minimum and the concentration of ionic forms of metals is at a maximum.

A model based assessment on data from five streams in northern Sweden indicated that reduced emissions of acid precursors have generated significant improvements in surface water chemistry during episodes associated with spring runoff. Although there was no significant change in the annual average stream water chemistry at these sites, the anthropogenically driven episodic acidification associated with spring flood runoff decreased by 40 to 80% between 1990 and 1999. A regional scale model application indicated that the 65% reduction in sulfur deposition between 1970 and 1990 has reduced the area of acidified spring floods across 250 000 km<sup>2</sup> of northern Sweden by 75%. It is also likely that future reductions in acid deposition will further improve the spring flood acidification situation in this region.

The difficulties of assessing episodic acidification have contributed to many previous acidification recovery assessments having focused on changes in *average* lake conditions. The failure to account for an episodic response may greatly underestimate the immediate benefits of reducing acid deposition. Also, since spring flood responds directly to snow acidity in those northern regions that are not chronically acidified, delays in deposition reductions translate directly into recovery delays.

## 6.4. Evidence from paleolimnological studies

Monitoring of northern lakes often started after industrial pollution had been ongoing for tens of years. In the absence of long-term data on water quality, as is often the case in the Arctic and subarctic, paleolimnological reconstructions using microfossils preserved in lake sediments provide a powerful chronology of acidification history and recovery and a tool for separating anthropogenic impact from the natural pH succession. Microfossils are surprisingly accurate proxy sources for reconstructing past environmental conditions. Modern distributions of organisms that preserve well in lake sediments can be related to the hydrochemistry of the water body using a large set of lakes ('training set') and environmental optima and tolerance ranges for the individual species. These ecological optima and tolerance ranges can then be applied to fossil communities by means of mathematical calibration (transfer) functions, which enable changes in the hydrochemistry of a lake to be quantitatively determined from the fossil assemblages deposited in the sediment over a certain time period. In particular, the strong linkage between diatoms and lake-water pH has long been recognised (for a historical review, see Battarbee *et al.*, 2001), which is why diatoms have been widely used as indicators in bio-monitoring present and past changes in the acidification status of surface waters. Using paleolimnological techniques, it

is possible to obtain a temporal perspective on the lake ecosystems with regard to acidification (Smol, 1992).

In the previous AMAP assessment (AMAP, 1998), paleolimnological assessments of acidification in arctic Canada and Alaska showed that the lakes had been unproductive throughout their entire existence and that long-term natural acidification was still occurring. However, anthropogenically-induced acidity was reported to have affected the present-day acid-sensitive plankton species, invertebrates, and fish on the Kola Peninsula and in the neighboring areas of Finland and Norway (AMAP, 1998). Critical loads were reported to have been exceeded in large areas of northern Finland and Norway due to the low critical load values in these systems (the lakes in the region are characterized by a low buffering capacity and are sensitive to acidification, see section 6.1) and the influence of emissions from industrial areas on the Kola Peninsula. At present, the diatom-based pH-reconstructions cover extensive areas of arctic Fennoscandia, the Kola Peninsula, Siberia (Norilsk), Svalbard, and arctic Canada.

### 6.4.1. Millennial trends in lake acidification

#### 6.4.1.1. Fennoscandia and the Kola Peninsula

Natural long-term acidification is a common feature of lakes in cold environments with thin soils and acid bedrock in the catchment. Most of the available pH reconstructions are based on changes in the subfossil diatom flora, since diatoms are very responsive to changes in acidity (e.g., Weckström *et al.*, 1997; Bigler and Hall, 2002). Seppä and Weckström (1999) studied the acidification history of Lake Tsuolbmajavri, located just above the present pine treeline in northwestern Finnish Lapland. The current pH of the lake is around 7.4. Diatom-inferred pH (hereafter referred to as DI-pH) was observed to have decreased slowly and gradually throughout the Holocene (i.e., the last 10 000 years) (Figure 6.13). The periods of slightly more evident drops in DI-pH values were related to climate-driven changes in soil-forming processes and catchment vegetation patterns, for example the immigration of pine and the initiation of paludification. However, the influence of peatlands on the acidification status of lakes is probably less clear in northern Fennoscandia than in more southerly boreal environments owing to the less acidic nature of the mire vegetation and water in the northern rich fens (Korhola *et al.*, 2002; Sjörs and Gunnarsson, 2002). A similar slow and gradually decreasing pH trend was reconstructed from another lake (Lake Toskaljavri) in the barren tundra region of Finnish Lapland (Figure 6.13), with a pH decrease of ca. 0.3 to 0.4 pH units during the Holocene (Seppä *et al.*, 2002).

An even weaker trend in natural acidification was evident in data from Solovieva and Jones (2002), who studied the Holocene history of a small upland lake (Lake Chuna) on the Kola Peninsula (Figure 6.13). The lake water is currently dilute, clear and slightly acidic with a pH of around 6.4. The lake experienced slow natural acidification in the early Holocene, with the acid-base balance achieved about 4000 years ago after which there appeared to have been no further acidification. By applying the diatom models developed by Weckström *et al.* (1997) and Solovieva (2000), Grönlund and Kauppila (2002) recorded a similar slight trend of progressively declining DI-pH towards the present in Lake Soldatskoje, a small tundra site located in the northern coastal area of the Kola Peninsula. Similarly,




Figure 6.13. Diatom-inferred Holocene pH histories of lakes in arctic Fennoscandia and the Kola Peninsula. Toskaljavri (a) is redrawn from Seppä *et al.* (2002), Tsuolbmajavri (b) from Seppä and Weckström (1999), Lake Chuna (c) from Solovieva and Jones (2002), and Njargajavri (d) from Sarmaja-Korjonen *et al.* (2006). Note the different scales for pH and time (cal. BP means calibrated years before present).

Bigler *et al.* (2002a, 2003) found that two small subarctic lakes near Abisko in northern Sweden had gone through a slow long-term acidification during the Holocene from about pH 7.2 to 6.8 and 7.2 to 6.5, respectively. No evidence was found that changes in land-use or reindeer herding would have affected the acidity status of these lakes during the last 1000 years.

Korsman (1999) reconstructed the late Holocene pH history from five currently acidic lakes in northern Sweden using diatoms and reported a slight natural acidification trend beginning thousands of years ago as a result of soil-forming processes and natural changes in vegetation. One of these lakes is within the AMAP region. In a recent study, Sarmaja-Korjonen *et al.* (2006) reported relatively strong mid-Holocene acidification of around two pH units from 7.5 to 5.5 in Lake Njargajavri, a small shallow lake in Finnish Lapland (Figure 6.13). The reasons for the rapid decline in pH may be due to specific characteristics of this poorly buffered, acid (pH 5.3) and oligotrophic lake and its catchment that made the system particularly susceptible to climatically-induced changes in pH. Climate is known to modulate lake acidity through links between lake ice cover, primary productivity, and DIC dynamics (Wolfe, 2002, see section 6.5.1).

#### 6.4.1.2. Concluding comments on millennial-scale acidification

Millennial-scale changes in lake water acidity in arctic Fennoscandia and the Kola Peninsula are mainly slow or non-existent during the Holocene. Excluding the initial transient alkaline period following deglaciation evident at some sites, Korhola and Weckström (2004) estimated that the long-term natural rate of pH decline in arctic lakes in this region has been around 0.005 to 0.01 pH units per 100

years, which is considerably less than the corresponding 0.01 to 0.02 pH units per 100 years in boreal lakes. Some arctic lakes may, however, be especially vulnerable to changes in climate, which can modulate pH.

#### 6.4.2. Recent acidification

##### 6.4.2.1. Fennoscandia and the Kola Peninsula

In 1990, critical loads for surface waters in northern Europe were exceeded almost everywhere. Henriksen *et al.* (1997a,b) reported that critical loads of acid deposition were exceeded in 50 to 70% of lakes on the Kola Peninsula and in the Norwegian–Russian border areas. Regional geochemical mapping from 1992 to 1998 demonstrated that although the local influence of industry on the Kola Peninsula can be seen, distance from the coast dominates the distribution of pH and sulfur trends in lake water at the regional scale (Reimann *et al.*, 2000a).

A quick and efficient paleolimnological means to establish the extent of lake acidification at the regional level is the so called ‘top–bottom approach’. From a number of lakes a single sediment sample is analysed (e.g., for diatoms) from the top of the sediment layer, to represent present-day conditions, with another taken from deeper down the core to represent pre-industrial conditions. By comparing these samples it is possible to estimate the extent of the change in species assemblages and thus the DI-pH values since pre-industrial times. This method was applied to 118 northern Swedish lakes (Korsman, 1999) and to 32 lakes around the smelters on the Kola Peninsula (Weckström *et al.*, 2003). The results indicate that in the majority of the Swedish lakes DI-pH was unchanged, while on the Kola Peninsula the decrease in DI-pH occurred only within the immediate vicinity of the smelters. These diatom data do not sup-



port the hypothesis of large-scale modern acidification in northern Sweden, nor the widespread acidification of arctic lakes due to sulfur pollution from the Kola smelting and mining industries. However, in four lakes out of 15 studied by Dauvalter (1997) in Finnish Lapland, acidification was postulated to have altered the geochemical cycling of potentially harmful metals by dissolution from sediments back to water or by reducing the adsorption of metals onto sedimenting particles.

In addition to these broad-scale top-bottom studies, several downcore pH reconstructions covering the past 200 to 300 years have been made for a number of individual lakes (see Figure 6.14). Collectively, these studies indicate that no substantial changes in DI-pH have taken place. The next paragraph summarises some of these studies.

Data from three small, potentially acid-sensitive (autumn alkalinity values of 20, 20, and 60  $\mu\text{eq/L}$ ) lakes, two in eastern Finnish Lapland 40 km west and 150 km southwest from the Nikel smelter and a reference lake in western Lapland, suggested no substantial changes in DI-pH despite the relatively high acid deposition in the east (Korhola *et al.*, 1999) (Figure 6.14, lakes A-C). The other lakes (Figure 6.14, lakes D-N), dilute clearwater lakes with varying geology, were not chosen to evaluate their acidification history and so their sensitivity to acidification was not a major criterion for selection. Sorvari *et al.* (2002) did not find change in DI-pH over the last 200 years in five tundra lakes in northwestern Finnish Lapland (Figure 6.14, lakes H-L) investigated using high-resolution sediment sampling. Similarly, Weckström (Environmental Change Research Unit, University of Helsinki, unpubl. data) found no decrease in DI-pH in two lakes in western Finnish Lapland (Figure

6.14, M-N), in fact one even showed a slight increase in DI-pH since 1800 AD.

Sediments representing the industrial time of Lake Chuna, a small upland lake on the Kola Peninsula about 30 km from the Monchegorsk smelter, were studied by Moiseenko *et al.* (2000) and Ilyashuk and Ilyashuk (2001). Moiseenko *et al.* (2000) found that toward present times the acidobiontic diatom taxa (as well as abnormal forms of some diatom species) increased as species diversity decreased, these changes running in parallel with the start of heavy metal accumulation due to industrial development of the region. Ilyashuk and Ilyashuk (2001) studied changes in the subfossil benthic invertebrate (Diptera: Chironomidae) communities in the sediments of Lake Chuna. They concluded that compositional changes in the assemblages were due to inputs of airborne contaminants and climate change. According to their interpretation, the first changes that took place in chironomid communities around 1950 were caused by the decrease in pH and the accumulation of heavy metals in bottom sediments. In the uppermost sediment layers climate change may have lead to the decrease in the predominant species, to increased taxon evenness, and thus to an increase in species diversity over the last two decades. Solovieva and Jones (2002) studied the same lake using modern quantitative approaches but did not find any significant decline in DI-pH over recent decades/centuries. This discrepancy between the pH reconstructions for Lake Chuna from the two different diatom studies is still to be resolved. It may in part result from methodological differences in reconstructing pH. Nevertheless, over the last 100 years the sediments have accumulated some toxic elements (lead, nickel, copper, cobalt, and cadmium) despite their




Figure 6.14. Diatom-inferred recent pH-histories of 14 lakes and ponds in northern Fennoscandia. Note the different scales for age and pH. Figures A-C are from Korhola *et al.* (1999), D-E from Erola (1999), F-G from Bigler and Hall (2003), H-L from Sorvari *et al.* (2002), and M-N from J. Weckström (Environmental Change Research Unit, University of Helsinki, unpubl. data).

low concentrations in water. The increasing appearance of deformations in diatom and chironomid specimens towards recent times may indicate that the accumulation of heavy metals has had an adverse effect on the lake biota.

In their study of two bays of Lake Imandra on the Kola Peninsula (Monche Bay in the near vicinity of the Monchegorsk copper-nickel smelter and Kunchast Bay, an internal reference site located about 90 km southwest from the smelter), Ilyashuk *et al.* (2003) found an increase in the metal contamination of the sediments in Monche Bay that was accompanied by marked changes in chironomid communities towards more toxic-tolerant species and a decline in diversity and total abundance. Also, the presence of increasingly frequent morphological deformities among the detritus-eating *Chironomus* and the decreased Benthic Quality Index (BQI) both served to indicate increasingly toxic conditions for the biota. At the reference site (Kunchast Bay) no obvious changes in chironomid communities were noted that could be connected to metal contamination and acidification. The mouthpart deformities of *Chironomus* may reflect a switch from non-genetic stress in midge larvae to an increasing amount of stress at genetic level under prolonged stress conditions (Ilyashuk *et al.*, 2003).

Dauvalter and Rognerud (2001) studied the impact of heavy metals on the watershed of the Pasvik River, the largest river system in northern Fennoscandia. They found increased concentrations of heavy metals in recent sediment layers in the lower reaches of the river. This was presumably due to the atmospheric emissions of nickel, copper, cobalt, zinc, cadmium and mercury from the smelters and to wastewaters from tailing dams and mines. In the upper river reaches no significant changes in vertical distribution of heavy metals were apparent.

The geochemical and biological data from the Kola Peninsula suggest localized effects of pollution from the smelter industries within a few tens of kilometers from the actual emission sources. However, no precise dates were available for the sediment cores studied by Dauvalter and Rognerud (2001) and Ilyashuk *et al.* (2003) and the time resolution of the samples (1 cm sediment slices) was not high enough to allow estimates of any recovery.

#### 6.4.2.2. Siberia

Using the top–bottom approach, Michelutti *et al.* (2001) found that diatom assemblages in the sediments of 17 lakes had experienced relatively little change since pre-industrial times in the Norilsk area, in Russian Siberia. Lakes seemed well buffered against acidification due to the surrounding alkaline bedrock and overlying glacial deposits. According to their investigations, the effects of the massive mining activities on the water quality of these lakes have been minimal, and the alkaline nature of the lake water has resulted in the incorporation of the insoluble metallic complexes into the lake sediments. However, the mining activities may have caused increased erosion, which had altered the species assemblages to some extent (Michelutti *et al.*, 2001).

There is no evidence of widespread lake acidification during the industrial period in the Usa Basin of the East-European Russian Arctic according to the subfossil diatom assemblages studied by Solovieva *et al.* (2002). However, they did find evidence of alkalization of the lakes due to atmospheric deposition, as shown by the increased DI-

pH values towards the present. Similarly, Solovieva *et al.* (2005) did not find any detectable signs of acidification in two lakes within this region studying both diatoms and chironomids from the high-resolution sediment record extending back to about 1800.

#### 6.4.2.3. Svalbard

Elevated levels of atmospheric contaminants such as spherical carbonaceous particles (from fossil fuel combustion), polyaromatic hydrocarbons, polychlorinated biphenyls, and possibly lead from both long-range and local sources are recorded in Svalbard lake sediments for the past 30 to 40 years (Rose *et al.*, 2004). Lakes in Svalbard are potentially particularly sensitive to acid deposition as they are located on deep permafrost and there is little groundwater interaction (Betts-Piper *et al.*, 2004). Exceedance of critical loads for acidity (Lien *et al.*, 1995) also suggests that acidification effects are possible, but no ecological impacts on diatom (Jones and Birks, 2004) or chironomid (Brooks and Birks, 2004) assemblages from the lake sediments have been observed. Trends in DI-pH values in five Svalbard lakes have been stable over the past few centuries. However, top–bottom analysis of chrysophycean stomatocysts showed marked shifts in assemblages in most of the lakes studied. Lakes on granitic bedrock, which are presently acidic, may have become more acidic over time. On the other hand, lakes on carbonate bedrock that are presently alkaline appear to have increased their pH in recent times. These changes may be related to atmospheric contamination from local and remote sources, but climate change may also play a role (Betts-Piper *et al.*, 2004).

#### 6.4.2.4. Concluding comments on recent acidification

Broad-scale recent acidification of arctic lakes is not apparent through DI-pH reconstructions. This may be partly because most of the lakes studied are not particularly sensitive to acidification or are not located in areas of high deposition. Also, there are no accurately dated high-resolution DI-pH reconstructions from the near vicinity of the local emission sources to allow a paleolimnological evaluation of acidification status in these lakes. However, within a few tens of kilometers from the local emission sources there is clear evidence of the effects of metal accumulation on the lake biota as a function of time, as indicated by the increased proportion of deformed diatoms and chironomids towards the core top. As documented in a recent Arctic-wide study (Smol *et al.*, 2005), climate change is already affecting lakes and ponds and their biota in arctic areas. This is evident from the marked recent change in algal and zoological paleolimnological indicators that is inconsistent with atmospheric acidification or nutrient input but instead indicates the widespread impacts of climatic change (e.g., Douglas *et al.*, 1994; Sorvari *et al.*, 2002; Michelutti *et al.*, 2003).

### 6.5. Interaction between acidification and other environmental issues

Both the causes and effects of air pollutants are closely linked to other environmental problems and human activities. Many of the traditional air pollutants and green-

house gases have common sources. They interact chemically and physically in the atmosphere and cause a variety of interrelated environmental effects at different spatial scales (Swart *et al.*, 2004; Table 6.5). Politically these different air pollutants and greenhouse gases have been treated separately; they also have different spatial and temporal scales. Others have a more local deposition pattern (e.g., many heavy metals) while others are spread over hundreds of kilometers or even globally (sulfur, greenhouse gases). The different spatial scale of each pollutant and its effects makes georeferenced modeling necessary, as this can handle different scales and resolutions simultaneously (van Rompaey, 1995 and references therein). There is also a need to seek synergies in emissions controls for air pollution and climate change to gain economic and political benefits (Swart *et al.*, 2004). Furthermore, the projected climate change-related alterations in temperature, wind patterns, and precipitation can change the routes of contaminant entry and the locations and amounts of deposition in the Arctic (AMAP, 2002; Macdonald *et al.*, 2003; ACIA, 2004).

Table 6.5. Impacts of various substances emitted to the air (redrawn from Seip and Aunan, 2002).

	Regional impacts			Local impacts		
	Climate change	Acidification	Ground level ozone	Health	Vegetation	Materials
CO <sub>2</sub>	x					
CH <sub>4</sub>	x		x			
SO <sub>2</sub>	x	x		x	x	x
NO <sub>x</sub>	x	x	x	x	x	x
NH <sub>3</sub>	x	x		x	x	x
NM VOC <sup>a</sup>	x		x	x	x	?
PM <sup>b</sup>	x	x		x	x	x

<sup>a</sup> non-methane volatile organic compounds; <sup>b</sup> particulate matter

Climate warming, acid deposition, the effects of toxic chemicals, and increasing exposure to ultraviolet (UV) radiation are all regarded as widespread problems in northern ecosystems. In the past, climate warming, acidification, and UV radiation were treated as if they had distinct and separate effects on ecosystems. It is now clear that these major stresses caused by man's alteration of the atmosphere cannot be studied in isolation (Schindler *et al.*, 1996; Schindler and Curtis, 1997). The recent Arctic Climate Impact Assessment (ACIA, 2004) concluded that the total impact of contaminants, excess UV radiation, and climate warming is greater than the sum of its parts. Thresholds of tolerance of biodiversity or of the structural and functional attributes of ecosystems to changes in atmospheric stressors are largely unknown (Freedman and Beauchamp, 1998). Multiple stresses, for example, concomitant changes in the atmosphere and land use, make these thresholds even more difficult to define.

This section summarizes the possible interactions between acidification and other environmental issues in high-latitude aquatic ecosystems in two key sectors: climate change and UV radiation, and heavy metals/contaminants.

## 6.5.1. Interactions concerning climate change and UV radiation

### 6.5.1.1. Anticipated changes in climate

The overarching stress on ecosystems in the future will be global climate change, which is projected to be greatest in the Arctic due to various feedback mechanisms (Chapman and Walsh, 1993; Overpeck *et al.*, 1997). The Arctic is highly likely to be warmer and moister in future. According to the Arctic Climate Impact Assessment (McBean, 2005), the average surface temperature in the Arctic is currently increasing by approximately 0.09 °C per decade and it is probable that there was an increase in arctic precipitation of 1% per decade during the past century. The Arctic Climate Impact Assessment projected (Kattsov and Källén, 2005) that the annual precipitation will increase by 7.5 to 18.1% by 2071-2090, depending on the model used. The projected increase is generally greatest in autumn and winter and smallest in summer. Temperature is projected to increase by 2.8 to 4.6 °C in the Arctic (north of 60° N) by 2071-2090. Climate change will influence water quality by altering the balance between the atmospheric, terrestrial, and aquatic processes in watersheds, and the effects of human resource use on these processes. Among the most important physical and chemical changes projected for freshwater ecosystems are: increasing water temperatures, thawing permafrost, changes in ice cover on rivers and lakes, and increasing levels of contaminants (ACIA, 2004).

### 6.5.1.2. Anticipated changes in hydrology and water quality

#### *Effects of increasing temperature and moisture on water quality*

Murdoch *et al.* (2000) summarized some of the potential water quality effects of increasing temperature and changing moisture conditions. If air temperatures increase, fewer arctic lakes and streams will freeze to the bottom and lakes will have an increased number of ice-free days. In both cases these changes will increase nutrient cycling and productivity. A shallower depth of freezing in arctic lakes during warm years has been shown to cause increased productivity through a lengthening of the growing season. If sufficient nutrient sources are available, lakes in arctic and alpine regions will also experience increasing productivity as a result of more frequent mixing and deeper thermoclines with increases in temperature. Longer thaw seasons will enhance decomposition and greenhouse gas releases from northern wetlands and peatlands. Increased moisture is predicted to decrease permafrost thawing compared to warm-dry conditions, but increased erosion from high runoff may yield greater nutrient, DOC, and sediment loads from thawing permafrost terrain.

Dissolved organic carbon is derived from terrestrial vegetation, especially wetland vegetation. Concentrations in high latitude lakes are low. In small boreal lakes DOC is the most important determinant of thermocline depth (Perez-Fuentetaja *et al.*, 1999). Changes in the levels of DOC are likely to be one of the most important changes of climate warming in surface waters (Schindler, 2001). The effects of climate change on treeline lakes are likely to be strong since shifts in vegetation cover near the treeline will affect DOC in lake waters as well as hydrology (Rouse *et al.*, 1997). The projected increase in supply of organic material and nutrients will enhance primary production



in arctic freshwaters dramatically (Schindler, 1997; Hobbie *et al.*, 1999; Flanagan *et al.*, 2003).

Thawing permafrost increases the flow into ground-water, which can lead to the disappearance of water bodies in some areas but an increase the area of wetlands and small ponds in others. Longer ice-free periods and the northward advance of vascular plants will increase evapo(transpi)ration, leading to lower water levels. On the other hand, increased precipitation and cloudiness may counteract this. However, lakes, ponds, and wetlands are more likely to dry out during summer (Wrona *et al.*, 2005). Earlier ice break-up affects supplies of nutrients, sediments, and water quality, but increases the amount of harmful UV-radiation during spring. These multiple stressors will impact on the aquatic biota at a time when they are having to compete with new species moving north from lower latitudes.

#### *pH changes in high-latitude and high-altitude lakes*

Thawing permafrost and a deepening of the active layer are very likely to increase geochemical weathering and nutrient release (Wrona *et al.*, 2005). Warmer wetter conditions will also favor erosion and greater runoff of weathering products. This could increase the buffering capacity of lakes against strong acids.

As previously described, climate change probably alters the chemical properties of high-latitude lakes, potentially influencing their susceptibility to anthropogenic atmospheric deposition. A small change in temperature can have a profound impact on aquatic environments, especially in areas where the duration of snow and ice cover has a dominant role in determining ecosystem functioning. It has been suggested that climate (e.g., temperature and precipitation) indirectly controls the pH of oligotrophic alpine lakes (Psenner and Schmidt, 1992) as well as poorly buffered arctic lakes (Wolfe, 2002). In the Arctic, this is explained by dissolved inorganic carbon (DIC) dynamics, which are affected by lake ice cover and primary productivity (Wolfe, 2002). Thus DIC regulates the pH of dilute lakes. Globally, lakes are supersaturated by carbon dioxide (CO<sub>2</sub>) and under prolonged ice conditions this situation is exaggerated, which leads to a decline in pH. With a warmer climate and longer ice-free season CO<sub>2</sub> supersaturation is eliminated and enhanced algal production further reduces limnetic CO<sub>2</sub>, shifting the balance in inorganic forms of dissolved carbon away from CO<sub>2</sub> and toward HCO<sub>3</sub><sup>-</sup>, resulting in an increase in pH (Wolfe, 2002). If the catchment to lake area ratio is small (e.g., < 10) in-lake processes become especially important in determining the lake pH status. Otherwise changes in the catchment, such as vegetation, land use, erosion, and weathering, would mainly drive changes in lake water quality.

Climate change may also affect the seasonality of the stress. A rise in algal productivity under warmer conditions may lead to larger seasonal fluctuations in pH as organic matter production influences the redox potential and consequently the acid-base equilibrium (Psenner and Schmidt, 1992), thus acting as an additional stress on the biota with pulses of strong acids into lakes in spring (see section 6.3). The relationship between climate and pH is, however, dependent on the unbuffered character of remote high-latitude lakes, which allows the direct control of pH by DIC dynamics. In general, it seems that a warmer and moister future climate would increase the alkalinity of arctic lakes.

At present, water below the winter ice in subarctic lakes is warmer than in lakes further south due to the rapid ice over. This sustains efficient decomposition on the sediment surface and causes anoxia in lakes (especially mesopolyhumic lakes) at the end of winter. Climate warming may remove this problem by shortening the ice-on period and increasing wind mixing in shallow lakes. On the other hand, increased production and DOC concentrations in shallow lakes may strengthen thermal stratification and sedimentation of organic matter to the bottom, thus increasing oxygen demand. Deep lakes are projected to be more strongly stratified. Changing the redox situation near the bottom will affect the alkalinity of the lakes.

#### *Multiple stressors and effects on biota*

The combined effects of multiple stressors may be antagonistic or synergistic, i.e., smaller than expected or larger than expected, respectively (Folt *et al.*, 1999). The cumulative impacts of anthropogenic stressors on ecosystems are especially worrying in relation to a loss of biodiversity and to changes in ecosystem functioning (Sala *et al.*, 2000; Schindler, 2001; Vinebrooke *et al.*, 2004). When the critical load for multiple environmental stresses is exceeded, the ecosystem may change abruptly. Sala *et al.* (2000) projected very little change in biodiversity in arctic areas if the interactions between driving factors (land use, climate, nitrogen deposition, biotic exchange and atmospheric CO<sub>2</sub>) are synergistic, and much change if the interactions are antagonistic.

For example, acidification typically shifts phytoplankton communities towards larger dinoflagellates and filamentous algae, while suppressing cyanobacteria and smaller chrysophytes and larger eukaryotic algae that are less well adapted to higher temperatures, suggesting that acidification and environmental warming have synergistic negative impacts on phytoplankton. It has been hypothesized that since acidification favors smaller zooplankton species, such as rotifers and certain copepods, the effects of several other stressors on zooplankton will be reduced in acidified lakes. For example, smaller zooplankton species experience lower metabolic costs per capita than larger species during warming, and are less vulnerable to UV radiation, visually feeding vertebrate predators, and pesticides. On the other hand, some planktonic crustaceans, which are UV-tolerant due to their ability to produce photo-protective pigmentation, are likely to face an increased risk of predation by visually feeding planktivorous fish, especially if large populations of alien sportfish are introduced into clear lakes (Vinebrooke *et al.*, 2004). In their experiments on multiple stress effects on daphnids, Folt *et al.* (1999) concluded that effects of toxins and low food supply would probably be enhanced by thermal stress. Hypothesized antagonistic and synergistic effects of anthropogenic acidification and other major abiotic and biotic stressors on phytoplankton and zooplankton are shown in Table 6.6.

#### **6.5.1.3. Recovery from acidification in surface waters**

There is good documentation of a large-scale chemical recovery process from surface water acidification in Europe and North America (Stoddard *et al.*, 1999; Forsius *et al.*, 2001; Evans *et al.*, 2001; Skjelkvåle *et al.*, 2001a). Recovery is also documented in the northern regions (section 6.1.2). Modeling studies based on current emission reduction

Table 6.6. Hypothesized antagonistic (+) and synergistic (-) effects of major abiotic and biotic stressors (first column) and anthropogenic acidification on phytoplankton and zooplankton (redrawn from Vinebrooke *et al.*, 2004).

	Phytoplankton	Zooplankton	Rationale
Elevated temperature	-	+	Acid-sensitive cyanobacteria are thermophilic; small acid-tolerant zooplankton (e.g., rotifers) experience lower metabolic costs than large acid-sensitive cladocerans.
Ultraviolet radiation	+	+	Acid-tolerant species must also be UV-tolerant as they experience elevated UV exposure during lake acidification and loss of UV-attenuating DOC. Small acid-tolerant rotifers show higher UV-tolerance than other zooplankton.
Eutrophication	-	?	Acid-sensitive cyanobacteria are better competitors for nutrients than other groups of phytoplankton.
Toxins	+	+	Acid-tolerant, small zooplankton species are more resistant to contaminants than large species.
Predation	+	-	Large, acid-tolerant phytoplankton (e.g., cyanobacteria, dinoflagellates) are less edible; smaller zooplankton are more susceptible to predatory invertebrates like <i>Chaoborus</i> , while pigmented UV- and acid-tolerant zooplankton are easily detected by introduced fish.

plans project further chemical recovery (Jenkins *et al.*, 2003; Wright *et al.*, 2005). Uncertainties in these scenarios are mainly related to the effects of climate change and future behavior of nitrogen in the ecosystem. Other uncertainties are related to the biological response.

Present-day climatic conditions are commonly assumed in model projections of future scenarios. However, as previously discussed large changes in climate are anticipated for the Arctic and the direction and degree of change may significantly affect the dynamics of terrestrial and aquatic ecosystems. Skjelkvåle *et al.* (2003) identified four key climate-related factors that may influence recovery from acidification: (a) increased frequency and severity of sea-salt episodes, (b) increased frequency and severity of drought, (c) increased turnover of organic carbon, and (d) increased mineralization of nitrogen. Although most empirical evidence is from boreal or temperate regions, it is likely that these processes are also relevant for most arctic environments.

#### *Increased frequency and severity of sea-salt episodes*

The 'sea-salt effect' in surface waters is important in areas receiving substantial inputs of sea-salts, in particular coastal areas of Norway, the United Kingdom and the United States. The sea-salt effect may temporarily increase the acidity of runoff. However, a prerequisite for the lake and stream acidification effect to occur from sea-salt episodes is that the catchment soil is acidic. Recent climate forecasts project a dramatic increase in the North Atlantic Oscillation index (see section 3.7.3) over the next 80 years, implying that warm, westerly conditions in winter may become more prevalent. A greater frequency and intensity of sea-salt episodes may therefore be expected in coastal surface waters. Sea-salt episodes may have important regional biological effects. Massive regional fish kills were reported for the first time after the episode in southwestern Norway during winter 1993 (Barlaup and Åtland, 1996). The fish deaths occurred in moderately acidified systems that suddenly became extremely acid.

#### *Increased frequency and severity of drought*

In parts of North America, the reduction and storage of sulfate in wetlands, and its subsequent re-oxidation and release, have been shown to have a major impact on runoff water quality and hence recovery (Dillon and LaZerte, 1992; Jeffries *et al.*, 2003). Immobilization and re-mineralization of sulfur within soil organic matter are both impor-

tant in European soils. Both stores of sulfur are sensitive to drought. Relationships between sulfate pulses and drought associated with El Niño events have been shown to occur in lakes in Ontario, Canada (Dillon *et al.*, 1997). In the UK, large flushes of sulfate were widely observed in streams following a drought in 1995 (e.g., Harriman *et al.*, 2001). Climate-regulated sulfur retention and release represent 'noise' within an overall recovery trend. Release of stored sulfur will delay recovery where pools are large. Also, sulfate flushes following drought (particularly if droughts become more severe with climate change) may continue to generate acidic episodes, despite improvements in baseline water quality. These drought-driven episodes may be more extreme or frequent in future climate scenarios and may, like sea-salt episodes, contribute to a delay in chemical and biological recovery in surface waters.

#### *Increased turnover of organic carbon*

Regional trends of increasing DOC concentrations over the last two decades have been documented across substantial parts of northern and central Europe (Evans and Monteith, 2001; Skjelkvåle *et al.*, 2001a,b, 2005; Evans *et al.*, 2005; Vuorenmaa *et al.*, 2006) and eastern North America (Stoddard *et al.*, 2003). For Canada the picture is less straightforward (Jeffries *et al.*, 2003). The widespread occurrence of these trends indicates a regional cause and various hypotheses have been put forward to explain them. It has been proposed that these increases may be coupled to change in potential drivers, such as increasing temperature (Freeman *et al.*, 2001; Hejzlar *et al.*, 2003), changes in hydrological regimes (Tranvik and Jansson, 2002; Hejzlar *et al.*, 2003), increasing atmospheric CO<sub>2</sub> concentration (Freeman *et al.*, 2004), airborne nitrogen enrichment in soils (Findlay, 2005), or decreasing sulfur deposition (Stoddard *et al.*, 2003; Evans *et al.*, 2005; Vuorenmaa *et al.*, 2006). More research is needed to establish the relative significance of the various drivers in the different regions.

Elevated DOC concentrations in surface waters have raised concerns that terrestrial carbon stores may be becoming unstable, with unpredictable consequences for the global carbon cycle and with complex consequences for surface waters. Recovery from acidification along with decreasing acid deposition is being partially offset by increasing organic acidity. Increasing DOC concentrations may also cause increased buffering of changes in pH, increased water coloration, and decreased visible light and UV-B

penetration within the water column (see sections 6.5.1.2 and 6.5.1.4). While increased organic acidity may delay chemical recovery from acidification in surface waters, the other factors may influence biological recovery.

#### *Increased mineralization of nitrogen*

Additional uncertainties with regard to nitrogen processes relate to the influence of climate (short and long term) on nitrate leaching, which may alter the long-term trend, or simply add 'noise' to the anthropogenic signal. Because internal ecosystem cycling of nitrogen greatly exceeds system inputs and outputs, any disturbance of this cycle has the potential to completely obscure the relationship between nitrogen deposition and runoff. Both in the UK and the US, large pulses of nitrate have been observed in surface waters following severe winters possibly as a result of soil freezing (Mitchell *et al.*, 1996; Monteith *et al.*, 2000). The frequency of such pulses may change in future in response to altered climate. Results from the CLIMEX project, where ambient air and soil temperature was increased over three years, show increased leaching of inorganic nitrogen, probably due to increased mineralization and nitrification rates in soils (Wright and Jenkins, 2001). Continued high deposition of nitrogen is likely to increase nitrogen saturation and generate increased nitrate concentrations in runoff, thereby delaying recovery due to reductions in sulfur emissions. Increased temperature due to climate change may increase nitrate in runoff and thereby contribute to delay in recovery. The role of nitrogen in acidification and recovery remains uncertain.

#### 6.5.1.4. Impacts of DOC changes on UV radiation in lakes

Climate effects on exposure to UV radiation were discussed extensively by Wrona *et al.* (2005). Humic substances strongly absorb UV radiation and act as a screen, thus protecting organisms from its detrimental effects. Under normal conditions all wavelengths of solar radiation in freshwater ecosystems, including UV radiation, are attenuated to some degree by DOC (Schindler, 1999) and long term variation in underwater UV irradiance is primarily controlled by the amount of dissolved organic material. Long lasting ice cover has a similar protective role, protecting organisms from UV radiation during the most intensive radiation period in spring. Interactions between climate

change and UV radiation potentially modify the underwater UV radiation regime via: (a) changes in stratospheric ozone levels, (b) changes in snow- and ice-cover duration, and (c) changes in DOC levels in natural waters (Wrona *et al.*, 2005). A reduction in snow and ice cover on the surface of rivers, lakes, or oceans is likely to increase the exposure of many organisms to UV radiation (Weatherhead *et al.*, 2005). On the other hand, UV penetration in dilute lakes that are presently within catchments dominated by tundra and forest-tundra may be significantly reduced if climate warming induces a northward shift in the treeline or increases in the density of forest cover, leading to increased DOC inputs (Pienitz and Vincent, 2000; Figure 6.15). Lakes with very high catchment to lake area ratios are likely to be most affected by vegetation changes in treeline areas (Pienitz and Vincent, 2000). Export of humic substances is related to climate and to the balance between production and decomposition. However, climatic as well as deposition impacts on DOC export are under debate (Evans *et al.*, 2002; section 6.5.1.3).

Thus, both acid precipitation and climate warming may enhance the exposure of aquatic organisms to increased UV radiation, which is usually attributed only to depletion of stratospheric ozone (Schindler *et al.*, 1996; Gorham, 1998; Schindler, 1999). Levels of DOC decrease rapidly when lakes become acidified. Lower pH values lead to increased protonation of functional groups and cause the dissolved materials to become hydrophobic and to settle out of the water column onto bottom sediments. Increased photo-degradation also seems to occur. Acidification, climate warming, and stratospheric ozone depletion may thus act together in a 'three-pronged attack' of UV radiation on aquatic systems, particularly in dry conditions (Schindler *et al.*, 1996; Schindler, 1999; Figure 6.15).

#### 6.5.2. Interactions concerning heavy metals/contaminants

In principle, the atmospheric transport routes for acidic compounds should be similar to the pathways of other contaminants, such as heavy metals. Trace metals can catalyze the oxidation of SO<sub>2</sub> to sulfate by sunlight (Kellogg, 1995). Acid precipitation contains a variety of trace metals. However, the deposition pattern of heavy metals is different from SO<sub>2</sub> in that they (usually) deposit within




Figure 6.15. Climate warming will induce environmental changes in the Arctic that alter the exposure of aquatic organisms to the increasing levels of UV radiation caused by stratospheric ozone depletion: (a) a warmer wetter climate is likely to result in higher DOC levels in arctic lakes which, because humic substances strongly absorb UV radiation, will offset the effects of increased UV radiation from stratospheric ozone depletion (modified from Schindler, 1999), whereas (b) in arctic areas with a warmer drier climate the effects of stratospheric ozone depletion, acid rain, and climate warming can combine to increase the exposure of aquatic biota to UV radiation by causing a decrease in the DOC levels in lakes (redrawn from Schindler, 1999).



the immediate vicinity of the sources (mercury being a clear exception). Therefore, the interactive effects of metals and acid precipitation are especially visible locally near the emitters. The co-emissions of the smelters (base cations) may be adequate to prevent environmental acidification at the regional scale (e.g., Kashulina *et al.*, 2003). The AMAP heavy metals assessment (AMAP, 2005) included little information on the potential interactions of heavy metals with other pollutants, for example sulfate, or the cumulative impacts of multiple pollutants. This section addresses potential interactions of heavy metal/contaminant pollution with other air pollution issues, including climate change. The complementary AMAP assessment on contaminant pathways (Macdonald *et al.*, 2003) addressed recent global changes and their effects on the distribution of various contaminants.

#### 6.5.2.1. Processes in air

Warming, increased precipitation, and thawing permafrost will increase the transport and deposition of contaminants to the Arctic. This was discussed in detail by Wrona *et al.* (2005) in relation to mercury and persistent organic pollutants. The timing of weather events also affects the transfer of contaminants to the Arctic. For example, snowfall occurring at the time of arctic haze would increase the transfer of contaminants to the ground at that time of year.

The northward movement of toxic metals and organic pollutants by cold condensation was confirmed by evidence from tree bark (Simonich and Hites, 1995) and lake sediments (Muir *et al.*, 1995). Differential distillation, which pollutants undergo, allows them to be released in vapor form from warm areas and to re-condense in colder regions. However, Givellet *et al.* (2004) concluded that there is no evidence for the 'cold condensation hypothesis' and that the Arctic was not an important natural sink for mercury during pre-industrial times. Moreover, while the anthropogenic emissions of mercury are thought to have decreased by 30% in the past 20 years (Pacyna and Pacyna, 2002), there is evidence that mercury fluxes may have doubled over the past 100 years in the Arctic (Lockhart *et al.*, 1995, 1998; Jackson, 1997) and that the average concentration of heavy metals has more than doubled on the Kola Peninsula over the last 20 years (Dauvalter, 2003). According to Givellet *et al.* (2004), springtime mercury depletion events are the chief mechanism for transferring atmospheric mercury to the arctic environment. Climate warming enhances mercury depletion events, which in turn increases the transfer of mercury into the food webs.

Synergistic interactions seem to occur between acidification, climate warming, and stratospheric ozone depletion that enhance the global mercury cycle (reviewed by Schindler, 2001). Global change and air pollutant levels influence the scale of mercury depletion (mercury transport and deposition from the air) (AMAP, 2002). The release of mercury to the atmosphere might be expected to increase as lakes become clearer (due to changes in DOC concentrations), particularly if increased methyl mercury is available due to warming lakes, and as reservoirs are constructed (Schindler, 1999).

#### 6.5.2.2. Processes in terrestrial areas

Terrestrial processes have a key role in determining the impacts of heavy metals on surface waters. On the Kola Peninsula, direct exposure to SO<sub>2</sub> seems to be the main

reason for ecosystem deterioration (Kashulina *et al.*, 2003). The pine ecosystems there act as a biogeochemical barrier against metals by accumulating them in plant tissue and humus horizons (Goryainova and Nikonov, 1997). The function of forest ecosystems as a regulator of the cycles of these heavy metals (copper and nickel) is disrupted if the vegetation is lost (e.g., by acidification) and more toxic substances can then drain into watercourses.

Significant interactions between acidic rain and metal exposure were also observed in the lichen *Bryoria fuscescens* by Tarhanen *et al.* (1999). They established that although the metal load had a very important role in the decline of epiphytic lichen cover, acidic precipitation further disturbs the symbiosis between the photobiont and mycobiont of a lichen. Kashulina *et al.* (2003) concluded that the whole spectrum of emitted elements needs to be studied in the Kola region, in order to understand the effect of anthropogenic activities, including acidification.

#### 6.5.2.3. Processes in surface waters

Changes in the timing of spring freshet, ice melt, and productivity are very likely to alter the efficiency of arctic lakes in capturing contaminants, for example mercury. Episodes of high contaminant levels are likely to occur in freshwaters when stored contaminants are released due to warming climate (Wrona *et al.*, 2005). A wetter climate will expand the spatial extent of direct runoff to surface waters thus significantly increasing pollutant loads from point and non-point sources that are hydrologically isolated or filtered through groundwater aquifers under current flow conditions (Murdoch *et al.*, 2000). Inundation of wetlands, riparian zones, and low-lying soils results in increased mobilization of trace metals and organic compounds from soils, increased mobilization and methylation of mercury, and greater anaerobic activity in saturated soils (sulfate reduction, denitrification) (Murdoch *et al.*, 2000).

A change in flow patterns can alter the capability of a lake to receive and restore contaminants. At present, most of the contaminant load is carried through the lake surface layers under the ice before lake turnover and peak primary production and little is retained in the lake itself (Macdonald *et al.*, 2000). In a warming climate the main runoff pulse may couple with lake mixing and primary production and hence cause more contaminants to be captured within the lake. In the case of acidifying compounds, lakes in catchments with thin soils, especially on slowly weathering bedrock, may thus face strong effects as the substances remain in the lake, rather than being carried rapidly through the lake.

Acidification and heavy metal contamination often work synergistically because the solubility of heavy metals in water increases as pH falls. As a result, heavy metals leach more quickly from contaminated soils in contact with acidic water. Dauvalter (1995) studied 15 lakes in Finnish Lapland. In four acidic lakes, the low pH had changed the geochemical cycling of potentially harmful metals by increasing desorption of metals from sediments back to water and/or decreasing the adsorption of metals onto settling sediment particles. Cadmium and nickel mobilization from sediments during acidification was observed by Nuorteva *et al.* (1987) and Fjeld *et al.* (1994).

Heavy metals are absorbed by plankton at the base of the food web and biomagnified to significant amounts at higher trophic levels. The influence of acidification and eutrophication on metal behavior must therefore be consid-

ered. It is well known that inorganic monomeric aluminum acts synergistically with pH to cause embryo mortality (e.g., Clark and Hall, 1985; Clark and LaZerte, 1985; Freda and McDonald, 1990; Blaustein *et al.*, 2003). UV radiation also may affect the toxicity of contaminants (Wrona *et al.*, 2005).

Water chemistry, especially acidity, and food web structure also affect mercury availability and uptake. Acidification can greatly enhance methylation, producing a higher proportion of bioavailable methylmercury (Miskimmin *et al.*, 1992; Gilmour and Henry, 1991). Lake acidification is known to favor higher methylation to demethylation ratios, both directly and via effects on DOC (see references in Schindler *et al.*, 1995). Climate warming alters the thermal conditions of lakes in a way that increases the methylation of mercury (Ramlal *et al.*, 1993; Schindler *et al.*, 1995). On the other hand, increased penetration of shortwave solar radiation due to decreases in DOC may cause increased conversion of methylmercury to elemental mercury, which is then released back to the atmosphere (Sellers *et al.*, 1996). Once back in the atmosphere, mercury is susceptible to long-range transport and biomagnification in distant food chains (Schindler, 1999).

Future climate change, in interaction with other environmental problems, is thus thought to influence the distribution patterns and mobility of organic pollutants and toxic metals in freshwater systems and to lead to changes in the uptake and accumulation of these substances in

freshwater food chains. This is one of the main hypotheses tested in an ongoing EU project EURO-LIMPACS (<http://www.eurolimpacs.ucl.ac.uk/index.php>).

### **Acidification of the oceans**

Owing to increasing atmospheric CO<sub>2</sub> levels, CO<sub>2</sub> levels in the oceans have increased since pre-industrial times. Added CO<sub>2</sub> decreases the CO<sub>3</sub><sup>2-</sup> (carbonate) ion concentration and makes the ocean more acidic. Ocean pH has decreased by 0.1 units, equivalent to a 30% increase in hydrogen ion concentration over the last two centuries (Caldeira and Wickett, 2003; The Royal Society, 2005). If global emissions of CO<sub>2</sub> continue to rise on current trends, then the average pH of the oceans could fall by 0.5 pH units by 2100 (a 0.5 unit decrease in ocean acidity means a three-fold increase in hydrogen ions). This could have substantial effects on the biological processes in surface oceans. Recent studies indicate that future undersaturation of aragonite and calcite, especially in polar oceans, may have potential biological impacts on calcifying organisms, especially shelled pteropods, the densities of which are high in cold water regions (Orr *et al.*, 2005). Shell dissolution rates of pteropods increase in waters that have become undersaturated with aragonite (e.g., Byrne *et al.*, 1984). Also, benthic calcareous organisms such as cold-water corals are in danger of becoming surrounded by water masses that are undersaturated with aragonite. Orr *et al.* (2005) suggested that some high latitude surface waters will probably become undersaturated within the next 50 years, leading to detrimental conditions for many organisms.

## Chapter 7

# Air Pollution and Health Impacts in the Arctic

Jon Øyvind Odland

In addition to the detrimental ecosystem effects of acidifying compounds and haze precursors described in previous chapters, elevated concentrations of air pollutants may also cause serious human health impacts. Concern over health effects is currently one of the most important factors driving international air pollutant emission reduction policies (Sliggers and Kakebeeke, 2004; UNECE, 2004b).

### 7.1. Major air pollutants of health concern

The sources, health effects, and vulnerable groups for major air pollutants are summarized in Table 7.1. It is very difficult to isolate the health effects of individual pollutants. It is more useful to consider each of the major pollutants as 'indicators' of the mixture of air pollution created by motor vehicles, home heating, and industry. Continuous monitoring of particulate matter, nitrogen oxides, ozone, and carbon monoxide is now established in the big cities (Simpson *et al.*, 2001), but these data provide only general estimates of actual exposures in individuals. Carbon monoxide is the only major air pollutant for which a biomarker of exposure (carboxyhaemoglobin in erythrocytes) is available.

Researchers are currently investigating the importance of the size and chemical composition of particles as a causal factor for cardiorespiratory effects (Brunekreef and Holgate, 2002). The focus is now on the very small particles; PM<sub>2.5</sub> and PM<sub>1</sub> (i.e., particles smaller than 2.5 µm or 1 µm, respectively). As very small particles penetrate further into lungs than larger particles, they are believed to be more strongly associated with adverse health effects. Diesel engine emissions contribute disproportionately to the very-small-particle fraction of urban air pollution (WHO, 1996).

The recent documentation of lung cancer as an effect of long-term exposure to urban air pollution (Pope *et al.*, 2002) highlights carcinogenic chemicals in the smallest air particles and in carcinogenic gases (e.g., benzene; benz[a]pyrene) as possible causal agents.

Carbon dioxide (CO<sub>2</sub>), another air pollutant created by fuel combustion, has no direct health effects at the concentrations occurring in the ambient environment. However, it is the main 'greenhouse gas' causing global climate change (McMichael *et al.*, 1996) and, as such, indirectly contributes to the global health impact of such change. Efforts to reduce urban air pollution by reducing the use of cars would have the added benefit of reducing CO<sub>2</sub> emissions.

Health effects of the major air pollutants are listed in Tables 7.1 and 7.2. In any particular study, establishing

Table 7.1. Selected outdoor air pollutants and their effects on health (adapted from WHO, 2001).

Source	Known health effects	Contributing or potentiating factors	Vulnerable populations
<b>Particulate matter</b>			
Biomass and fossil fuel combustion in home heating, industry and motor vehicle engines; cigarette smoke	Upper respiratory tract irritation and infection; exacerbation of and increased mortality from cardiorespiratory diseases	Sulfur dioxide, sulfuric acid; cold, heat, humidity	Elderly people with respiratory and cardiovascular diseases; children with asthma
<b>Sulfur dioxide and acid aerosols</b>			
Fossil fuel combustion; metal smelting and petrochemical industries; home heating/cooking with coal	Throat irritation; exacerbation of cardiorespiratory diseases, including asthma	Exercise, particulates, asthma	People with respiratory diseases (e.g., children with asthma); elderly people with respiratory and cardiovascular diseases
<b>Oxides of nitrogen</b>			
Fuel combustion at high temperature (e.g., from vehicle engines, gas cooking and heating)	Eye irritation; upper respiratory tract infection (especially in children); exacerbation of asthma; irritation of bronchi	Exercise, respiratory tract infection, asthma	People with respiratory diseases (e.g., children with asthma)
<b>Ozone</b>			
Reaction product of sunlight and vehicle pollutants; hydrocarbons and oxides of nitrogen	Eye and throat irritation; reduced exercise capacity; exacerbation of respiratory disease	Exercise, respiratory tract infection, asthma	People with respiratory diseases (e.g., children with asthma)
<b>Carbon monoxide</b>			
Biomass and fossil fuel combustion; cigarette smoke and vehicle exhaust	Headache, nausea, dizziness, breathlessness, fatigue, visual disturbance, confusion; angina, coma, death; low birth weight (after maternal exposure during pregnancy)	Coronary artery disease	People with ischemic heart disease
<b>Lead</b>			
Smelting; leaded petrol	In children: neuropsychological and cognitive effects. In adults: hypertension, classic lead poisoning	Other sources of lead; iron deficiency	Children, pregnant women
<b>Other pollutants; 'air toxics'</b>			
(hydrocarbons, aldehydes, other organic compounds, asbestos)	Eye irritation; lung cancer; asthma	Smoking, occupational exposures	Smokers, asbestos workers, people with asthma, children

Table 7.2. Adverse respiratory health effects of air pollution (American Thoracic Society, 2000).

- 
- Increased mortality
  - Increased incidence of lung cancer
  - Increased frequency of symptomatic asthma attacks
  - Increased incidence of lower respiratory tract infections
  - Increased exacerbation of chronic cardiopulmonary or other diseases, reflected in various ways, including reduced ability to cope with daily activities, increased hospitalization, increased physician visits and medication, and decreased pulmonary function
  - Reduction of FEV1 (forced expiratory volume in one second) or FVC (forced vital capacity) associated with clinical symptoms
  - Increased prevalence of wheezing unrelated to colds, or wheezing on most days or nights
  - Increased prevalence or incidence of chest tightness
  - Increased prevalence or incidence of cough/phlegm production requiring medical attention
  - Increased incidence of acute upper respiratory tract infections that interfere with normal activity
  - Acute upper respiratory tract infections that do not interfere with normal activity
  - Eye, nose, and throat irritation that may interfere with normal activities (e.g., driving a car), if severe
- 

whether there is an association between air pollution and one or more of the effects listed depends on exposure level, the background health status of the population exposed, and their age.

## 7.2. Key epidemiological findings

Most recent epidemiological studies of air pollution and mortality have used time-series analysis to relate daily mortality rates to daily air pollution levels (on the same day or previous days). However, this approach cannot be used to ascertain whether increased mortality reflects a significant reduction in life expectancy (Brunekreef and Holgate, 2002). A few studies have documented associations between mortality and air pollution exposure over longer periods. The Harvard 'six cities' study involved a 14–16-year prospective cohort of more than 8000 adults in the United States, and the American Cancer Society (ACS) study collected data on over 500 000 people living in 51 different US metropolitan areas between 1982 and 2000 (Pope

*et al.*, 1995, 2002). These studies reported significant associations between annual average particle pollution levels (PM10 or PM2.5) and annual all-cause mortality rates: an average increase of 10 µg/m<sup>3</sup> of PM10 or PM2.5 was associated with a 3–4% increase in mortality. The ACS study also found a significant association between levels of PM2.5 and deaths due to cardiorespiratory diseases and lung cancer (Pope *et al.*, 2002). A 'natural intervention' study using long-term exposure and effect data analyzed mortality in Dublin before and after coal-burning was banned in 1990 (Clancy *et al.*, 2002). The annual average particle pollution level declined by 36% after the ban, while adjusted mortality rates decreased by 15.5% and 10.3% for respiratory and cardiovascular deaths, respectively.

A review of epidemiological studies concluded that there may be a relationship between acute ozone exposure and increased mortality, especially in elderly people (Thurston and Ito, 2001). However, the concurrence of high ozone levels with hot weather makes it difficult to separate the effect of heat from the effect of ozone on mortality.

Acute time-series studies have shown associations between particle pollution and daily hospital admissions, mainly for respiratory diseases (especially asthma and chronic obstructive pulmonary disease) but also for cardiovascular diseases (Morgan *et al.*, 1998; McGowan *et al.*, 2000; Petroeschovsky *et al.*, 2001; Denison *et al.*, 2001; Department of Environment, 2003).

## 7.3. The arctic perspective

The arctic perspective is, in general, the same as the global perspective but with the addition of point sources related to heavy industrial complexes, especially in the area of the former Soviet Union (Smith-Sivertsen *et al.*, 1998). The associations between air pollution and human health in different arctic areas are briefly described in the recent AMAP assessment on human health (AMAP, 2003), and in more detail in many publications related to the Russian–Norwegian Health Group's reports (Smith-Sivertsen *et al.*, 1998, 2001, 2003; Dotterud *et al.*, 2001). The results reported were mostly very optimistic. Studies to date have been unable to show any significant health effects that are directly associated with emissions from the nickel refineries. No

Table 7.3. Reflections and predictions (after Kjellström *et al.*, 2002).

### Circa 1900

Cities being rapidly industrialized. Large amounts of sulfur dioxide and particulates being emitted in heavily populated areas from inefficient combustion of coal in power stations and industrial and domestic furnaces.

Air pollution and resultant lack of sunshine in industrialized areas causing widespread lung damage, high mortality, and an upsurge in diseases such as rickets in children. Indoor air pollution, from cooking with coal and wood, even worse for health than outdoor pollution.

### Circa 2100 (a pessimistic view)

Increasing air pollution in developing countries.

Limited progress on motor vehicle emissions in developed countries.

### Circa 2100 (an optimistic view)

Intelligent global stewardship of our natural resources has led to a major shift towards alternative energy sources such as wind power, solar energy, and fuel-cell engines for vehicles.

More wood burning to heat houses, as this creates less greenhouse gas emissions than fossil fuel.

Coal burning in super efficient and clean-burning electric power stations is still continuing in countries with large coal reserves (e.g., China, India, United States of America).

Many governments have put pressure on motor vehicle manufacturers to produce less-polluting cars.

High-quality public transport systems and advanced telecommunications systems have made daily commuting in private cars largely obsolete.

Private vehicles are mainly used for leisure activities.

---



reduction in lung function between exposed and non-exposed groups could be demonstrated (Smith-Sivertsen *et al.*, 2001). Sensitization described by serum IgE-levels was more common in Russia than Norway, but the Russians did not report any more atopic diseases (Smith-Sivertsen *et al.*, 2003). Atopic diseases seem to be even less prevalent in the Russian study groups than in studies of other northern European countries (Dotterud *et al.*, 2001). Urinary nickel excretion was considerably higher in the Russian populations, but only partly associated with vicinity to the nickel smelters or to occupational exposure (Smith-Sivertsen *et al.*, 1998). Differences in health status of the Norwegian and Russian border populations seem to be more associated with socio-economic conditions than to environmental pollution (Odland *et al.*, 2004).

#### 7.4. The shifting panorama

From a global and arctic perspective, many countries are at the same stage of industrial and urban development as Western European countries 50 to 80 years ago, when high levels of ambient air pollution from coal-burning were common. At the household level, promoting energy-efficient and less-polluting cooking stoves constructed from local materials would be an important step in reducing air pollution (Table 7.3). Switching the energy source for cooking to less-polluting kerosene, gas, and electricity is another solution, often out of reach for poor communities in the short term (Kjellström *et al.*, 2002). Worldwide, a major change in priorities is needed to steer economic development towards low-pollution policies.

## Chapter 8

# Conclusions and Recommendations

*Martin Forsius, John Derome, Lars R. Hole, Sirkku Manninen, Marjut Nyman, Patricia Quinn, Brit Lisa Skjelkvåle, and John L. Stoddard*

### 8.1. Sources of acidifying pollutants and arctic haze precursors

AMAP assessments have clearly documented that air pollutants, including persistent organic pollutants, heavy metals such as mercury, and acidifying substances can reach the Arctic as a result of long-range transport from source regions in Europe, North America, and Asia. In relation to acidifying substances, the industrial areas of northern Europe and parts of Russia and the northeastern United States are responsible for most of the pollution exported to the Arctic. There are also significant sources of acidifying substances within the Arctic, including industrial sources (the metallurgical industry, power plants, oil and gas related activities) and diffuse sources associated with, for example, shipping. Emissions from natural sources within the Arctic (volcanoes, marine algae, and forest fires) are very difficult to quantify and almost impossible to project. However, the frequency, severity, and duration of boreal forest fires do appear to be increasing, possibly related to the influence of changing climate.

Nonferrous metal production remains the dominant source of emissions of acidifying gases to the atmosphere within the Arctic. Non-ferrous metal smelters located at Norilsk, and at Nikel, Zapolyarnyy, and Monchegorsk on the Kola Peninsula (all operated by MMC Norilsk Nickel) together account for 68% of the total 'harmful emissions' (including sulfur dioxide, dust, and nitrogen oxides) from non-ferrous metal production in Russia. These smelters are all located within or close to the Arctic. Sulfur dioxide emissions from these smelters have reduced substantially due to changes in production and better technology for controlling emissions. The main reductions have occurred since 1995 and have been considerably greater on the Kola Peninsula than at Norilsk.

*It is recommended that improved information be obtained on emissions from arctic point sources, in particular for the non-ferrous metal smelters on the Kola Peninsula and at Norilsk.*

Recent modeling results suggest that southeast Asia is not likely to be a major source of acidification-related atmospheric pollution at ground level in the Arctic. Efforts are, however, needed to confirm this finding. Future changes in atmospheric transport patterns due to climate change could also alter this conclusion. Europe (Northern Russia in particular) and to a lesser degree North America continue to be the main source regions for acidifying pollutants carried into the Arctic by long-range transport.

Shipping activities in arctic waters may increase substantially in the future. Projected reductions in sea-ice thickness and extent due to climate change have raised the possibility that shipping activities in arctic waters may increase substantially over the coming decades (as the navigation season lengthens and new sea routes open). Emissions associated with this increased activity are difficult to project.

*The impacts of acidification from arctic shipping should be addressed, and future scenarios for emissions associated with this source should be included in this work.*

The Arctic is thought to contain at least a quarter of the world's undiscovered petroleum resources: most of these in Russia, Alaska, Canada, Greenland, and Norway (off-shore). A significant increase in arctic oil and gas activity can be expected, including activity in offshore areas.

*The impacts associated with oil and gas related activities on acidification and arctic haze in the Arctic should be assessed together with the assessment of the impacts of oil and gas development.*

### 8.2. Trends in air concentrations and deposition

Ice core drillings on Svalbard show that significant changes in atmospheric pollution within the Arctic have only occurred since the beginning of the industrial era, indicated by increased levels of non-sea-salt sulfate, nitrate, acidity, fly ash, and organic contaminants. Acidifying pollutants in rain, snow, dust, and gases have been monitored regularly at purpose-built stations in some parts of the Arctic, mostly since the 1980s. Fennoscandia has several background monitoring stations, while the vast Siberian region and the Canadian Arctic and Alaska have relatively few. Some of the stations have now generated time series datasets that are long enough to show whether concentrations are increasing, decreasing, or staying the same over time.

#### 8.2.1. Air and precipitation

Sulfate concentrations measured at those air monitoring stations with the best air quality time series in the High Arctic (Alert, Canada; and Ny-Ålesund, Svalbard) and at several monitoring stations in subarctic areas of Europe (Oulanka, Finland; Tustervann, Svanvik, Karasjok/Jergul, Norway; Bredkälen, Sweden) show decreasing trends since the 1990s. Although there are relatively few background stations within the Arctic measuring sulfate concentrations, most record a decrease in concentrations since the 1990s. These observations are supported by modeling results. There are few signs of significant trends in precipitation for the period studied. However, expected future occurrence of rain events in both summer and winter will result in increasing wet deposition in the Arctic.

For nitrate and ammonia the pattern is unclear, with increases at some stations and decreases at others. The increasing trends in nitrate are particularly apparent in recent years indicating a decoupling between the trends in sulfur and nitrogen. Time series of sulfur and nitrogen concentrations in precipitation at Norilsk since 1990 do not show any significant trends.

In general, sulfur deposited originates from local point sources and long-range transport. In the AMAP region, high levels of deposition only occur close to large point sources in the vicinity of the Nikel and Monchegorsk smelters on the Kola Peninsula and in Norilsk in north-western Siberia. With the exception of areas affected by emissions from the industrial centers, average atmospheric deposition loads are much lower in arctic Russia than in more southerly parts of the country. Background sulfate levels in precipitation decrease from west to east across the Russian Arctic. There is a similar pattern for background levels of nitrate in precipitation across the Russian Arctic. A trans-Arctic snow study found no overall relationship between pH and levels of sulfate from human activities.

Short-term episodes of winds carrying very high concentrations of air pollutants can result in pulses of pollution entering the arctic environment. Studies at Finnish background stations showed that the five worst days of the year can bring 20 to 30% of the annual sulfate load of bulk deposition.

### 8.2.2. Model projections

The decreasing trends in levels of acidifying pollutants observed at many sites throughout the Arctic are supported by model results. Models indicate that mean concentrations of sulfur oxides and total sulfur deposition within the Arctic almost halved between 1990 and 2000. These decreasing trends are supported by empirical data from the several monitoring stations around the Arctic (see above). The modeled results for airborne oxidized nitrogen are similar to those for sulfur. The models also confirm earlier findings that emissions in Eurasia continue to make the greatest contribution to acid deposition within the Arctic.

Model projections based on future emissions scenarios indicate that the decreasing trends observed between 1990 and 2000 are likely to level off and that only small reductions in concentrations and deposition can be expected after 2020, even if maximum feasible reductions in emissions are achieved. Although further recovery and continuing improvement in the acidification status of the Arctic can be expected during the period until 2020, this is dependent on the implementation of existing international agreements to reduce emissions of acidifying substances. The Gothenburg Protocol is the most important agreement in this connection.

*All arctic countries should be encouraged to ratify the UN ECE LRTAP protocol to Abate Acidification, Eutrophication, and Ground-level Ozone (the 'Gothenburg Protocol') and to support its implementation.*

### 8.3. Arctic haze

Based on measurements of sulfate aerosol and light scattering by aerosols, the amount of arctic haze reaching the Arctic was either relatively constant or decreasing between the 1980s and early 1990s. Levels of sulfate aerosol have decreased since the late 1990s and the indications are that levels are still decreasing. In contrast, nitrate aerosol

appears to be increasing in Alert (Canada), with an unclear trend at Barrow (Alaska). However, trend analysis on light scattering data collected since the late 1990s indicates an increase in the amount of haze reaching the Alaskan Arctic (Barrow). There is also evidence of increasing light absorption, due to black carbon aerosols at Barrow (Alaska) and Alert (Canada). Forest fires are a possible source of this black carbon. The frequency, severity and duration of boreal forest fires seem to be increasing and the pollution plumes from these summer fires can extend over vast areas. Boreal forest fires may even dominate the black carbon budget in the Arctic in years of strong burning.

## 8.4. Effects

### 8.4.1. Human health

Epidemiological studies indicate that differences in health status of the Norwegian and Russian border populations are more associated with socio-economic conditions than environmental pollution. Studies to date have been unable to show any significant health effects that are directly associated with emissions from the nickel refineries.

### 8.4.2. Terrestrial ecosystems

The present deposition of acidifying compounds from long-range, transboundary transport of anthropogenic emissions at lower latitudes does not appear to be a threat to terrestrial ecosystems in most of the Arctic.

The non-ferrous metal smelters are responsible for the vast majority of acidification-related effects in plants and soils within the European Arctic. Information on the effects of emissions from the smelters on the Kola Peninsula on terrestrial ecosystems is accumulating rapidly and is more accessible to the international scientific community than information on the situation in the Norilsk region, which is less detailed and published mostly in Russian.

Acidified soils on the Kola Peninsula mostly occur immediately around the smelters and coincide with the areas where the vegetation has been completely destroyed. Strongly acidic precipitation only falls within about 30 km of the smelters; outside this zone lower sulfur dioxide levels and the presence of alkaline particles in the atmosphere prevent the precipitation becoming acidic. Sparse data on soil microorganisms show that adverse effects are also concentrated in restricted areas around the smelters but are more associated with excessive heavy metal deposition than soil acidification.

Outside the area immediately around the smelters, there is no clear evidence of soil acidification due to sulfur dioxide emissions (and subsequent deposition of acidifying compounds) on the Kola Peninsula. The lack of widespread soil acidification despite high sulfur dioxide emissions appears to result from the simultaneous emission of alkaline fly ash from the power plants and the apatite fertilizer complex, the low interception of acidifying compounds by the sparse cover of coniferous trees, and the low rate of conversion of sulfur dioxide to sulfuric acid in the Arctic. Soils in the Norilsk area are not considered acid sensitive.

In the European Arctic there are direct effects of sulfur dioxide on trees, dwarf shrubs, and epiphytic lichens. Effects on vegetation, some of which are evident at a regional level, are due to changes in air quality and in most cases are not related to changes in soil acidification. Direct effects of sulfur dioxide include visible leaf damage, decrease in needle life span in conifers, and elevated sulfur concentrations in plant tissues. The observed changes in plant community structure (species composition and coverage) also indicate the indirect effects of sulfur and nitrogen deposition. In terms of their effects on plants, it is difficult to differentiate between the effects of acidifying air pollutants and elevated heavy metal levels in soils. Large quantities of heavy metals are emitted by the non-ferrous metal smelters.

The most likely explanation for the changes observed in vertebrate populations near Monchegorsk is habitat deterioration. Winter mortality may be especially high in the contaminated habitats. Changes in insect populations are mostly due to changes in host plant abundance and physiology, accompanied by a decline in some natural predators, and to changes in habitat structure.

The critical loads of acidity for terrestrial ecosystems have not been exceeded in Canada north of 60° N. In the European Arctic the widespread exceedance of critical loads of acidity in 1990 should have almost disappeared by 2010 if the agreed emissions reduction policies are fully implemented. However, the critical levels of sulfur dioxide and critical loads for acidity in highly sensitive forest ecosystems are expected to be exceeded locally and regionally near the non-ferrous metal smelters. (In this context, 'critical level' refers to the atmospheric concentration above which direct adverse effects will occur on plants, while 'critical load' refers to the maximum amount that can be deposited on an ecosystem without causing that ecosystem to become damaged). It must be remembered however, that non-exceedance of critical loads/levels does not necessarily mean that there will be no environmental change.

#### 8.4.3. Freshwater ecosystems

Anthropogenic acidification of lakes is only of concern in areas with both sensitive geology and elevated levels of acid deposition. Because the areas with the highest levels of acid deposition, for example Norilsk, are less geologically sensitive the biggest impacts on lake chemistry and biology have mostly been observed in small sensitive ecosystems in local areas.

Available lake chemistry data provide irregular and incomplete coverage of the Arctic. Only a few of the lakes monitored in the Canadian Arctic are acid sensitive; these are mostly on Baffin Island or the central mainland. There are no lake chemistry data for large areas of the North American Arctic. Acid sensitive lakes are scattered throughout the European Arctic but are most abundant in northern Fennoscandia and the Kola region. Acidification by natural organic acids is important in some lakes.

Chemical monitoring data show that lakes in the Euro-Arctic Barents region are showing clear signs of a regional-scale recovery from acidification caused by long-range

transported sulfur. Lakes close to the pollution sources on the Kola Peninsula are showing the clearest signs of recovery. There are insufficient data to make similar conclusions about regional-scale biological recovery.

Assessments of biological effects of acidification in arctic surface waters are largely based on sparse and isolated data. There are far more chemical data available than biological data. The best biological effects data are from the impacted areas in northeastern Norway and Finland – outside this area it is impossible to draw robust conclusions about biological effects in surface waters. It is important to note that lakes subjected to acidic inputs, even those with measured decreases in acid neutralizing capacity, alkalinity, and pH, have not necessarily been acidified to the point where measurable damage to the biota can be observed.

Biological recovery is occurring in at least two isolated areas. Both sites are directly downwind of point sources on the Kola Peninsula. A comparison of minnow populations in lakes and streams in the Vätsäri area of northeastern Finland between 1993 and 2000 and increasing abundance of the acid-sensitive cladoceran *Daphnia longiremis* since 1999 in Lake Dalvatn, an acid-sensitive lake in Finnmark, both indicate improvement in the acidification status of these waters. A clear recovery of the brown trout population of lake Otervatn in Eastern Finnmark has occurred.

Changes in the microscopic fossil record in lake sediments indicate that most arctic lakes have experienced very slow natural acidification over the last 10 000 years. These paleolimnological studies do not support the hypothesis of recent and widespread acidification of arctic lakes due to sulfur pollution from the smelting and mining industries. Many of the lakes studied are outside areas of high acid deposition or are not particularly sensitive to acidification. Within a few tens of kilometers from the local emission sources there is however clear evidence of the effects of metal accumulation on the lake biota as a function of time, as indicated by the increased proportion of deformed specimens of diatoms and chironomids towards the top section of the sediment. The algal and zoological paleolimnological indicators also show that the global warming is affecting the lakes and ponds in the Arctic.

Sulfur deposition has important effects on the severity of acidic episodes in spring. Data on streams of the Kola North showed that in the periods of pH depression during floods (acidic episodes) the total concentration of toxic aluminum in water increased by 50 to 88%. A model based assessment on data from streams in northern Sweden indicated that reduced emissions of acid precursors have generated significant improvements in the surface water chemistry during episodes associated with spring runoff, and that episodic acidification decreased by between 40 and 80% between 1990 and 1999. It is also likely that expected future reductions in acid deposition will further improve the spring flood acidification situation in the northern regions.

In 1990, critical loads of acidity for surface waters in the European Arctic were exceeded almost everywhere. Implementation of agreed emission reduction policies will reduce the area in which critical loads are exceeded and, to a greater extent, the amount by which the critical loads



are exceeded. But even with the implementation of the maximum feasible emissions reductions (the 'MFR 2020' scenario) there are still very likely to be areas where the critical loads for surface waters are exceeded. The required deposition reduction is up to 3.2 kilograms of sulfur per hectare per year in the most exposed area of the Kola Peninsula.

### 8.5. Links between acidification, arctic haze, and other environmental issues

The effects of haze aerosols on the arctic climate are complicated by feedbacks between aerosols, clouds, radiation, sea ice, and vertical and horizontal transport processes. Whether the pollutant aerosols cause an overall warming or an overall cooling is not yet known. Changes in the light scattering and absorbing properties of the haze directly affect the amount of solar radiation passing through the haze.

The causes and the effects of air pollutants are closely linked to other environmental problems. For example, climate change, changes in exposure to ultraviolet radiation, the effects of toxic chemicals, and land-use. Complex interactions between these factors can strongly affect ecosystem responses to air pollutants.

Recent studies indicate that if global emissions of carbon dioxide continue to rise at current rates, then the average pH of the surface oceans could fall by 0.5 pH units by 2100. This could cause detrimental conditions for calcifying organisms, especially in high latitude surface oceans, within the next 50 years.

*Future AMAP assessments should view acidification and arctic haze in a wider context. Particularly in relation to toxic pollutants, particulate matter, and climate change and climate feedback effects (i.e., reduced snow albedo as a result of increasing black carbon deposition).*

*Chemical and biological variables should be monitored and studies conducted to assess the risk of acidification of high latitude surface oceans due to increased carbon dioxide emissions.*

### 8.6. Gaps in knowledge and recommendations concerning monitoring and research needs

#### 8.6.1. Geographical gaps

New information from the areas around Norilsk has provided valuable insight into the sensitivity, and potential vulnerability of this region. The few samples taken, however, are not sufficient to enable an overall assessment of the geographic extent or trends in environmental effects in the Norilsk region.

*Continued effort should be made to collect data and information for the Norilsk region. This should involve both MMC Norilsk Nickel and the relevant local and regional authorities.*

While Fennoscandia has several background air monitoring stations for acidification parameters, most areas of the Arctic have few, if any, background air monitor-

ing stations. The vast regions of Siberia east of the Urals, Alaska, and the Canadian Arctic are covered by relatively few stations, and some of these are being closed. A better distribution of background monitoring/research stations will enable a more complete and accurate circumpolar assessment to be made of acidification and arctic haze, as well as of other air pollution issues.

*Additional background monitoring stations for air and precipitation chemistry should be established in northern Canada and Alaska. More regional and background air monitoring and precipitation chemistry stations should be established in the Russian Arctic. These background air monitoring stations should be extended to form 'multi-purpose' monitoring stations that monitor acidification parameters together with, for example, persistent organic pollutants and heavy metals, ozone and UV, and precipitation chemistry and relevant meteorological and ecosystem effects parameters. This will help to ensure cost-effective monitoring at remote sites that are expensive to operate and maintain, and will allow data to be assessed in a more multidisciplinary context.*

#### 8.6.2. Data availability

Both AMAP and EMEP have benefited from a close cooperation with respect to monitoring levels of acidifying pollutants in air and precipitation within the European Arctic. The new EANET (Acid Deposition Monitoring Network in East Asia) initiative represents an opportunity to develop similar cooperation in relation to monitoring in the Far East of Asia, including assessment of the acidification potential and arctic haze impacts of long-range transported air pollutants from southeast Asia. Programs addressing acidification issues also exist in North America, although their coverage of arctic areas is limited.

*AMAP should continue to develop a better linkage with programs such as EMEP and EANET and with appropriate national experts (particularly in Russia and North America) to extend the geographical areas and time series datasets available for use in AMAP assessments.*

*Improved efforts should be made to ensure that relevant data, both existing and future, from North American monitoring stations are reported to the AMAP database at NILU according to agreed procedures. Including more North American data in the AMAP database would enable a better understanding of hemispheric pollutant transport to the Arctic and would be a useful source of data for groups engaged in hemispheric transport modeling.*

*Monthly data from Russian precipitation chemistry monitoring sites should be reported to the AMAP database at NILU according to agreed procedures. This will enable a better understanding of acid deposition at the regional level within the European Arctic.*

#### 8.6.3. Trends in air and precipitation

Long-term monitoring is extremely important, even when trends are decreasing. For example, background monitoring stations that measure sulfate have mostly recorded a decrease in concentrations since the 1990s. In contrast, many stations have recorded an increasing trend in nitrate

over recent years. This indicates a decoupling between the trends in sulfur and nitrogen. There is also evidence of increasing light absorption due to black carbon aerosols at Barrow (Alaska) and Alert (Canada). Natural climate fluctuations (such as the North Atlantic Oscillation) affect the concentrations of acidifying air pollutants and levels of arctic haze. Time series datasets need to be long enough to resolve the effects of natural climate fluctuations and to identify the underlying trends. Measurements linked to chemical transport models are particularly important for helping to establish the causes of trends.

*Monitoring of trends in acidic species in the Arctic and other high latitude stations should continue and sulfur dioxide should be added to the list of chemicals monitored at Barrow (Alaska).*

*Monitoring of aerosol composition, light scattering, and light absorption should continue at Barrow (Alaska) and Alert (Canada), as well as at other locations to better define changes in arctic haze over time.*

#### 8.6.4. Pathways

Recent modeling studies yield conflicting results as to whether southern Asia is a source of pollutants to the Arctic. Rapid industrialization in this region has caused an increase in the emissions of many air pollutants and these increases are likely to continue. The transport pathways to the Arctic are likely to be affected by the projected changes in the climate system.

*Co-operation with relevant international organizations, such as EMEP, the LRTAP Convention Task Force on Hemispheric Transport of Air Pollutants, and EANET should be enhanced to obtain more precise data on emissions from southeast Asia and to investigate the possible impact of these emissions on the Arctic.*

*Further research is needed into pollution transport pathways to the Arctic. In particular, the potential effects of future changes in transport pathways due to climate change should be assessed.*

#### 8.6.5. Effects monitoring and research

The work on ecosystem effects has clearly shown that it is very difficult to differentiate the effects caused by different pollutants, for example effects caused by acidification, poor air quality, and toxic metals. Data on effects on biota are sparse and isolated and key chemical data for cause-effect assessments are often incomplete. Episodic effects are very important in the Arctic due to the extreme climate and the large emission point sources. Surface waters are generally more sensitive to inputs of acidifying pollutants than soils. For this reason there is a greater and more extensive exceedance of critical loads to lakes than soils. Recovery will initially be observed as decreases in acid-

ity followed by recovery of aquatic biota. The latter is the ultimate proof for successful action.

*More biological data from sites close to the main emission point sources and less affected sites as reference are needed to assess effects and verify recovery. Systematic studies (monitoring programs) should be conducted on key species and variables that indicate changes in the structure and function of terrestrial ecosystems (including possible recovery) in very polluted areas as well as in 'control' areas. Similarly, more biological data on effects and trends in surface waters should be collected in the most impacted areas of the Arctic. The effects of acidic episodes on biota should be studied with high temporal resolution. In these contexts, AMAP should co-operate and co-ordinate with other relevant programs and projects, such as CAFF and CBMP in obtaining biological data. Systematic lake surveys should be conducted in acid-sensitive regions of the Arctic.*

*Sites for effects monitoring studies should be located close to air quality, deposition, and climate monitoring stations ('multi-purpose stations', see above). The study sites should be representative of the different terrestrial ecosystem types (tundra, mountain birch, and coniferous forest ecosystems) because these differ in their sensitivity to air pollution impacts.*

*Limnological and accurately-dated high-resolution paleolimnological studies are needed in lakes near emission sources in the environmentally sensitive areas of the Kola Peninsula and Norilsk to assess recovery.*

#### 8.6.6. Models

Modeling is one of the most important tools available for gaining insight into the possible pollution status of the extensive areas of the Arctic where the observational networks are absent or poorly developed. Models also allow investigation of scenarios for future trends, and for linkages between contaminant pathways and, for example, climate change.

*Measurements of sulfur species, nitrogen species, and black carbon should be used to further validate and improve existing air quality and deposition models. Measurements should be conducted during field campaigns to improve models, for example determining dry deposition velocities during summer.*

*There is a need to integrate aircraft, ground-based, and long-term data sets for use in three-dimensional arctic climate models designed to evaluate climate forcing by arctic haze. Existing data are primarily long-term ground-based measurements of aerosol composition and, at a few sites, scattering and absorption. To estimate aerosol radiative forcing in the Arctic requires additional data on the vertical and horizontal distributions of composition and amount of the haze as well as its variability. Such data are only derived from periodic aircraft measurements and long-term ground-based observations.*

# References

- Aamlid, D., 1993. Symptoms of sulphur dioxide injuries on some boreal plants. In: R. Jalkanen and T. Aalto (eds.). Forest pathological research in northern forests with special reference to abiotic stress factors. Extended SNS meeting in forest pathology in Lapland, Finland, 3-7 August 1992. Metsäntutkimuslaitoksen tiedonantoja, 451:93-102.
- Aamlid, D. and I. Skogheim, 2001. The occurrence of *Hypogymnia physodes* and *Melanelia olivacea* lichens on birch stems in northern boreal forest influenced by local air pollution. Norsk geografisk Tidsskrift, 55:94-98.
- Aamlid, D., H. Tømmervik, M. Gytarsky, R. Karaban, K. Venn, T. Rindal, N. Vassilieva, G. Koptsik and E. Løbersli, 1995. Determination of exceedance of critical levels in the border area between Norway and Russia. In: E. Løbersli and K. Venn (eds.). Effect of air pollution on terrestrial ecosystems in the border area between Norway and Russia. Proceedings of the second symposium, Svanvik, Norway, 3-5 October 1994, pp. 19-23. Directorate for Nature Management, Trondheim, Norway.
- Aamlid, D., N. Vassilieva, P.A. Aarrestad, M.L. Gytarsky, S. Lindmo, R. Karaban, V. Korotkov, T. Rindal, V. Kuzmicheva and K. Venn, 2000. The ecological state of the ecosystems in the border area between Norway and Russia. Boreal Environment Research, 5:257-278.
- Aarrestad, P.A. and D. Aamlid, 1999. Vegetation monitoring in South-Varanger, Norway – species composition of ground vegetation and its relation to environmental variables and pollution impact. Environmental Monitoring and Assessment, 58:1-21.
- ACIA, 2004. Impacts of a Warming Arctic: Arctic Climate Impact Assessment. Cambridge University Press. 139p.
- Aherne, J. and S.A. Watmough, 2005. Critical loads of acidity for North America. Conference Abstracts, Acid Rain 2005, 7th International Conference on Acid Deposition, p. 476. Prague, Czech Republic, 12-17 June 2005.
- Ahlbrandt, T.S., 2002. Future petroleum energy resources of the world. International Geology Review, 44:1092-1104.
- Albrecht, B.A., 1989. Aerosols, cloud microphysics, and fractional cloudiness. Science, 245:1227-1230.
- Alekseev, A.S. and A.R. Soroka, 2002. Scots pine growth trends in northwestern Kola Peninsula as an indicator of positive changes in the carbon cycle. Climatic Change, 55:183-196.
- Alexeyev, V.A., 1995. Impacts of air pollution on far north forest vegetation. Science of the Total Environment, 160/161:605-617.
- Allen, D.J., J.E. Dibb, B. Ridley, K.E. Pickering and R.W. Talbot, 2003. An estimate of the stratospheric contribution to springtime tropospheric ozone maxima using TOPSE measurements and beryllium-7 simulations. Journal of Geophysical Research, 108:8355, doi:10.1029/2001JD001428.
- Allen-Gil, S.M., C.P. Gubala, D.H. Landers, B.K. Lasorsa, E.A. Crecelius and L.R. Curtis, 1997. Heavy metal accumulation in sediment and freshwater fish in U.S. Arctic lakes. Environmental Toxicology and Chemistry, 16:733-741.
- AMAP, 1997. Arctic Pollution Issues: A State of the Arctic Environment Report. Arctic Monitoring and Assessment Programme. xii+188p.
- AMAP, 1998. AMAP Assessment Report: Arctic Pollution Issues. Arctic Monitoring and Assessment Programme. xii+859p.
- AMAP, 2002. Arctic Pollution 2002. Persistent Organic Pollutants, Heavy Metals, Radioactivity, Human Health, Changing Pathways. Arctic Monitoring and Assessment Programme, Oslo. xii+112p.
- AMAP, 2003. AMAP Assessment 2002: Human Health in the Arctic. Arctic Monitoring and Assessment Programme, Oslo, Norway. xiv + 137p.
- AMAP, 2005. AMAP Assessment 2002: Heavy Metals in the Arctic. Arctic Monitoring and Assessment Programme, Oslo, Norway. xvi + 265p.
- American Thoracic Society, 2000. What constitutes an adverse health effect of air pollution? Official statement of the American Thoracic Society. American Journal of Respiratory and Critical Care Medicine; 161:665-673.
- Andreae, M.O., D. Rosenfeld, P. Artaxo, A.A. Costa, G.P. Frank, K.M. Longo and M.A.F. Silva-Dias, 2004. Smoking rain clouds over the Amazon. Science, 303:1337-1342.
- Antoniades, D., M.S.V. Douglas and J.P. Smol, 2003a. Comparative physical and chemical limnology of two Canadian High Arctic regions: Alert (Ellesmere Island, NU) and Mould Bay (Prince Patrick Island, NWT). Archiv für Hydrobiologie, 158:485-516.
- Antoniades, D., M.S.V. Douglas and J.P. Smol, 2003b. The physical and chemical limnology of 24 ponds and one lake from Isachsen, Ellef Ringnes Island, Canadian High Arctic. International Review of Hydrobiology, 88:519-538.
- Äyräs, M. and G. Kashulina, 2000. Regional patterns of element contents in the organic horizon of podzols in the central part of the Barents region (Finland, Norway and Russia), with special reference to heavy metals (Co, Cr, Fe, Ni, Pb, V and Zn) and sulphur as indicators of airborne pollution. Journal of Exploration Geochemistry, 68:127-144.
- Äyräs, M., H. Niskavaara, I. Bogatyrev, V. Chekushin, V. Pavlov, P. de Caritat, J.H. Halleraker, T.E. Finne, G. Kashulina and C. Reimann, 1997. Regional patterns of heavy metals (Co, Cr, Cu, Fe, Ni, Pb, V and Zn) and sulphur in terrestrial moss samples as indication of airborne pollution in a 188,000 km<sup>2</sup> area in northern Finland, Norway and Russia. Journal of Geochemical Exploration, 58:269-281.
- Bååth, E., 1989. Effects of heavy metals in soil on microbial processes and populations (a review). Water Air and Soil Pollution, 47:335-379.
- Bækken, T. and K.J. Aanes, 1990. Bruk av vassdragets bunnsfauna i annkvalitetsklassifisering. NIVA-rapport O-87119, Norwegian Institute for Water Research, Oslo. (In Norwegian)
- Baker, J.P., D.P. Bernard, S.W. Christensen, M.J. Sale, J. Freda, K. Heltcher, D. Marmorek, L. Rowe, P. Scanlon, G. Suter, W. Warren-Hicks and P. Welbourn, 1990. Biological effects of changes in surface water acid-base chemistry. NAPAP Rep. 13. In: P.M. Irving (ed.), Acidic Deposition: State of Science and Technology. Vol. II. Aquatic Processes and Effects, U.S. National Acid Precipitation Assessment Program, Washington, D.C. 381p.
- Barlaup, B.T. and Å. Åtland, 1996. Episodic mortality of brown trout (*Salmo trutta* L.) caused by sea-salt-induced acidification in western Norway: effects on different life stages within three populations. Canadian Journal of Fisheries and Aquatic Science, 53:1835-1843.
- Barrie, L.A., 1986. Arctic air pollution: An overview of current knowledge. Atmospheric Environment, 20:643-663.
- Barrie, L.A., 1996. Occurrence and trends of pollution in the Arctic troposphere. In: E.W. Wolff and R.C. Bales (eds.), Chemical Exchange between the Atmosphere and Polar Snow, pp. 93-130. NATO ASI Series 1: Global Environmental Change 43.
- Barrie, L.A. and R.M. Hoff, 1984. The oxidation rate and residence time of sulphur dioxide in the Arctic atmosphere. Atmospheric Environment, 18:2711-2722.
- Barrie, L.A., R.M. Hoff and S.M. Daggupaty, 1981. The influence of mid-latitude pollution sources on haze in the Canadian Arctic. Atmospheric Environment, 15:1407-1419.
- Battarbee, R.W., V.J. Jones, R.J. Flower, N.G. Cameron, H. Bennion, L. Carvalho and S. Juggins, 2001. Diatoms. In: E.F. Stoermer, H.J.B. Birks, W.M. Last (eds.). Tracking Environmental Change Using Lake Sediments, Vol. 3, Terrestrial, Algal and Siliceous Indicators, pp. 155-202. Kluwer Academic Publishers.
- Belikova T.V., V.N. Vasilenko, I.M. Nararov, A.N. Pegoeva and Kh. D. Fridman, 1984. Characterization of snow cover background pollution by sulphur at the territory of USSR. Meteorology and hydrology, 9.
- Berresheim, H., P.H. Wine and D.D. Davis, 1995. Sulphur in the atmosphere. In: H.B. Singh (ed.), Composition, Chemistry, and Climate of the Atmosphere, pp. 251-307. Van Nostrand Reinhold.
- Betts-Piper, A.M., B.A. Zeeb and J.P. Smol, 2004. Distribution and autecology of chrysophyte cysts from high Arctic Svalbard lakes: preliminary evidence of recent environmental change. Journal of Paleolimnology, 31:467-481.
- Bigler, C. and R.I. Hall, 2002. Diatoms as indicators of climatic and limnological change in Swedish Lapland: a 100-lake calibration set and its validation for paleoecological reconstructions. Journal of Paleolimnology, 27:97-115.
- Bigler, C. and R.I. Hall, 2003. Diatoms as quantitative indicators of July temperature: a validation attempt at century-scale with meteorological data from northern Sweden. Palaeogeography Palaeoclimatology Palaeoecology, 189:147-160.
- Bigler, C., I. Larocque, S.M. Peglar, H.J.B. Birks and R. Hall, 2002a. Quantitative multiproxy assessment of long-term patterns of Holocene environmental change from a small lake near Abisko, northern Sweden. The Holocene, 12:481-496.
- Bigler, M., D. Wagenbach, H. Fischer, J. Kipfstuhl, H. Miller, S. Sommer and B. Stauffer, 2002b. Sulphate record from a northeast Greenland ice core over the last 1200 years based on continuous flow analysis. Annals of Glaciology, 35:250-256.
- Bigler, C., E. Grahm, I. Larocque, A. Jeziorski and R.I. Hall, 2003. Holocene environmental change at Lake Njulla (999 m a.s.l.), northern Sweden: a comparison with four small nearby lakes along an altitudinal gradient. Journal of Paleolimnology, 29:13-29.
- Bishop, K., H. Laudon and S. Köhler, 2000. Separating the natural and anthropogenic components of spring flood pH decline: A method



- for areas that are not chronically acidified. *Water Resources Research*, 30:1873-1889.
- Blais, J.M., K.E. Duff, T.E. Laing and J.P. Smol, 1999. Regional contamination in lakes from the Noril'sk region in Siberia, Russia. *Water Air and Soil Pollution*, 3-4:389-404.
- Blanchet, J.-P. and E. Girard, 1995. Water-vapor temperature feedback in the formation of continental arctic air: implications for climate. *Science of the Total Environment*, 160/161:793-802.
- Blanchet, J.-P. and R. List, 1987. Estimation of optical properties of Arctic haze using a numerical model. *Atmosphere-Ocean*, 21:444-464.
- Blaustein, A.R., J.M. Romansic, J.M. Kiesecker and A.C. Hatch, 2003. Ultraviolet radiation, toxic chemicals and amphibian population declines. *Diversity and Distributions*, 9:123-140.
- Bobbink, R., M. Ashmore, S. Braun, W. Flückiger and I.J.J. Van den Wyngaert, 2002. Empirical nitrogen critical loads for natural and semi-natural ecosystems: 2002 update. Background document for the Expert Workshop on Empirical Critical Loads for Nitrogen on (Semi-)natural Ecosystems – Berne Switzerland, 11-13 November. UN/ECE Convention on Long-range Transboundary Air Pollution.
- Bodhaine, B.A., 1989. Barrow surface aerosol: 1976–1986. *Atmospheric Environment*, 23:2357-2369.
- Bodhaine, B.A. and E.G. Dutton, 1993. A long-term decrease in Arctic haze at Barrow, Alaska. *Geophysical Research Letters*, 20:947-950.
- Boltneva, L.I., A.A. Ignatiev, R.T. Karaban, I.M. Nazarov, I.A. Rudneva and T.I. Sysygina, 1982. Aerosol and gaseous industrial emission impact on Northern Taiga Pine Forests. *Ecology*, 4:37-43. (In Russian)
- Borys, R.D., 1989. Studies of ice nucleation by arctic aerosol on AGASP-II. *Journal of Atmospheric Chemistry*, 9:169-185.
- Bottenheim, J.B., L.A. Barrie and E. Atlas, 1993. The partitioning of nitrogen oxides in the lower arctic troposphere during spring 1988. *Journal of Atmospheric Chemistry*, 17:15-28.
- Bowling, S.A. and G.E. Shaw, 1992. The thermodynamics of pollution removal as an indicator of possible source areas for Arctic Haze. *Atmospheric Environment*, 26A:2953-2961.
- Boyd, R.S. and S.N. Martens, 1994. Nickel hyperaccumulated by *Thlaspi montanum* var. *montanum* is acutely toxic to an insect herbivore. *Oikos*, 70:21-25.
- Brakefield, P.M., 1987. Industrial melanism: do we have the answer? *Trends in Ecology and Evolution*, 2:117-121.
- Brigham, L. and B. Ellis (eds), 2004. Arctic Marine Transport Workshop. Proceedings of a Workshop (Cambridge, UK, 28-30 September 2004). Institute of the North/US Arctic Research Commission/IASC. 18+31p.
- Brock, C.A., L.F. Radke, J.H. Lyons and P.V. Hobbs, 1989. Arctic hazes in summer over Greenland and the North American Arctic, I. Incidence and origins. *Journal of Atmospheric Chemistry*, 9:129-148.
- Brook, J.R., 1995. Wet acid deposition episodicity in eastern North America and the influence of deposition episodes on annual deposition amounts. *Atmospheric Environment*, 29:1795-1807.
- Brooks, S.J. and H.J.B. Birks, 2004. The dynamics of Chironomidae (Insecta: Diptera) assemblages in response to environmental change during the past 700 years on Svalbard. *Journal of Paleolimnology*, 31:483-498.
- Brunekreef, B. and S.T. Holgate, 2002. Air pollution and health. *Lancet*, 360:1233-1242.
- Brunner, I., 2001. Ectomycorrhizas: their role in forest ecosystems under the impact of acidifying pollutants. *Perspectives in Plant Ecology, Evolution and Systematics*, 4:13-27.
- Byrne, R.H., J.G. Acker, B.R. Betzer, R.A. Feely and M.H. Cates, 1984. Water column dissolution of aragonite in the Pacific Ocean. *Nature*, 312:321-326.
- Caldeira, K. and M.E. Wickett, 2003. Anthropogenic carbon and ocean pH. *Nature*, 425:365-365.
- Callaghan, V., 2005. Chapter 7. Arctic Tundra and Polar Desert Ecosystems. *In: Arctic Climate Impact Assessment*, pp. 243-352. Cambridge University Press.
- Carlson, T.N., 1981. Speculations on the movement of polluted air to the Arctic. *Atmospheric Environment*, 15:1473-1477.
- Cess, C.R., 1983. Arctic aerosol: model estimates of the interactive influences upon the surface-troposphere radiation budget. *Atmospheric Environment*, 17:2555-2564.
- Chapman, W.E. and J.E. Walsh, 1993. Recent variations in sea ice and air temperature in high latitudes. *Bulletin of the American Meteorological Society*, 74:33-47.
- Chertov, O.G., M. Nadporozhskaya, I. Lapshina and O. Grigorieva, 1993. Soil degradation in surrounding of "Pechenganikel" smelter complex. *In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.)*. Aerial Pollution on Kola Peninsula, pp.159-162. Proceeding of the International Workshop, April 14-16 1992. St. Petersburg. Kola Science Centre, Apatity.
- Christensen, J., 1997. The Danish Eulerian Hemispheric Model - A Three Dimensional Air Pollution Model used for the Arctic. *Atmospheric Environment*, 31:4169-4191.
- Christensen, J., 1999. An overview of modelling the Arctic mass budget of metals and sulphur: Emphasis on source apportionment of atmospheric burden and deposition. *In: Modelling and Sources: A Workshop on Techniques and Associated Uncertainties in Quantifying the Origin and Long-Range Transport of Contaminants to the Arctic*, Annex 15, 1-5. Report and extended abstracts. Bergen, 14-16 June 1999.
- Christensen, J.H., J. Brandt, L.M. Frohn and H. Skov, 2004. Modelling of mercury in the Arctic with the Danish Eulerian Hemispheric Model. *Atmospheric Chemistry and Physics*, 4:2251-2257.
- Clancy, L., P. Goodman, H. Sinclair and D.W. Dockery, 2002. Effect of air-pollution control on death rates in Dublin, Ireland: an intervention study. *Lancet*, 360:1210-1214.
- Clark, K.L. and R.J. Hall, 1985. Effects of elevated hydrogen ion and aluminum concentration on the survival of amphibian embryos and larvae. *Canadian Journal of Zoology*, 63:116-123.
- Clark, K.L. and B. LaZerte, 1985. A laboratory study on the effects of aluminum and pH on amphibians eggs and tadpoles. *Canadian Journal of Fisheries and Aquatic Sciences*, 42:1544-1551.
- Clarke, A.D., 1989. In-situ measurements of the aerosol size distributions, physicochemistry, and light absorption properties of Arctic haze. *Journal of Atmospheric Chemistry*, 9:255-267.
- Clarke, A.D. and K.J. Noone, 1985. Soot in the Arctic snowpack: A cause for perturbation in radiative transfer. *Atmospheric Environment*, 19:2045-2053.
- Coppola, E., F. Kucharski, F. Giorgi and F. Molteni, 2005. Bimodality of the North Atlantic Oscillation in simulations with greenhouse gas forcing. *Geophysical Research Letters* 32:L23709, doi:10.1029/2005GL024080.
- Cornelissen, J.H.C., T.V. Callaghan, J.M. Alatalo, A. Michelsen, E. Graglia, A.E. Hartley, D.S. Hik, S.E. Hobbie, M.C. Press, C.H. Robinson, G.H.R. Henry, G.R. Shaver, G.K. Phoenix, D.G. Jones, S. Jonasson, F.S. Chapin III, U. Molau, C. Neill, J.A. Lee, J.M. Melillo, B. Sveinbjörnsson and R. Aerts, 2001. Global change and arctic ecosystems: is lichen decline a function of increases in vascular plant biomass? *Journal of Ecology*, 89:984-994.
- Curry, J.A., 1995. Interactions among aerosols, clouds, and climate of the Arctic Ocean. *Science of the Total Environment*, 160/161:777-791.
- Curry, J.A. and G.F. Herman, 1985. Infrared radiative properties of Arctic stratus clouds. *Journal of Climate and Applied Meteorology*, 24:525-538.
- Damoah, R., N. Spichtinger, C. Forster, P. James, I. Mattis, U. Wandinger, S. Beirle and A. Stohl, 2004. Around the world in 17 days – hemispheric-scale transport of forest fire smoke from Russia in May 2003. *Atmospheric Chemistry and Physics*, 4:1311-1321.
- Danielson, R.M. and S. Visser, 1989. Effects of forest soil acidification on ectomycorrhizal and vesicular-arbuscular mycorrhizal development. *New Phytologist*, 112:41-47.
- Dauvalter, V., 1995. Influence of pollution and acidification on metal concentrations in Finnish Lapland lake sediments. *Water Air and Soil Pollution*, 85:853-858.
- Dauvalter, V., 1997. Metal concentrations in sediments in acidifying lakes in Finnish Lapland. *Boreal Environment Research*, 2:369-379.
- Dauvalter, V., 2003. Impact of mining and refining on the distribution and accumulation of nickel and other heavy metals in sediments of subarctic Lake Kuetsjärvi, Murmansk Region, Russia. *Journal of Environmental Monitoring*, 5:210-215.
- Dauvalter, V.A. and S. Rognerud, 2001. Heavy metal pollution in sediments of the Pasvik River drainage. *Chemosphere*, 42:9-18.
- de Caritat, P., M. Åyräs, H. Niskavaara, V. Chekushin, I. Bogatyrev and C. Reimann, 1998. Snow composition in eight catchments in the central Barents Euro-Arctic Region. *Atmospheric Environment*, 32:2609-2626.
- de Caritat, P., G. Hall, S. Gislason, W. Belsey, M. Braun, N.I. Golubeva, H.K. Olsen, J.O. Scheie and J.E. Vaive, 2005. Chemical composition of arctic snow: concentration levels and regional distribution of major elements. *Science of The Total Environment*, 336:183-199.
- Denison, L., R. Simpson, A. Petroeschovsky, L. Thalib and G. Williams, 2001. Ambient air pollution and daily hospital admissions in Melbourne 1994-1997. Melbourne: EPA Victoria; 2001 November. Report No. 789.
- Department of Environment, 2003. Research on air pollution in Perth, morbidity and mortality: A case-crossover analysis 1992–1997. Technical Series 114. Government of Western Australia.
- Deyeva, N.M. and E.A. Maznaja, 1993. The state of the bilberry in polluted and unpolluted forests of the Kola Peninsula. *In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.)*. Aerial pollution on Kola Peninsula, pp. 308-311. Proceedings of the International Workshop, April 14-16 1992. St. Petersburg. Kola Science Centre, Apatity.
- Dibb, J.E., L.D. Meeker, R.C. Finkel, J.R. Southon, M.W. Caffee and



- L.A. Barrie, 1994. Estimation of the stratospheric input to the Arctic troposphere:  $^7\text{Be}$  and  $^{10}\text{Be}$  in aerosols at Alert, Canada. *Journal of Geophysical Research*, 99:12855-12864, doi:10.1029/94JD00742.
- Dibb, J.E., R.W. Talbot, S.I. Whitlow, M.C. Shipham, J. Winterle, J. McConnell and R. Bales, 1996. Biomass burning signatures in the atmosphere and snow at Summit, Greenland: An event on 5 August 1994. *Atmospheric Environment*, 30:553-561.
- Dibb, J.E., R.W. Talbot, E. Scheuer, G. Seid, L. DeBell, B. Lefer and B. Ridley, 2003. Stratospheric influence on the northern North American free troposphere during TOPSE:  $^7\text{Be}$  as a stratospheric tracer. *Journal of Geophysical Research*, 108:8363, doi:10.1029/2001JD001347.
- Dillon, P.J. and B.D. LaZerte, 1992. Response of Plastic Lake catchment, Ontario, to reduced sulphur deposition. *Environmental Pollution*, 77:211-217.
- Dillon, P.J., H.E. Evans and P.J. Scholer, 1988. The effects of acidification on metal budgets of lakes and catchments. *Biogeochemistry*, 5:201-220.
- Dillon, P.J., L.A. Molot and M. Futter, 1997. The effect of El Niño-related drought on the recovery of acidified lakes. *Environmental Monitoring and Assessment*, 46:105-111.
- Disney, R.H.L., E.L. Zvereva and M. Mostovsky, 2001. A scuttle fly (Diptera: Phoridae) parasitizing a beetle (Coleoptera: Chrysomelidae) in Russia. *Entomologica Fennica*, 12:59-63.
- DNV, 1999. Reference values for ship pollution. Report 99-2034. Det Norske Veritas, Norway.
- DNV, 2003. Utredning av helårig petroleumsvirksomhet i området Lofoten – Barentshavet: Konsekvenser for og fra skipstrafikk. ULB Studie Nr. 14, Rapport 2003-0331. Det Norske Veritas, Norway.
- Doka, S.E., D.K. McNicol, M.L. Mallory, I. Wong, C.K. Minns and N.D. Yan, 2003. Assessing potential for recovery of biotic richness and indicator species due to changes in acidic deposition and lake pH of five areas of southeastern Canada. *Environmental Monitoring and Assessment*, 88:53-101.
- Dotterud, L.K., J.O. Odland and E.S. Falk, 2001. Atopic diseases among school children in Nikel, Russia, an Arctic area with heavy air pollution. *Acta Dermato-Venereologica*, 81:198-201.
- Douglas, M.S.V., J.P. Smol and W. Blake, Jr., 1994. Marked post-18th century environmental change in high-arctic ecosystems. *Science*, 266:416-419.
- Douglas, T.A. and M. Sturm, 2004. Arctic haze, mercury, and the chemical composition of snow across northwestern Alaska. *Atmospheric Environment*, 38:805-820.
- Downing, R.J., J.-P. Hettelingh and P.A.M. de Smet, 1993. Calculation and mapping of critical loads in Europe: Status report 1993. Coordination Center for Effects, National Institute for Public Health and Environmental Protection, Bilthoven, The Netherlands.
- Dudley, N. and S. Stolton, 1996. Air pollution and Biodiversity: a review. WWF International.
- Duff, K.E., T.E. Laing, J.P. Smol and D.R.S. Lean, 1998. Limnological characteristics of lakes located across arctic treeline in northern Russia. *Hydrobiologia*, 391:205-222.
- Duncan, B.N. and I. Bey, 2004. A modeling study of the export pathways of pollution from Europe: Seasonal and interannual variations (1987-1997). *Journal of Geophysical Research*, 109, D08301, doi:10.1029/2003JD004079.
- Dutton, E.G., J.J. DeLuisi and B.A. Bodhaine, 1984. Features of aerosol optical depth observed at Barrow, March 10-20, 1983. *Geophysical Research Letters*, 11:385-388.
- Eckhardt, S., A. Stohl, S. Beirle, N. Spichtinger, P. James, C. Forster, C. Junker, T. Wagner, U. Platt and S.G. Jennings, 2003. The North Atlantic Oscillation controls air pollution transport to the Arctic. *Atmospheric Chemistry and Physics*, 3:1769-1778.
- Eeva, T., E. Lehtikoinen and M. Nikinmaa, 2003. Pollution-induced nutritional stress in birds: An experimental study of direct and indirect effects. *Ecological Applications*, 13:1242-1249.
- Eilers, J.M., D.H. Landers, A.D. Newell, M.E. Mitch, M. Morrison and J. Ford, 1993. Major ion chemistry of lakes on the Kenai Peninsula, Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*, 50:816-826.
- Ekimov, S.V., I.V. Samodova, I.M. Petrov, V.V. Troitsky and M.A. Burstein, 2001. Russian smelter emissions. *Mining Journal*, London, November 23, 2001, pp. 393.
- EMEP, 2004. EMEP Assessment. Part I, European perspective. G. Löwblad, L. Tarrason, K. Tørseth and S. Dutchak (eds). Norwegian Meteorological Institute, Oslo.
- Emery, C.A., R. Haberle and T.P. Ackermann, 1992. A one-dimensional modeling study of carbonaceous haze effects on the springtime Arctic environment. *Journal of Geophysical Research*, 97:20599-20613.
- Erkinaro, J., E. Niemelä, H. Erkinaro and M. Rask, 1992. Monitoring of the possible effects of acidification on fish populations and zoobenthos of rivers and lakes in northeastern Finnish Lapland. In: K. Kinnunen and M. Varmola (eds.). *Effects of Air Pollutants and Acidification in Combination with Climatic Factors on Forests, Soils and Waters in Northern Fennoscandia*, pp. 168-171. Nordic Council of Ministers, Copenhagen.
- Erkinaro, H., J. Erkinaro, M. Rask and E. Niemelä, 2001. Status of zoobenthos and fish populations in subarctic rivers of northernmost Finland: possible effects of acid emissions from Russian Kola Peninsula. *Water Air and Soil Pollution*, 130:831-836.
- Erola, U.-R., 1999. *Herkät pohjoiset alueet ilmastomuutoksen kuvastajina – piilevästratigrafinen tutkimus subarktista lammista Kilpisjärvellä*. Unpublished M.Sc. Thesis, University of Helsinki, Helsinki, Finland, 90p. (In Finnish).
- Evans, C.D. and D. Monteith, 2001. Trends in surface water chemistry at AWMN sites 1988–2000: Evidence for recent recovery at a national scale. *Hydrology and Earth System Sciences*, 5:351-366.
- Evans, C.D., J. Cullen, C. Alewell, J. Kopáček, A. Marchetto, F. Moldan, A. Prectel, M. Rogora, J. Vesely and R.F. Wright, 2001. Recovery from acidification in European surface waters. *Hydrology and Earth System Sciences*, 5:283-298.
- Evans, C.D., C. Freeman, D. Monteith, B. Reynolds and N. Fenner, 2002. Climate change – terrestrial export of organic carbon: Communication arising. *Nature*, 415:861-862.
- Evans, C.D., D.T. Monteith and D.M. Cooper, 2005. Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution*, 137:55-71.
- Evdokimova, G.A., 2000. The impact of heavy metals on the microbial diversity of podzolic soils in the Kola Peninsula. Forest dynamics in heavily polluted regions. Report no. 1 of the IUFRO Task Force on Environmental Change. pp. 67-76.
- Evdokimova, G.A., 2001. Fluorine in the soils of the White Sea Basin and bioindication of pollution. *Chemosphere*, 42:35-43.
- Evdokimova, G.A., E.V. Beresneva, E.V. Lebedeva and M.V. Scjatkovskaja, 2004. The influence of fluorine on the soil fungi of forest podzols in the impact zone of aluminium plant. *Mikologija I Fitopatologija*, 38:66-70. (In Russian)
- Evseev, A.V., V.I. Nikolaev and T.G. Sukhova, 2000. Geochemical pattern in macro and microelements distribution in snow cover and glaciers of Severnaya Zemlya. *Data of Glaciological Studies*, 89:11-17.
- Ferek, R.J., P.V. Hobbs, L.F. Radke, J.A. Herring, W.T. Sturges and G.F. Cota, 1995. Dimethyl sulfide in the arctic atmosphere. *Journal of Geophysical Research*, 100:26093-26104.
- Findlay, S.E.G., 2005. Increased carbon transport in the Hudson River: unexpected consequence of nitrogen deposition? *Frontiers in Ecology and the Environment*, 3:133-137.
- Fjeld, E., S. Rognerud and E. Steinnes, 1994. Influence of environmental factors on heavy metal concentration in lake sediments in southern Norway indicated by path analysis. *Canadian Journal of Fisheries and Aquatic Sciences*, 51:1708-1720.
- Flanagan, K.M., E. McCauley, F. Wrona and T. Prowse, 2003. Climate change: the potential for latitudinal effects on algal biomass in aquatic ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences*, 60:635-639.
- Folt, C.L., C.Y. Chen, M.V. Moore and J. Burnaford, 1999. Synergism and antagonism among multiple stressors. *Limnology and Oceanography*, 44:864-877.
- Forsius, M., S. Kleemola, J. Vuorenmaa and S. Syri, 2001. Fluxes and trends of nitrogen and sulphur compounds at Integrated Monitoring Sites in Europe. *Water Air and Soil Pollution*, 130:1641-1648.
- Forsius, M., J. Vuorenmaa, J. Mannio and S. Syri, 2003. Recovery from acidification of Finnish lakes: regional patterns and relation to emission reduction policy. *Science of The Total Environment*, 310:121-132.
- Forster, C., U. Wandering, G. Wotawa, P. James, I. Mattis, D. Althausen, P. Simmonds, S. O'Doherty, C. Kleefeld, S.G. Jennings, J. Schneider, T. Trickl, S. Kreipl, H. Jäger and A. Stohl, 2001. Transport of boreal forest fire emissions from Canada to Europe. *Journal of Geophysical Research*, 106:22887-22906.
- Foster, J.S., J.W. Winchester and E.G. Dutton, 1992. The date of snow disappearance on the Arctic tundra as determined from satellite, meteorological station and radiometric in situ observations. *IEEE Transactions on Geoscience and Remote Sensing*, 30:793-798.
- Framstad, E., 2002. *Terrestrisk naturovervåking. Markvegetasjon, epifytter, smånagere og fugl i TOV-områdene*, 2002. Norsk Institutt for Naturforskning, 70p. (In Norwegian)
- Frantzen, B. and A. Bambulyak, 2003. Oil transport from the Russian part of the Barents Region. On July 2003. Report, Svanhovd Miljøsester, Planteforsk. Barents Secretariat, Norway.
- Freda, J. and D. McDonald, 1990. Effects of aluminum on the leopard frog *Rana pipiens*: life stage comparisons and aluminum uptake. *Canadian Journal of Fisheries and Aquatic Sciences*, 47:210-216.
- Freedman, B. and S. Beauchamp, 1998. Implications of atmospheric change for biodiversity of aquatic ecosystems in Canada. *Environmental Monitoring and Assessment*, 49:271-280.
- Freeman, C., C.D. Evans, D.T. Monteith, B. Reynolds and N. Fenner, 2001. Export of organic carbon from peat soils. *Nature*, 412:785-785.
- Freeman, C., N. Fenner, N.J. Ostle, H. Kang, D.J. Dowrick, B. Reynolds,

- M.A. Lock, D. Sleep, S. Hughes and J. Hudson, 2004. Export of dissolved organic carbon from peatlands under elevated carbon dioxide levels. *Nature*, 430:195-198.
- Fremstad, E., J. Paal and T. Möls, 2005. Impacts of increased nitrogen supply on Norwegian lichen-rich alpine communities: a 10-year experiment. *Journal of Ecology*, 93:471-481.
- Frohn, L.M., J.H. Christensen and J. Brandt, 2002. Development of a high resolution nested air pollution model – the numerical approach. *Journal of Computational Physics*, 179:68-94.
- Frohn, L.M., J.H. Christensen, J. Brandt, C. Geels and K.M. Hansen, 2003. Validation of a 3-D hemispheric nested air pollution model. *Atmospheric Chemistry and Physics Discussions*, 3:3543-3588.
- Fromm, M., R. Bevilacqua, R. Servranckx, J. Rosen, J.P. Thayer, J. Herman and D. Larko, 2005. Pyro-cumulonimbus injection of smoke to the stratosphere: Observations and impact of a super blowup in northwestern Canada on 3-4 August 1998. *Journal of Geophysical Research*, 110:D08205, doi:10.1029/2004/JD005350.
- Gaare, E. and H. Tømmervik, 2000. Monitoring lichen grazing in East Finnmark. NINA Oppdragsmelding 669. Norwegian Institute For Nature, pp. 1-28.
- Galloway, J.N., 1995. Acid deposition: perspectives in time and space. *Water, Air and Soil Pollution*, 85:15-24.
- Garland, J.A., 1977. The dry deposition of sulphur dioxide to land and water surfaces. *Proceedings of the Royal Society of London A*, 354:245-268.
- Garrett, T.J., L.F. Radke and P.V. Hobbs, 2002. Aerosol effects on the cloud emissivity and surface longwave heating in the Arctic. *Journal of Atmospheric Sciences*, 59:769-778.
- Garrett, T.J., C. Zhao, X. Dong, G.G. Mace and P.V. Hobbs, 2004. Effects of varying aerosol regimes on low-level Arctic stratus. *Geophysical Research Letters*, 31, doi:10.1029/2004GL019928.
- Geels, C., S. Doney, R. Dargaville, J. Brandt and J.H. Christensen, 2004. Investigating the sources of synoptic variability in atmospheric CO<sub>2</sub> measurements over the northern hemisphere continents: A regional model study. *Tellus*, 56B:35-50.
- Gilbert, R.O., 1987. Statistical methods for environmental pollution monitoring. Van Nostrand Reinhold, 320p.
- Gilmour, C.C. and E.A. Henry, 1991. Mercury methylation in aquatic systems affected by acid deposition. *Environmental Pollution*, 71:131-169.
- Gilyazov, A.S., 1993. Air pollution impact on the bird communities of the Lapland biosphere reserve. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 383-390. *Proceedings of the International Workshop*, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Gilyazov, A.S. and G.D. Kataev, 1990. An experience of zooinidication of industrial pollution in the Kola Peninsula. In: Y.D. Nukhimovskaja and L.A. Bibikova (eds.). *Anthropogenic Impacts on the Nature of Reserves*. Central Research Laboratory of Game and Reserves, Moscow, pp. 5-24. (In Russian)
- Gilyazova, E.V., 1993. Pollution impact on soil mesofauna of Lapland Biosphere Reserve. In: M.V. Kozlov, E. Haukioja, V.T. Yarmishko (eds.). *Aerial Pollution in the Kola Peninsula*, pp. 366-368. *Proceedings of the International Workshop*, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Girard, E., J.-P. Blanchet and Y. Dubois, 2005. Effects of arctic sulphuric acid aerosols on wintertime low-level atmospheric ice crystals, humidity and temperature at Alert, Nunavut. *Atmospheric Research*, 73:131-148.
- Givélet, N., F. Roos-Barraclough, M.E. Goodsite, A.K. Cheburkin and W. Shoty, 2004. Atmospheric mercury accumulation rates between 5900 and 800 calibrated years BP in the high arctic of Canada recorded by peat hummocks. *Environmental Science and Technology*, 38:4964-4972.
- Głowacki, P., 1997. The mass balance of Hans glacier in the light of cryochemical investigations. *Polish Polar Studies (24th Polar Symposium)*, Warszawa, pp. 75-79.
- Gordon, C., J.M. Wynn and S.J. Woodin, 2001. Impacts of increased nitrogen supply on high Arctic heath: the importance of bryophytes and phosphorus availability. *New Phytologist*, 149:461-471.
- Gorham, E., 1998. Acid deposition and its ecological effects: a brief history of research. *Environmental Science and Policy*, 1:153-166.
- Goryainova, V. and V. Nikonov, 1997. Heavy metals in pine ecosystems of the Kola Peninsula subjected to acid precipitation. In: K. Aoyama, K. Katoh, T. Murano, T. Paces and Y. Taguchi (eds.). *Acid Snow and Rain*, pp. 463-467. *Proceedings of International Congress of Acid Snow and Rain*, Niigata University, Japan.
- Goto-Azuma, K., R.M. Koerner and D.A. Fisher, 2002. An ice core record over the last two centuries from Penny Ice Cap, Baffin Island, Canada. *Annals of Glaciology*, 35:29-35.
- Greenaway, K.R., 1950. Experiences with Arctic flying weather, Royal Meteorological Society Canadian Branch, Toronto.
- Gregorauskiene, V., R. Salminen, C. Reimann and V. Chekushin, 2000. Field Manual for Barents Ecogeochemistry project. Geological Survey of Finland, Report S/44/0000/2/2000.
- Gregurek, D., F. Melcher, V. Pavlov, C. Reimann and E.F. Stumpfl, 1999. Mineralogy and mineral chemistry of snow filter residues in the vicinity of the nickel-copper processing industry, Kola Peninsula, NW Russia. *Mineralogy and Petrology*, 65:87-111.
- Grell, G.A., J. Dudhia and D.R. Stauffer, 1994. A description of the fifth-generation Penn State/NCAR mesoscale model (MM5). NCAR Tech. Note TN-398+STR, 122 pp. [Available from UCAR Communications, P.O. Box 3000, Boulder, CO 80307.]
- Grönlund, T. and T. Kauppila, 2002. Holocene history of Lake Soldatskoje (Kola Peninsula, Russia) inferred from sedimentary diatom assemblages. *Boreas*, 31:273-284.
- Guerreiro, C., T. Larssen and P.A. Aarrestad, 2003. Konsekvenser av utslipp til luft av helårig petroleumsvirksomhet i området Lofoten – Barentshavet. NILU OR 16/2003. (In Norwegian)
- Gytarsky, M.L., R.T. Karaban, I.M. Nazarov, T.I. Sysygina and M.V. Chemeris, 1995. Monitoring of forest ecosystems in the Russian subarctic: effects of industrial pollution. *Science of the Total Environment*, 164:57-68.
- Gytarsky, M.L., R.T. Karaban and I.M. Nazarov, 1997. On the assessment of sulphur deposition on forests growing over the areas of industrial impact. *Environmental Monitoring and Assessment*, 48:125-137.
- Hagen, L.O., B. Sivertsen and K. Arnesen, 2005. Grenseområdene i Norge og Russland – Luft- og nedbørkvalitet, april 2003-mars 2004. Norsk institutt for luftforskning (NILU). Rapport NILU OR 61/2004. 74 pp. (In Norwegian with English Summary: Air Quality Monitoring in the Border Areas of Norway and Russia - Progress Report April 2004-March 2005).
- Hämäläinen, J. and P. Huttunen, 1998. Macroinvertebrate-inferred stream-water acidity in north eastern Finland along a sulphur deposition gradient. *Water Air and Soil Pollution*, 104:223-236.
- Hammar, J., 1989. Freshwater ecosystems of polar regions: Vulnerable resources. *Ambio*, 19:6-22.
- Hansen, A.D.A. and H. Rosen, 1984. Vertical distribution of particulate carbon, sulphur and bromine in the Arctic haze and comparison with ground-level measurements at Barrow, Alaska. *Geophysical Research Letters*, 11:381-384.
- Hansen, J. and L. Nazarenko, 2004. Soot climate forcing via snow and ice albedos. *Proceedings of the National Academy of Sciences of the United States of America*, 101(2):423-428, doi:10.1073/pnas.2237157100
- Hansen, K.M., J.H. Christensen, J. Brandt, L.M. Frohn and C. Geels, 2004. Modelling atmospheric transport of I-hexachlorocyclohexane in the Northern Hemisphere with a 3-D dynamic model: DEHM-POP. *Atmospheric Chemistry and Physics*, 4:1-13.
- Hanski, I., H. Henttonen, E. Korpimäki, L. Oksanen and P. Turchin, 2001. Small-rodent dynamics and predation. *Ecology*, 82:1505-1520.
- Hansson, L., 1999. Intraspecific variation in dynamics: small rodents between food and predation in changing landscapes. *Oikos*, 86:159-169.
- Harriman, R., A.W. Watt, A.E.G. Christie, P. Collen, D.W. Moore, A.G. McCartney, E.M. Taylor and J. Watson, 2001. Interpretation of trends in acidic deposition and surface water chemistry in Scotland during the last three decades. *Hydrology and Earth System Sciences*, 5:407-420.
- Havas, M. and T.C. Hutchinson, 1983. The smoking hills: natural acidification of an aquatic ecosystem. *Nature*, 301:23-27.
- Heidam, N.Z., J. Christensen, P. Wahlin and H. Skov, 2004. Arctic atmospheric contaminants in NE Greenland: levels, variations, origins, transport, transformations, and trends 1990-2001. *Science of the Total Environment*, 331:5-28.
- Heintzenberg, J., 1980. Particle size distribution and optical properties of Arctic haze. *Tellus*, 32:251-260.
- Heintzenberg, J. and S. Larssen, 1983. SO<sub>2</sub> and SO<sub>4</sub> in the Arctic. Interpretation of observations at three Norwegian Arctic-subarctic stations. *Tellus*, 35B:255-265.
- Heintzenberg, J., T. Tuch, B. Wehner, A. Wiedensohler, A. Ansmann, I. Mattis, D. Müller, M. Wendisch, S. Eckhardt and A. Stohl, 2002. Arctic Haze over Central Europe. *Tellus*, 55B:796-807.
- Heinz, G.H., 1989. How lethal are sublethal effects. *Environmental Toxicology and Chemistry*, 8:463-464.
- Hejzlar, J., M. Dubrovský, J. Buchtele and M. Ružička, 2003. The apparent and potential effects of climate change on the inferred concentration of dissolved organic matter on a temperate stream (The Málse River, South Bohemia). *Science of the Total Environment*, 310:143-152.
- Heliövaara, K. and R. Väisänen, 1993. *Insects and Pollution*. CRC Press.
- Helle, T. and I. Kojola, 1992. Harmaaporonjäkälän kasvuvaihtelu Itä-Fennoskandiassa. In: H. Kauhanen and M. Varmola (eds.). *Itä-Lapin metsävaurioprojektin väliraportti*. The Lapland Forest Damage Project – Interim Report, pp. 106-114. Finnish Forest Research Institute, Rovaniemi, Research Paper 413. (In Finnish, English abstract: The variation in the growth rate of *Cladonia rangiferina* in eastern Fennoscandia).
- Helsel, D.R. R.M. and Hirsch, 1995. *Statistical Methods in Water Re-*

## References

- sources. *Studies in Environmental Sciences* 49. Elsevier Science B.V.
- Henriksen, A. and M. Posch, 2001. Steady-state models for calculating critical loads of acidity for surface waters. *Water Air and Soil Pollution: Focus* 1:375-398.
- Henriksen, A., L. Lien, T.S. Traaen, I. Sevaldrud and D.F. Brakke, 1988. Lake acidification in Norway – present and predicted chemical status. *Ambio*, 17:259-266.
- Henriksen, A., J. Kämäri, M. Posch and A. Wilander, 1992. Critical loads of acidity: Nordic surface waters. *Ambio*, 21:356-363.
- Henriksen, A., J. Kämäri, M. Posch, M. Forsius, A. Wilander, H. Baxendale and T. Tarvainen, 1994. Critical loads for acidification of surface waters in Northern Fennoscandia (Nordkalotten): Nordkalottkommiteens publikasjonsserie Rapport 33.
- Henriksen, A., B.L. Skjelkvåle, J. Mannio, A. Wilander, J.P. Jensen, T. Moiseenko, R. Harriman, T.S. Traaen, E. Fjeld, J. Vuorenmaa, P. Kortelainen and M. Forsius, 1997a. Results of national lake surveys 1995 in Finland, Norway, Sweden, Denmark, Russian Kola, Russian Karelia, Scotland and Wales, Report 47/97. Norwegian Institute of Water Research, Oslo, Norway, 43p.
- Henriksen, A., J. Mannio, A. Wilander, T. Moiseenko, T.S. Traaen, B.L. Skjelkvåle, E. Fjeld and J. Vuorenmaa, 1997b. Regional Lake Surveys in The Barents region of Finland – Norway – Sweden and Russian Kola 1995 – Results. NIVA-report SNO 3633-97, Acid Rain Research Report 45/97. 36p.
- Henriksen, A., B.L. Skjelkvåle, J. Mannio, A. Wilander, R. Harriman, C. Curtis, J.P. Jensen and T. Moiseenko, 1998. Northern European Lake Survey – 1995. Finland, Norway, Sweden, Denmark, Russian Kola Russian Karelia, Scotland and Wales. *Ambio*, 27:80-91.
- Herber, A., L.W. Thomason, H. Gernandt, U. Leiterer, D. Nagel, K.-H. Schulz, J. Kaptur, T. Albrecht and J. Not, 2002. Continuous day and night aerosol optical depth observations in the Arctic between 1991 and 1999. *Journal of Geophysical Research*, 107(D10): 10.1029/2001JD000536.
- Hesthagen, T., H.M. Berger and B.M. Larsen, 1992. Fish population studies in acid-sensitive areas in eastern Finnmark, northern Norway. In: K. Kinnunen and M. Varmola (eds.). *Effects of Air Pollutants and Acidification in Combination with Climatic Factors on Forests, Soils and Waters in Northern Fennoscandia*, pp. 172-176. Nordic Council of Ministers, Copenhagen.
- Hesthagen, T., A. Langeland and H.M. Berger, 1998. Effects of acidification due to emissions from the Kola Peninsula on fish populations in lakes near the Russian border in northern Norway. *Water Air and Soil Pollution*, 102:17-36.
- Hettelingh J.-P., M. Posch and P.A.M. De Smet, 2001. Multi-effect critical loads used in multi-pollutant reduction agreements in Europe. *Water Air and Soil Pollution*, 130:1133-1138.
- Hettelingh, J.-P., Slootweg, J., and M. Posch, 2004. Critical Loads and Dynamic Modelling Results. 259101014/2004, RIVM, Bilthoven. 134p.
- Hillamo, R.E., V.-M. Kerminen, W. Maenhaut, J.-L. Jaffrezou, S. Balachandran and C.I. Davidson, 1993. Size distributions of atmospheric trace elements at Dye 3, Greenland – 1. Distribution characteristics and dry deposition velocities. *Atmospheric Environment*, 27A:2787-2803.
- Hilmo, O. and H.C. Larsen, 1994. Morphology of epiphytic lichens in areas with different air quality. *Stiftelsen Forskning I Trondheim, Universitetet Trondheim, ALLFORSK Report 2*, pp. 1-44.
- Hirsch, R.M., J.R. Slack and R.A. Smith, 1982. Techniques of trend analysis for monthly water quality analysis. *Water Resources Research*, 18:107-121.
- Hobbie, J.E., 1984. Polar limnology. In: F.E. Taub (ed.). *Ecosystems of the World. 23. Lakes and Rivers*. Science Publication 22, Elsevier.
- Hobbie, J.E., B.J. Paterson, N. Bettez, L. Deegan, W.J. O'Brien, G.W. Kling, G.W. Kipphut, W.B. Bowden and A.E. Hershey, 1999. Impact of global change on the biogeochemistry and ecology of an Arctic freshwater system. *Polar Research*, 18:207-214.
- Hobbs, P.V. and A.L. Rangno, 1998. Microstructures of low and middle-level clouds over the Beaufort Sea. *Quarterly Journal of the Royal Meteorological Society*, 124:2035-2071.
- Hoff, R.M., 1988. Vertical structure of Arctic haze observed by Lidar. *Journal of Applied Meteorology*, 27:125-139.
- Hoff, R.M., W.R. Leaitch, P. Fellin and L.A. Barrie, 1983. Mass-size distributions of chemical constituents of the winter Arctic aerosol. *Journal of Geophysical Research*, 88:10947-10956.
- Høgda, K.A., H. Tømmervik, I. Solheim and Ø. Marhaug, 1995. Use of multitemporal Landsat image data for mapping the effects from air pollution in the Kirkenes-Pechenga area in the period 1973-1994. University of Tromsø, Foundation for applied Research, NORUT-IT Report, IT2039/1-95.
- Holopainen, T., H. Heinonen-Tanski and A. Halonen, 1996. Injuries to Scots pine mycorrhizas and chemical gradients in forest soil in the environment of a pulp mill in Central Finland. *Water, Air and Soil Pollution*, 87:111-130.
- Holt, C. and N.D. Yan, 2003. Recovery of crustacean zooplankton communities from acidification in Killarney Park, Ontario, 1971-2000: pH 6 as a recovery goal. *Ambio*, 32:203-207.
- Hopper, J.E., D.E.J. Worthy, L.A. Barrie and N.B.A. Trivett, 1994. Atmospheric observations of aerosol black carbon, carbon dioxide and methane in the High Arctic. *Atmospheric Environment*, 28:3047-3054.
- Hörnfeldt, B., 2004. Long-term decline in number of cyclic voles in boreal Sweden: analysis and presentation of hypotheses. *Oikos*, 107:376-392.
- Hughes, P.R., 1988. Insect populations on host plants subjected to air pollution. In: E.A. Heinrichs (ed.). *Plant Stress-Insect Interactions*, pp. 249-319. J. Wiley and Sons.
- Hurrell, J.W. and H. van Loon, 1997. Decadal variations in climate associated with the North Atlantic Oscillation. *Climatic Change*, 36:301-326.
- Ilyashuk, B.P. and E.A. Ilyashuk, 2001. Response of alpine chironomid communities (Lake Chuna, Kola Peninsula, northwest Russia) to atmospheric contamination. *Journal of Paleolimnology*, 25:467-475.
- Ilyashuk, B.P., E.A. Ilyashuk and V.A. Dauvalter, 2003. Chironomid responses to long-term metal contamination: a paleolimnological study in two bays of Lake Imandra, Kola Peninsula, northern Russia. *Journal of Paleolimnology*, 30:217-230.
- Isaeva, L., 2004. Influence of air pollution on species diversity of aphylloporaceous fungi. In: V. Kulikov, V. Razumovskij, O. Chervyakov and O. Kislova (eds.). *Man and environment in the Barents region at the threshold of the XXI century*, pp. 180-182. International Conference Proceedings, Aug. 6-11, 2001. Karelian Research Centre, Russian Academy of Sciences, Petrozavodsk. (In Russian)
- Isaksson, E., M. Hermanson, S. Hicks, M. Igarashi, K. Kamiyama, J. Moore, H. Motoyama, D. Muir, V. Pohjola, R. Vaikmae, R.S.W. van de Wal and O. Watanabe, 2003. Ice cores from Svalbard – useful archives of past climate and pollution history. *Physics and Chemistry of the Earth*, 28:1217-1228.
- Ivarsson, H. and M. Jansson, 1995. Sources of acidity in running waters in Central-Northern Sweden. *Water, Air and Soil Pollution*, 84:233-251.
- Iversen, T., 1984. On the atmospheric transport of pollution to the Arctic. *Geophysical Research Letters*, 11:457-460.
- Iversen, T. and E. Joranger, 1985. Arctic air pollution and large scale atmospheric flows. *Atmospheric Environment*, 19:2099-2108.
- Jackson, T.A., 1997. Long range atmospheric transport of mercury in ecosystems, and the importance of anthropogenic emissions: A critical review and evaluation of the published evidence. *Environmental Reviews*, 5:99-120.
- Jaffe, D., T. Iversen and G. Shaw, 1995. Comment on “A long term decrease in arctic haze at Barrow, Alaska” by B.A. Bodhaine and E.G. Dutton. *Geophysical Research Letters*, 22:739-740.
- James, P., A. Stohl, C. Forster, S. Eckhardt, P. Seibert and A. Frank, 2003. A 15-year climatology of stratosphere-troposphere exchange with a Lagrangian particle dispersion model: 2. Mean climate and seasonal variability. *Journal of Geophysical Research*, 108:8522, doi:10.1029/2002JD002639.
- Jeffrey, S.K., S.A. Norton, T.A. Haines, E.A. Rochette, R.H. Heath and S.C. Nodvin, 1992. Mechanisms of episodic acidification in low-order streams in Maine, USA. *Environmental Pollution*, 78:37-44.
- Jeffries, D.S., 1995. A preliminary assessment of nitrogen-based fresh water acidification in southeastern Canada. *Water, Air and Soil Pollution*, 85:433-438.
- Jeffries, D.S. (ed.), 1997. *Canadian acid rain assessment, Vol. 3: The Effects on Canada's Lakes, Rivers and Wetlands*. Environment, Ottawa, Ontario. 178p.
- Jeffries, D.S., T.A. Clair, S. Couture, P.J. Dillon, J. Dupont, W. Keller, D.K. McNicol, M.A. Turner, R. Vet and R. Weeber, 2003. Assessing the recovery of lakes in Southeastern Canada from the effects of acid deposition. *Ambio*, 32:176-182.
- Jenkins, A., L. Camarero, B.J. Cosby, R. Ferrier, M. Forsius, R. Helliwell, J. Kopacek, V. Majer, F. Moldan, M. Posch, M. Rogara, W. Schöpp and R.F. Wright, 2003. A modelling assessment of acidification and recovery of European surface waters. *Hydrology and Earth System Sciences*, 7:447-455.
- Jerre, J., 1994. Mapping terrestrial effects in the border area. Summary of activities and results. In: B. Sivertsen (ed.). *Air pollution problems in the northern region of Fennoscandia included Kola*, pp. 151-156. Proceedings from the Seminar at Svanvik, Norway, 1-3 June 1993.
- Jones, V.J. and H.J.B. Birks, 2004. Lake-sediment records of recent environmental change on Svalbard: results of diatom analysis. *Journal of Paleolimnology*, 31:445-466.
- Joynt, E.H. and A.P. Wolfe, 2001. Paleoenvironmental inference models from sediment diatom assemblages in Baffin Island lakes (Nunavut, Canada) and reconstruction of summer water temperature. *Canadian Journal of Fisheries and Aquatic Sciences*, 58:1222-1243.
- Julin, A.C., 2003. Atmospheric transport of nitrate to Svalbard, a trajec-



- tory and model study. Master thesis, University of Stockholm. (In Swedish).
- Kähkönen, A.-M., 1996. Soil geochemistry in relation to water chemistry and sensitivity to acid deposition in Finnish Lapland. *Water, Air and Soil Pollution*, 87:311-327.
- Kahl, J.D., 1990. Characteristics of the low-level temperature inversion along the Alaskan Arctic coast, *International Journal of Climatology*, 10:537-548.
- Kahl, J.D. and A.D.A. Hansen, 1989. Determination of regional sources of aerosol black carbon in the Arctic. *Geophysical Research Letters*, 16:327-330.
- Kalabin, G., 1991. Ecological situation in the Pechenga Region. Report INEP, Kola Science Center, Apatity, Russia.
- Kalela, O., 1957. Regulation of reproduction rate in sub-arctic populations of the vole *Clethrionomys rufocanus* (Sund.). *Annales Academiae Scientiarum Fennicae, Series A, IV Biologica*, 34:1-60.
- Kämäri, J. (ed.), 1998. Acidifying Pollutants, Arctic Haze, and Acidification in the Arctic. In: AMAP Assessment Report: Arctic Pollution Issues, Chapter 9, pp. 621-659. Arctic Monitoring and Assessment Programme, Oslo.
- Kapitsa, A.P. and E.I. Golubeva, 1997. The structure of tundra plant cover as an ecological indicator in the Kola Peninsula. In: R.M.M. Crawford (ed.). *Disturbance and Recovery in Arctic Lands*. NATO ASI Series, Kluwer Academic Publishers, 620p.
- Kaplan, J.O., N.H. Bigelow, I.C. Prentice, S.P. Harrison, P.J. Bartlein, T.R. Christensen, W. Cramer, N.V. Matveyeva, A.D. McGuire, D. F. Murray, V.Y. Razzhivin, B. Smith, D.A. Walker, P.M. Anderson, A.A. Andreev, L.B. Brubaker, M.E. Edwards and A.V. Lozhkin, 2003. Climate change and Arctic ecosystems: 1. Modeling, paleodata-model comparisons, and future projections. *Journal of Geophysical Research*, 108(D19):8171, doi:10.1029/2002JD002559.
- Karaban, R.T., I.M. Nazarov and N.P. Pavlinov, 1988. Principles of the environmental pollution and plant state integrated impact monitoring. *Inst. Appl. Geophys. Proc.* 72:4-10. (In Russian)
- Kashulina, G.M., 2002. Anthropogenic transformation of soils in subarctic European region. Kola Science Centre, Apatity, Russia. Part 1, 158 pp., Part 2, 234 pp. (In Russian)
- Kashulina, G. and V. Gregorauskiene, 2000. Complete soil profiles – technical report from the catchment stage of BARENTS Ecogeochemistry project. Geological Survey of Finland, Report No. S/41/0000/11/2000.
- Kashulina, G. and C. Reimann, 2001. Sulphur in the Arctic environment (1): results of a catchment-based multi-medium study. *Environmental Pollution*, 114:3-19.
- Kashulina, G. and C. Reimann, 2002. Sulphur in the Arctic environment (2): results of multi-medium regional mapping. *Environmental Pollution*, 116:337-350.
- Kashulina, G., C. Reimann, T.E. Finne, J.H. Halleraker, M. Äyräs and V.A. Chekushin, 1997. The state of ecosystems in the central Barents Region: scale, factors and mechanism of disturbance. *Science of the Total Environment*, 206:203-225.
- Kashulina, G., C. Reimann, T.E. Finne, P. de Caritat and H. Niskavaara, 1998a. Factors influencing NO<sub>3</sub> concentrations in rain, stream water, ground water and podzol profiles of eight small catchments in the European Arctic. *Environmental Pollution*, 102:559-568.
- Kashulina, G., M.-L. Räisänen, C. Reimann, P. de Caritat and I. Bogatyrev, 1998b. Acidity status and mobility of Al in podzols near SO<sub>2</sub> emission sources on the Kola Peninsula, NW Russia. *Applied Geochemistry*, 13/3:391-402.
- Kashulina, G., C. Reimann and D. Banks, 2003. Sulphur in the Arctic environment (3). environmental impact. *Environmental Pollution*, 124:151-171.
- Kasischke, E.S., E.J. Hyer, P.C. Novelli, L.P. Bruhwiler, N.H.F. French, A.I. Sukhinin, J.H. Hewson and B.J. Stocks, 2005. Influences of boreal fire emissions on Northern Hemisphere atmospheric carbon and carbon monoxide. *Global Biogeochemical Cycles*, 19, GB1012, doi:10.1029/2004GB002300.
- Kaste, Ø. and B.L. Skjelkvåle, 2002. Nitrogen dynamics in runoff from two small heathland catchments representing opposite extremes with respect to climate and N deposition in Norway. *Hydrology and Earth System Sciences*, 6:351-362.
- Kataev, G.D., 1995. Small mammals. In: W.W. Sytchev (ed.). *Forest ecosystems of Kola Peninsula under atmospheric pollution influence of smelters*, pp. 102-110. Russian Academy of Sciences. St. Petersburg, Russia. (In Russian)
- Kataev, G.D. and O.A. Makarova, 1984. Changes in fauna of terrestrial vertebrates of the Lapland reserve during a half of century. In: V.V. Kryuchkov, (ed.). *Monitoring of natural environment of the Kola North*, pp. 75-92. Kola Science Centre, Apatity.
- Kataev, G.D. and M.F. Popova, 1993. Cytological and morpho-physiological characteristics of voles from Monchegorsk area. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 379-382. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Kataev, G.D., J. Suomela and P. Palokangas, 1994. Densities of microtine rodents along a pollution gradient from a copper-nickel smelter. *Oecologia*, 97:491-498.
- Kattsov, V.M. and E. Källén, 2005. Chapter 4. Future climate change: modeling and scenarios for the Arctic. In: *Arctic Climate Impact Assessment*, pp. 99-150. Cambridge University Press.
- Kawamura, K., H. Kasukabe and L.A. Barrie, 1996. Source and reaction pathways of dicarboxylic acids, ketoacids, and dicarbonyls in Arctic aerosols: One year of observations. *Atmospheric Environment*, 30:1709-1722.
- Kekonen, T., J.C. Moore, R. Mulvaney, E. Isaksson, V. Pohjola and R. S.W. v.d Wal, 2002. An 800 year record of nitrate from the Lomonosovfonna ice core, Svalbard. *Annals of Glaciology*, 35:261-265.
- Keller, W., N.D. Yan, K.M. Somers and H. Heneberry, 2002. Crustacean zooplankton communities in lakes recovering from acidification. *Canadian Journal of Fisheries and Aquatic Science*, 59:726-735.
- Kellogg, W.W., 1995. Contaminants affecting the Arctic climate, and the role of the oceans. *Science of the Total Environment*, 160/161:769-775.
- Kim, Y., H. Hatsushika, R.R. Muskett and K. Yamazaki, 2005. Possible effect of boreal wildfire soot on Arctic sea ice and Alaska glaciers. *Atmospheric Environment*, 39:3513-3520.
- Kinnunen, K., 1992. Acidification of waters in Finnish Lapland. In: K. Kinnunen, M. Varmola (eds.). *Effect of Air and Acidification in Combination with Climatic Factors on Forest, Soils and Waters in Northern Fennoscandia*, pp. 72-78. Nordic Council of Ministers.
- Kirtsideli, I.J., N.I. Vorobeve and O.M. Tereshenkova, 1995. Influence of industrial pollution on the communities of soil micromycetes of forest tundra in Taimyr Peninsula. *Mikologiya i Fitopatologiya (Mycology and phytopathology)*, 29:12-19.
- Kjellström, T.E., A. Neller and R.W. Simpson, 2002. Air pollution and its health impacts: the changing panorama. *Medical Journal of Australia*, 177:604-608.
- Klonecki, A., P. Hess, L. Emmons, L. Smith, J. Orlando and D. Blake, 2003. Seasonal changes in the transport of pollutants into the Arctic troposphere- model study. *Journal of Geophysical Research*, 108:8367, doi:10.1029/2002JD002199.
- Koch, D. and J. Hansen, 2005. Distant origins of Arctic black carbon: A Goddard Institute for Space Studies ModelE experiment. *Journal of Geophysical Research*, 100, D04204, doi:10.1029/2004JD005296.
- Koneva, G.G., 1993. Changes in soil macrofauna around "Severonikel" smelter complex. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 362-364. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Koneva, G.G. and S. Koponen, 1993. Density of ground-living spiders (Araneae) near smelter in Kola Peninsula. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 365. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Koponen, S., 2005. Spiders (Araneae) along a pollution gradient. In: C. Deltshev (ed.). *22nd European Colloquium of Arachnology*, 1-6 August 2005, Blagoevgrad, Bulgaria, p. 36. Institute of Zoology & National Museum of Natural History, Bulgarian Academy of Sciences.
- Koptsik, G. and S. Koptsik, 1995. Critical loads of acid deposition for forest ecosystems in the Kola Peninsula. *Water, Air and Soil Pollution*, 85:2553-2558.
- Koptsik, G. and I. Muchina, 1995. Effect of acid deposition on acidity and exchangeable cations in the podzols of Kola Peninsula. *Water, Air and Soil Pollution*, 85:1209-1214.
- Koptsik, G.N., S.V. Koptsik and D. Aamlid, 1999a. Transformation of the elemental composition of plants in northern taiga biogeocoenoses under air pollution impact. *Moscow University Soil Science Bulletin*, 54:39-51.
- Koptsik, G.N., S.V. Koptsik, K. Venn, D. Aamlid, L. Strand and M.A. Zhuravleva, 1999b. Variations in acidity and cation exchange properties of forest soils under impact of acid atmospheric precipitation. *Eurasian Soil Science*, 32:787-798.
- Koptsik, G.N., S.V. Koptsik and D. Aamlid, 2001. Pine needle chemistry near a large point SO<sub>2</sub> source in northern Fennoscandia. *Water, Air and Soil Pollution*, 130:929-934.
- Koptsik, S., G. Koptsik, S. Livantsova, L. Eruslankina, T. Zhmelkova and Zh. Vologdina, 2003. Heavy metals in soils near the nickel smelter: chemistry, spatial variation, and impacts on plant diversity. *Journal of Environmental Monitoring*, 5:441-450.
- Korhola, A. and J. Weckström, 2004. Paleolimnological studies in Arctic Fennoscandia and the Kola Peninsula (Russia). In: R. Pienitz, M. Douglas and J.P. Smol (eds.). *Long-term environmental change in Arctic and Antarctic lakes*, pp. 381-418. Kluwer Academic Publishers.
- Korhola, A., J. Weckström and M. Nyman, 1999. Predicting the long-term acidification trends in small subarctic lakes using diatoms. *Journal of Applied Ecology*, 36:1021-1034.
- Korhola, A., J. Weckström and T. Blom, 2002. Relationships between



- lake and land-cover features along latitudinal vegetation ecotones in arctic Fennoscandia. *Archive für Hydrobiologie Supplement*, 139/2, 203-235.
- Korsman, T., 1999. Temporal and spatial trends of lake acidity in northern Sweden. *Journal of Paleolimnology*, 22:1-15.
- Kortelainen, P., 1993. Content of total organic carbon in Finnish lakes and its relationship to catchment characteristics. *Canadian Journal of Fisheries and Aquatic Sciences*, 50:1477-1483.
- Kozlov, M.V., 1987. Responses of Insect Populations to Human Impact. Institute of Forest and Timber. Krasnoyarsk (in Russian, English summary).
- Kozlov, M.V., 1996. Subalpine and alpine assemblages of Lepidoptera in the surroundings of a powerful smelter on the Kola Peninsula, NW Russia. *Nota Lepidopterologica*, 18:17-37.
- Kozlov, M.V., 1997. Pollution impact on insect biodiversity in boreal forests: evaluation of effects and perspectives of recovery. In: R.M.M. Crawford (ed.). *Disturbance and Recovery in Arctic Lands: An Ecological Perspective*, pp. 213-250. Proceedings of the NATO Advanced Research Workshop on Disturbance and Recovery of Arctic Terrestrial Ecosystems, Rovaniemi, Finland, 24-30 September 1995. NATO ASI series. Partnership subseries 2. Environment: vol. 25. Kluwer Academic Publishers.
- Kozlov, M.V., 2001. Snowpack changes around a nickel-copper smelter at Monchegorsk, northwestern Russia. *Canadian Journal of Forest Research*, 31:1684-1690.
- Kozlov, M.V., 2002. Changes in wind regime around a nickel-copper smelter at Monchegorsk, northwestern Russia. *International Journal of Biometeorology*, 46:76-80.
- Kozlov, M.V., 2003. Density fluctuations of the leafminer *Phyllonorycter strigulatella* (Lepidoptera: Gracillariidae) in the impact zone of a power plant. *Environmental Pollution*, 121:1-10.
- Kozlov, M., 2005. Pollution resistance of mountain birch, *Betula pubescens* subsp. *czerepanovii*, near the copper-nickel smelter: natural selection or phenotypic acclimation? *Chemosphere*, 59:189-197.
- Kozlov, M.V. and V.S. Barcan, 2000. Environmental contamination in the central part of the Kola Peninsula: history, documentation, and perception. *Ambio*, 29:512-517.
- Kozlov, M.V. and E. Haukioja, 1997. Microclimate changes along a strong pollution gradient in northern boreal forest zone. In: J.L. Uso, C.A. Brebbia and H. Power, (eds.). *Ecosystems and Sustainable development*, pp. 603-614. *Advances in Ecological Sciences*, vol. 1.
- Kozlov, M.V. and J. Jalava, 1994. Lepidoptera of Kola Peninsula, Northwestern Russia. *Entomologica Fennica*, 5:65-85.
- Kozlov, M.V. and A.V. Selikhovkin, 1997. No effects of sulphur dioxide on larval performance of *Epirrita autumnata* (Lepidoptera: Geometridae). *Environmental Entomology*, 26:1361-1363.
- Kozlov, M.V. and T. Whitworth, 2002. Population densities and diversity of Calliphoridae (Diptera) around a nickel-copper smelter at Monchegorsk, Northwestern Russia. *Entomologica Fennica*, 13:98-104.
- Kozlov, M.V. and E.L. Zvereva, 2004. Reproduction of mountain birch along a strong pollution gradient near Monchegorsk, Northwestern Russia. *Environmental Pollution*, 132:443-451.
- Kozlov, M.V., J. Jalava, A.L. Lvovsky and K. Mikkola, 1996a. Population densities and diversity of Noctuidae (Lepidoptera) along an air pollution gradient on the Kola Peninsula, Russia. *Entomologica Fennica*, 7:9-15.
- Kozlov, M.V., A.L. Lvovsky and K. Mikkola, 1996b. Abundance of day-flying Lepidoptera along an air pollution gradient in the northern boreal forest zone. *Entomologica Fennica*, 7:137-144.
- Kozlov, M.V., E.L. Zvereva and A.V. Selikhovkin, 1996c. Decreased performance of *Melasoma lapponica* (Coleoptera: Chrysomelidae) fumigated by sulphur dioxide: direct toxicity versus host plant quality. *Environmental Entomology*, 25:143-146.
- Kozlov, M.V., E. Haukioja and E.F. Kovnatsky, 2000. Uptake and excretion of nickel and copper by leaf-mining larvae of *Eriocrania semipurpurella* (Lepidoptera: Eriocraniidae) feeding on contaminated birch foliage. *Environmental Pollution*, 108:303-310.
- Kozlov, M.V., E.L. Zvereva, A.S. Gilyazov and G.D. Kataev, 2005a. Contaminated zone around a nickel-copper smelter: a death trap for birds and mammals? In: A.R. Burk (ed.). *Trends in Biodiversity*, pp. 81-99. Nova Science Publishers.
- Kozlov, M.V., N.K. Brodskaya, A. Haarto, K. Kuusela, M. Schäfer and V. Zverev, 2005b. Population densities and diversity of blood-sucking flies (Diptera: Ceratopogonidae, Culicidae, Tabanidae, Simuliidae) around a nickel-copper smelter at Monchegorsk, North western Russia. *Journal of Vector Ecology*, 30:263-271.
- Kuylenstierna, J.C.I., H. Rodhe, S. Cinderby and K. Hicks, 2001. Acidification in developing countries: Ecosystem sensitivity and the critical load approach on a global scale. *Ambio*, 30:20-28.
- Lappalainen, A., O. Mähönen, J. Erkinaro, M. Rask and E. Niemelä, 1995. Acid deposition from the Russian Kola Peninsula: are sensitive fish populations in north-eastern Finnish Lapland affected? *Water, Air and Soil Pollution*, 85:439-444.
- Laudon, H. and K.H. Bishop, 1999. Quantifying sources of ANC depression during spring flood episodes in Northern Sweden. *Environmental Pollution*, 105:427-455.
- Laudon, H. and K.H. Bishop, 2002. The rapid and extensive recovery from episodic acidification in northern Sweden due to declines in  $\text{SO}_4^{2-}$  deposition. *Geophysical Research Letters*, 29:1594-1597.
- Laudon, H. and H.F. Hemond, 2002. Recovery of streams from episodic acidification in northern Sweden. *Environmental Science and Technology*, 36:921-928.
- Laudon, H., S. Köhler and K. Bishop, 1999. Natural acidity or anthropogenic acidification in the spring flood of northern Sweden? *Science of the Total Environment*, 234:63-73.
- Laudon, H., O. Westling and K. Bishop, 2000. Cause of pH decline in stream water during spring melt runoff in Northern Sweden. *Canadian Journal of Fisheries and Aquatic Sciences*, 57:1888-1900.
- Laudon, H., O. Westling, A. Bergkvist and K. Bishop, 2004. Episodic acidification in northern Sweden: a regional assessment of the anthropogenic component. *Journal of Hydrology*, 297:162-173.
- Lavoué, D., 2000. Transport to the Arctic region of carbon aerosols emitted by biomass burning in boreal and temperate regions (In French: Transport vers la région arctique de l'aérosol carboné émis par les feux de biomasse des régions boreale et tempérée). PhD thesis at the University of Paris VII.
- Lavoué, D., C. Lioussé, H. Cachier, B.J. Stocks and J.G. Goldammer, 2000. Modeling of carbonaceous particles emitted by boreal and temperate wildfires at northern latitudes. *Journal of Geophysical Research*, 105:26871-26890.
- Leaith, W.R., R.M. Hoff, S. Melnichuk and A. Hogan, 1984. Some physical and chemical properties of the Arctic winter aerosol in northeastern Canada. *Journal of Climate and Applied Meteorology*, 23:916-928.
- Leaith, W.R., R.M. Hoff and J.L. MacPherson, 1989. Airborne and Lidar measurements of aerosol and cloud particles in the troposphere over Alert Canada in April 1986. *Journal of Atmospheric Chemistry*, 9:187-212.
- Lebedeva, E.V., 1993. Soil micromycetes of the non-ferrous metallurgy enterprise environs on the Kola Peninsula. *Mikologiya i Fitopatologiya* (Mycology and phytopathology), 27:12-17. (In Russian)
- Lee Behrens, H., S. Øftedal and P.W. Schive, 2000. PAME – Snap Shot Analysis of Maritime Activities in the Arctic. Revision No. 01, REPORT NO. 2000-3220. Norwegian Maritime Directorate, 75p.
- Lehtonen, J., 1987. Recovery and development of birch forests damaged by *Epirrita autumnata* in Utsjoki area, North Finland. Reports from the Kevo Subarctic Research Station, 20:35-39.
- Leighton, H., 1983. Influence of the Arctic haze on the solar radiation budget. *Atmospheric Environment*, 17:2065-2068.
- Leiterer, U., D. Nagel and R. Stolte, 1997. Typical vertical profiles of aerosol spectral extinction coefficients derived from observations of direct solar radiation extinction during the aircraft experiments Arctic Haze 94/95 and Merisec 93/94. *Atmospheric Research*, 44:73-88.
- Levina, V.I., 1969. Seasonal dynamic of humidity and chemical properties of podzolic, mountain-podzolic and mountain-tundra soils in Murmansk region. In: *Soils regimes on the Polar North*. Leningrad: "Nauka", pp. 5-58. (In Russian)
- Levine, S.Z. and S.E. Schwarz, 1982. In-cloud and below-cloud scavenging of nitric acid vapor. *Atmospheric Environment*, 16:1725-1734.
- Li, S.M. and L.A. Barrie, 1993. Biogenic sulphur aerosols in the Arctic troposphere. 1. Contributions to sulphate. *Journal of Geophysical Research*, 98D:20613-20622.
- Lidmar-Bergström, K. and J.-O. Näslund, 2005. Major Landforms and Bedrock. In: M. Seppälä (ed.). *The Physical Geography of Fennoscandia*, pp. 3-15. Oxford Regional Environments Series. Oxford University Press.
- Lien, L., A. Henriksen and T.S. Traaen, 1995. Critical loads of acidity to surface waters: Svalbard. *Science of the Total Environment*, 160/161:703-713.
- Lobersli, E. and K. Venn (eds.), 1995. Effects of air pollutants on terrestrial ecosystems in the border area between Norway and Russia. DN-utredning 1995-8 Directorate for Nature Management, Trondheim, 140p.
- Lockhart, W.L., P. Wilkinson, B.N. Billeck, R.V. Hunt, R. Wagemann and G.J. Brunskill, 1995. Current and historical inputs of mercury to high-latitude lakes in Canada and to Hudson Bay. *Water, Air and Soil Pollution*, 80:603-610.
- Lockhart, W.L., P. Wilkinson, B.N. Billeck, R.A. Danell, R.V. Hunt, G.J. Brunskill, J. Delaronde and V. St. Louis, 1998. Fluxes of mercury to lake sediments in central and northern Canada inferred from dated sediment cores. *Biogeochemistry*, 40:163-173.
- Löfgren, O., 1995. Niche expansion and increased maturation rate of *C. glareolus* in the absence of competitors. *Journal of Mammalogy*, 76:1100-1112.
- Logan, J.A., 1983. Nitrogen oxides in the troposphere; global and regional budgets. *Journal of Geophysical Research*, 88:10785-10807.
- Lukina, N. and V. Nikonov, 1996. Biogeochemical cycles in the northern

- forest ecosystems subjected to air pollution. Kola Science Centre. RAS. Part 1, 195 pp. Part 2, 216p.
- Luyssaert, S., M. Sulkava, H. Raitio and J. Hollmén, 2005. Are N and S deposition altering the mineral composition of Norway spruce and Scots pine needles in Finland? *Environmental Pollution*, 138:5-17.
- MacCracken, M.C., R.D. Cess and G. L. Potter, 1986. Climatic effects of anthropogenic Arctic aerosols: An illustration of climatic feedback mechanisms with one- and two-dimensional climate models. *Journal of Geophysical Research*, 91:14445-14450.
- Macdonald, R.W., B.L.A., T.F. Bidleman, M.L. Diamond, D.J. Gregor, R.G. Semkin, W.M.J. Strachan, Y.F. Li, F. Wania, M. Alae, L.B. Alexeeva, S.M. Backus, R. Bailey, J.M. Bewers, C. Gobeil, C.J. Halsall, T. Harner, J.T. Hoff, L.M.M. Jantunen, W.L. Lockhart, D. Mackay, D.C.G. Muir, J. Pudykiewicz, K.J. Reimer, J.N. Smith, G.A. Stern, W.H. Schroeder, R. Wagemann and M.B. Yunker, 2000. Contaminants in the Canadian Arctic: 5 years of progress in understanding sources, occurrence and pathways. *Science of the Total Environment*, 254:93-234.
- Macdonald, R.W., T. Harner, J. Fyfe, H. Loeng and T. Weingartner, 2003. AMAP Assessment 2002: The influence of global change on contaminant pathways to, within, and from the Arctic. Arctic Monitoring and Assessment Programme, Oslo, xii+65p.
- Macdonald, R.W., T. Harner and J. Fyfe, 2005. Recent climate change in the Arctic and its impact on contaminant pathways and interpretation of temporal trend data. *Science of the Total Environment*, 342:5-86.
- Mallory, M.L., A.J. Fontaine, P.A. Smith, M.O. W. Robertson and H.G. Gilchrist, 2006. Water chemistry of ponds on Southampton Island, Nunavut, Canada: effects of habitat and ornithogenic inputs. *Archiv für Hydrobiologie*, accepted.
- Manninen, S., 1995. Assessing the Critical Level of SO<sub>2</sub> for Scots Pine (*Pinus sylvestris* L.) in Northern Europe on the Basis of Needle Sulphur Fractions, Sulphur/Nitrogen Ratios and Needle Damage. PhD Thesis, University of Oulu, Finland.
- Manninen, S. and S. Huttunen, 2000. Response of needle sulphur and nitrogen concentrations of Scots pine versus Norway spruce to SO<sub>2</sub> and NO<sub>2</sub>. *Environmental Pollution*, 107:421-436.
- Manninen, S., S. Huttunen and M. Kontio, 1997a. Accumulation of sulphur in and on Scots pine needles in the subarctic. *Water, Air and Soil Pollution*, 95:147-164.
- Manninen, S., S. Huttunen and P. Perämäki, 1997b. Needle S fractions and S to N ratios as indices of SO<sub>2</sub> deposition. *Water, Air and Soil Pollution*, 95:277-297.
- Manninen, S., S. Huttunen and P. Perämäki, 1998. Effect of ambient SO<sub>2</sub> levels on S fractions in *Pinus sylvestris* foliage growing in the subarctic. *Scandinavian Journal of Forest Research*, 13:306-316.
- Masclat, P., V. Hoyau, J.L. Jaffrezou and H. Cachier, 2000. Polycyclic aromatic hydrocarbon deposition on the ice sheet of Greenland. Part 1: Superficial snow. *Atmospheric Environment*, 34:3195-3207.
- McBean, G., 2005. Chapter 2. Arctic climate – past and present. *In: Arctic Climate Impact Assessment*, pp. 21-60. Cambridge University Press.
- McGowan, J., P. Hider, E. Chacko and G. Town, 2000. Particulate air pollution and hospital admissions in Christchurch, New Zealand. *Australian and New Zealand Journal of Public Health*, 26:23-29.
- McMahon, T.A. and P.J. Denison, 1979. Empirical atmospheric deposition parameters – A survey. *Atmospheric Environment*, 13:571-585.
- McMichael, A.J., A. Haines, R. Sloof and S. Kovats, 1996. Climate Change and Human Health. World Health Organization.
- Meijer, S.N., W.A. Ockenden, A. Sweetman, K. Breivik, J.O. Grimalt and K.C. Jones, 2003. Global distribution and budget of PCBs and HCB in background surface soils: Implications for sources and environmental processes. *Environmental Science and Technology*, 37:667-672.
- Mendonça, B.G., J.J. DeLuise and J.A. Schroeder, 1981. Arctic Haze and perturbation in the solar radiation fluxes at Barrow, Alaska. *Proceedings from the 4th Conference on Atmospheric Radiation*. *Atm. Met. Soc.*, Toronto, pp. 95-96.
- Michelutti, N., T.E. Laing and J.P. Smol, 2001. Diatom assessment of past environmental changes in lakes located near the Noril'sk (Siberia) smelters. *Water, Air and Soil Pollution*, 125:231-241.
- Michelutti, N., M.S.V. Douglas, D.C.G. Muir, X. Wang and J.P. Smol, 2002a. Limnological characteristics of 38 lakes and ponds on Axel Heiberg Island, High Arctic Canada. *International Review of Hydrobiology*, 87:385-399.
- Michelutti, N., M.S.V. Douglas, D.R.S. Lean and J.P. Smol, 2002b. Physical and chemical limnology of 34 ultra-oligotrophic lakes and ponds near Wynniatt Bay, Victoria Island, Arctic Canada. *Hydrobiologia*, 482:1-13.
- Michelutti, N., M.S.V. Douglas and J.P. Smol, 2003. Diatom response to recent climatic change in a high arctic lake (Char Lake, Cornwallis Island, Nunavut). *Global and Planetary Change*, 38:257-271.
- Mikkola, K., 1996. A remote sensing analysis of vegetation damage around smelters in the Kola Peninsula, Russia. *International Journal of Remote Sensing*, 17(18):3675-3690.
- Miskimmin, B.M., J.W.M. Rudd and C.A. Kelly, 1992. Influence of dissolved organic carbon, pH, and microbial respiration rates on mercury methylation and demethylation in lake water. *Canadian Journal of Fisheries and Aquatic Sciences*, 49:17-22.
- Mitchell Jr., J.M., 1956. Visual range in the polar regions with particular reference to the Alaskan Arctic. *Journal of Atmospheric and Terrestrial Physics Special Supplement*, 1:195-211.
- Mitchell, M.J., D.J. Raynal and C.T. Driscoll, 1996. Biogeochemistry of a forested watershed in the central Adirondack mountains: temporal changes and mass balances. *Water, Air and Soil Pollution*, 88:355-369.
- Moberg, A., H. Tuomenvirta and P.Ø. Nordli, 2005. *In: M. Seppälä (ed.). The Physical Geography of Fennoscandia*, pp. 113-133. Oxford Regional Environments Series. Oxford University Press.
- Moiseenko, T.I., 1994. Acidification and critical loads in surface waters: Kola, northern Russia. *Ambio*, 23:418-424.
- Moiseenko, T.I., 1997. Theoretical bases of critical limit values of anthropogenic loads on subarctic surface waters. *Kola Science Center, Russian Academy of Science, Apatity*. 261p. (In Russian)
- Moiseenko, T.I., 1999. The fate of metals in Arctic surface waters. Method for defining critical levels. *Science of the Total Environment*, 236:19-39.
- Moiseenko, T.I., L.P. Kudryavtseva, I.V. Rodyushkin, V.A. Dauvalter, A.A. Lukin and N.A. Kashulin, 1995. Airborne contamination by heavy metals and aluminum in the freshwater ecosystems of the Kola Subarctic region (Russia). *Science of the Total Environment*, 160/161:715-727.
- Moiseenko, T.I., V.A. Dauvalter, B.P. Ilyashuk, L.Y. Kagan and E.A. Ilyashuk, 2000. Paleoeological reconstruction of the anthropogenic load. *Doklady Earth Sciences*, 307(1):102-105.
- Moiseenko, T.I., L. Kudryavtseva and I. Rodyushkin, 2001. The episodic acidification of small streams in the spring flood period of industrial polar region, Russia. *Chemosphere*, 42:45-50.
- Monteith, D.T., C.D. Evans and B. Reynolds, 2000. Are temporal variations in the nitrate content of UK upland freshwaters linked to the North Atlantic Oscillation? *Hydrological Processes*, 14:1745-1749.
- Moore, J.C., A. Grinsted, T. Kekonen and V. Pohjola, 2005. Separation of melting and environmental signals in an ice core with seasonal melt. *Geophysical Research Letters*, 32:L10501.
- Morgan, G., S. Corbett and J. Włodarczyk, 1998. Air pollution and hospital admissions in Sydney, Australia, 1990 to 1994. *American Journal of Public Health*, 88:1761-1766.
- Moser, K.A., J.P. Smol, D.R.S. Lean, G.M. MacDonald, 1998. Physical and chemical limnology of northern boreal lakes, Wood Buffalo National Park, northern Alberta and the Northwest Territories, Canada. *Hydrobiologia*, 377:25-43.
- Motova, A. and V.V. Nikonov, 1993. Aluminium compounds in podzolic Al-Fe-humus soils in the surroundings of the "Severonikel" smelter. *In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). Aerial pollution on Kola Peninsula*, pp.178-183. *Proceedings of the International Workshop*, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- MRCENR, 1995. Review of pollutant fallout in the atmosphere in the Murmansk region in 1993 and 1994. Murmansk Regional Committee for Ecology and Natural Resources, Murmansk. (In Russian)
- Muir, D.C.G., N.P. Grift, W.L. Lockhart, P. Wilkinson, B.N. Billeck and G.J. Brunskill, 1995. Spatial trends and historical profiles of organochlorine pesticides in Arctic lake sediments. *Science of the Total Environment*, 160/161:447-457.
- Murdoch, P.S., J.S. Baron and T.L. Miller, 2000. Potential effects of climate change on surface water quality in North America. *Journal of the American Water Resources Association*, 36:347-366.
- Mylona, S., 1996. Sulphur dioxide emissions in Europe 1880-1991 and their effect on sulphur concentrations and depositions. *Tellus*, B48:662-689.
- Naldrett, A.J., P.C. Lightfoot, V. Fedorenko, W. Doherty and N.S. Gorbachev, 1992. Geology and geochemistry of intrusions and flood basalts of the Noril'sk region USSR, with implications for the origin of the Ni-Cu ores. *Economic Geology*, 87:975-1004.
- Nash, III, T.H. and C. Gies, 1995. The response of lichens to atmospheric deposition with an emphasis on the Arctic. *Science of the Total Environment*, 160/161:737-747.
- Nelson, W.O. and P.G.C. Campbell, 1991. The effects of acidification on the geochemistry of Al, Cd, Pb and Hg in freshwater environments: A literature review. *Environmental Pollution*, 71:91-130.
- Newell, A.D. and M.E. Mitch, 1992. User's Guide and Data Dictionary for Kenai Lakes Investigation Project. EPA/600/R-92/070, U.S. Environmental Protection Agency, Washington, DC.
- Nikonov, V.V., N.V. Lukina, L.M. Polyanskaya and A.N. Panikova, 2001. Distribution of microorganisms in the Al-Fe-humus podzols of natural and anthropogenically impacted boreal spruce forests.



- Mikrobiologiya, 70:374-383. (In Russian)
- Nilsson, J. and P. Grennfelt (eds.), 1988. Critical loads for sulphur and nitrogen. Nord 1988:97, Nordic Council of Ministers, 418p.
- Noone, K.J. and A.D. Clarke, 1988. Soot scavenging measurements in Arctic snowfall. *Atmospheric Environment*, 22:2773-2779.
- Nordin, A., J. Strengbom, J. Witzell, T. Näsholm and L. Ericson, 2005. Nitrogen deposition and the biodiversity of boreal forests: implications for the nitrogen critical load. *Ambio*, 34:20-24.
- Norwegian Maritime Directorate, 2000. PAME – Snap Shot Analysis of Maritime Activities in the Arctic. Norwegian Maritime Directorate Report No. 2000-3220. 72p.
- Nøst, T., V. Yakovlev, H.M. Berger, N. Kashulin, A. Langeland, A. Lukin and H. Muladal, 1992. Impacts of pollution on freshwater communities in the border region between Russia and Norway. In: M. Varmola and T. Katermaa (eds.). Symposium on the state of the environment and environmental monitoring in Northern Fennoscandia and the Kola Peninsula, Oct. 6-8 1992, Rovaniemi, Extended Abstracts, pp. 152-155. Arctic Centre Publications 4.
- Novikov, G.A., 1981. Life on and under snow: Life of our birds and animals, Issue 3. Leningrad University. (In Russian)
- Nuorteva, P., S. Autio, K. Bergbom, H. Braunschweiler and E. Koski, 1987. Effect of acidification on mobility of heavy metals and aluminum in forest and lake ecosystems (Happamoitumisen vaikutus raskasmetallien ja alumiinin liikkuvuuteen metsä- ja järviökosysteemissa). *Aquilo Series Botanica* 25:122-128.
- Nyman, M., A. Korhola and S.J. Brooks, 2005. The distribution and diversity of Chironomidae (Insecta: Diptera) in western Finnish Lapland, with special emphasis on shallow lakes. *Global Ecology and Biogeography*, 14:137-153.
- Odland, J.O., E. Nieboer, N. Romanova and Y. Thomassen, 2004. Elements in placenta and pregnancy outcome in arctic and subarctic areas. *International Journal of Circumpolar Health*, 63:169-187.
- Ohtonen, R. and H. Väre, 1996. Importance of site homogeneity in gradient studies on soil ecology. II. Soil microbiology. In: H. Väre (ed.). Relationships between spatial heterogeneity in vegetation communities and in belowground ecosystems. *Acta Universitatis Ouluensis*, A 275, III:1-11.
- Olofsson, J., H. Kitti, P. Rautiainen, S. Stark and L. Oksanen, 2001. Effects of summer grazing by reindeer on composition of vegetation, productivity and nitrogen cycling. *Ecography*, 24:13-24.
- Orr, J.C., V.J. Fabry, O. Aumont, L. Bopp, S.C. Doney, R.A. Feely, A. Gnanadesikan, N. Gruber, A. Ishida, F. Joos, R.M. Key, K. Lindsay, E. Maier-Reimer, R. Matear, P. Monfray, A. Mouchet, R.G. Najjar, G.-K. Plattner, K.B. Rodgers, C.L. Sabine, J.L. Sarmiento, R. Schlitzer, R.D. Slater, I.J. Totterdell, M.-F. Weirig, Y. Yamanaka and A. Yool, 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, 437:681-686.
- Ottar, B., Y. Gotaas, O. Hov, T. Iversen, E. Joranger, M. Oehme, J. Pacyna, A. Semb, W. Thomas and V. Vitolas, 1986. Air pollutants in the Arctic. Final report of a research programme conducted on the behalf of British Petroleum, Ltd. Norwegian Institute of Air Research, NILU OR 30/86, 80p.
- Overpeck, J.K., K. Hughen, D. Hardy, R. Bradley, R. Case, M. Douglas, B. Finney, K. Gajewski, G. Jacoby, A. Jennings, S. Lamoureux, A. Lasca, G. MacDonald, J. Moore, M. Retelle, S. Smith, A. Wolfe and G. Zielinski, 1997. Arctic environmental change of the last four centuries. *Science*, 278:1251-1256.
- Pacyna, J.M. and B. Ottar, 1988. Vertical distribution of aerosols in the Norwegian Arctic. *Atmospheric Environment*, 22:2213-2223.
- Pacyna, E.G. and J.M. Pacyna, 2002. Global emission of mercury from anthropogenic sources in 1995. *Water, Air and Soil Pollution*, 137:149-165.
- Pacyna, J.M., V. Vitolis and J.E. Hanssen, 1984. Size differentiated composition of the Arctic aerosol at Ny Alesund Zeppelin. *Atmospheric Environment*, 18:2447-2459.
- Park, R.J., D.J. Jacob, P.I. Palmer, A.D. Clarke, R.J. Weber, M.A. Zondlo, F.L. Eisele, A.R. Bandy, D.C. Thornton, G.W. Sachse and T.C. Bond, 2005. Export efficiency of black carbon aerosol in continental outflow: Global implications. *Journal of Geophysical Research*, 110: D11205, doi:10.1029/2004JD005432.
- Penkett, S.A., N.J. Blake, P. Lightman, A.R.W. Marsh, P. Anwyl and G. Butcher, 1993. The seasonal variation of nonmethane hydrocarbons in the free troposphere over the North Atlantic Ocean: Possible evidence for extensive reaction of hydrocarbons with the nitrate radical. *Journal of Geophysical Research*, 98:2865-2885.
- Perez-Fuentetaja, A., P.J. Dillon, N.D. Yan and D.J. McQueen, 1999. Significance of dissolved organic carbon in the prediction of thermocline depth in small Canadian shield lakes. *Aquatic Ecology*, 33:127-133.
- Petroeschovsky, A., R.W. Simpson, L. Thalib and S. Rutherford, 2001. Associations between outdoor air pollution and hospital admissions in Brisbane, Australia. *Archives of Environmental Health*, 56:37-52.
- Pienitz, R. and W.F. Vincent, 2000. Effect of climate change relative to ozone depletion on UV exposure in subarctic lakes. *Nature*, 404:484-487.
- Pienitz, R., J.P. Smol and D.R.S. Lean, 1997a. Physical and chemical limnology of 59 lakes located between the southern Yukon and the Tuktoyaktuk Peninsula, Northwest Territories (Canada). *Canadian Journal of Fisheries and Aquatic Sciences*, 54:330-346.
- Pienitz, R., J.P. Smol and D.R.S. Lean, 1997b. Physical and chemical limnology of 24 lakes located between Yellowknife and Contwoyto Lake, Northwest Territories (Canada). *Canadian Journal of Fisheries and Aquatic Sciences*, 54:347-358.
- Poléo, A.B.S. and F. Bjerkely, 2000. Effect of unstable aluminium chemistry on Arctic char (*Salvelinus alpinus*). *Canadian Journal of Fisheries and Aquatic Sciences*, 57:1423-1433.
- Polyanskaya, L.M., V.V. Nikonov, N.V. Lukina, A.N. Panikova and D.G. Zvyagintsev, 2001. Microorganisms of Al-Fe-humus podzols of lichen pine forests as affected by aerotechnogenic pollution. *Pochovovedenie (Eurasian Soil Science)*, 2:215-226. (In Russian)
- Pope, C.A. III, M.J. Thun, M.M. Namboodiri, D.W. Dockery, J.S. Evans, F.E. Speizer and C.W. Heath, Jr., 1995. Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *American Journal of Respiratory and Critical Care Medicine*, 151:669-674.
- Pope, C.A., R.T. Burnett, M.J. Thun, E.E. Calle, D. Krewski, K. Ito and G.D. Thurston, 2002. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *The Journal of the American Medical Association*, 287:1132-1141.
- Porch, W.M. and M.C. MacCracken, 1982. Parametric study of the effects of Arctic soot on the solar radiation. *Atmospheric Environment*, 16:1365-1371.
- Psenner, R. and R. Schmidt, 1992. Climate-driven pH control of remote alpine lakes and effects of acid deposition. *Nature*, 356:781-783.
- Pueschel, R.F. and S.A. Kinne, 1995. Physical and radiative properties of Arctic atmospheric aerosols. *Science of the Total Environment*, 161:811-824.
- Quinn, P.K., T.S. Bates, T.L. Miller, D.J. Coffman, J.E. Johnson, J.M. Harris, J.A. Ogren, G. Forbes, T.L. Anderson, D.S. Covert and M.J. Rood, 2000. Surface submicron aerosol chemical composition: What fraction is not sulfate? *Journal of Geophysical Research*, 105:6785-6806.
- Quinn, P.K., T.L. Miller, T.S. Bates, J.A. Ogren, E. Andrews and G.E. Shaw, 2002. A three-year record of simultaneously measured aerosol chemical and optical properties at Barrow, Alaska. *Journal of Geophysical Research*, 107(D11). 10.1029/2001JD001248.
- Raatz, W.E. and G.E. Shaw, 1984. Long-range tropospheric transport of pollution aerosols into the Alaskan Arctic. *Journal of Climate and Applied Meteorology*, 23:1052-1064.
- Radke, F.S., J.H. Lyons, D.A. Hegg, P.V. Hobbs and I.H. Bailey, 1984. Airborne observations of arctic aerosols, I: Characteristics of arctic haze. *Geophysical Research Letters*, 11:393-396.
- Raguotis, A.D., 1989. Impact of industrial emissions on soil microflora in forests of extreme North. In: G.M. Kozubov (ed.). Ecology of northern forests: Abstracts of presentations of the 1st All-Union Meeting, 2-7 October 1989, vol. 2, pp 45-46. Institute of Biology, Komi Science Centre, Syktyvkar. (In Russian)
- Rahn, K.A., 1981. Relative importance of North America and Eurasia as sources of Arctic aerosol. *Atmospheric Environment*, 15:1447-1455.
- Rahn, K.A., 1989. Proceedings of the International Symposium on Arctic Air Chemistry. *Atmospheric Environment*, 23:2345-2347.
- Rahn, K.A. and R.J. McCaffrey, 1979. Long range transport of pollution aerosol to the Arctic. A problem without borders. Proceedings of the WMO symposium on the Long-range Transport of Pollutants and its relation to the General Circulation including Stratospheric/Tropospheric Exchange Processes, pp. 25-35, Sofia, 1-5 Oct 1979. WMO No. 538.
- Rahn, K.A., R. Borys and G.E. Shaw, 1977. The Asian source of Arctic Haze bands. *Nature*, 268:713-715.
- Rahn, K.A., R.D. Borys and G.E. Shaw, 1981. Asian desert dust over Alaska: Anatomy of an Arctic Haze Episode. In: T.L. Pewe (ed.). Desert dust: origin, characteristic and effect on man, pp. 37-70. Special paper No. 186. The Geological Society of America.
- Räsänen, M.L., G. Kashulina and I. Bogatyrev, 1997. Mobility and retention of heavy metals, arsenic and sulphur in podzols at eight locations in northern Finland and Norway and the western half of the Kola Peninsula. *Journal of Geochemical Exploration*, 59:175-195.
- Ramlal, P.S., C.A. Kelly, J.W.M. Rudd and A. Furutani, 1993. Sites of methyl mercury production in remote Canadian Shield lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, 50:972-979.
- Rao, T.N. and S. Kirkwood, 2005. Characteristics of tropopause folds over Arctic latitudes. *Journal of Geophysical Research*, 110:D18102, doi:10.1029/2004JD005374.
- Rask, M., M. Appelberg, T. Hesthagen, J. Tammi and A. Lappalainen, Fish Status Survey of Nordic Lakes – species composition, distribution, effects of environmental changes. Nordic Council of Ministers. TemaNord 2000:508.
- Rasmussen, S.O., K.K. Andersen, M.-L. Siggaard-Andersen and H.B.

- Clausen, 2002. Extracting the annual signal from Greenland ice-core chemistry and isotopic records *Annals of Glaciology*, 35:131-135.
- Rautio, P. and S. Huttunen, 2003. Total vs. internal element concentrations in Scots pine needles along a sulphur and metal pollution gradient. *Environmental Pollution*, 122:273-289.
- Rautio, M. and A. Korhola, 2002. Effects of ultraviolet radiation and dissolved organic carbon on the survival of subarctic zooplankton: Implications to climate change. *Polar Biology*, 25:460-468.
- Rautio, P., S. Huttunen and J. Lamppu, 1998a. Element concentrations in Scots pine needles on radial transects across subarctic area. *Water, Air and Soil Pollution*, 102:389-405.
- Rautio, P., S. Huttunen and J. Lamppu, 1998b. Seasonal foliar chemistry of northern Scots pines under sulphur and heavy metal pollution. *Chemosphere*, 37:271-287.
- Rautio, P., S. Huttunen, E. Kukkola, R. Peura and J. Lamppu, 1998c. Deposited particles, element concentrations and needle injuries on Scots pines along an industrial pollution transect in northern Europe. *Environmental Pollution*, 103:81-89.
- Reimann, C., P. de Caritat, J.H. Halleraker, T. Volden, M. Åyräs, H. Niskavaara, V.A. Chekushin and V.A. Pavlov, 1997. Rainwater composition in eight arctic catchments of Northern Europe (Finland, Norway and Russia). *Atmospheric Environment*, 31/2:159-170.
- Reimann, C., M. Åyräs, V. Chekushin, I. Bogatyrev, R. Boyd, P. de Caritat, R. Dutter, T.E. Finne, Ø. Jæger, G. Kashulina, O. Lehto, H. Niskavaara, V. Pavlov, M.L. Raisanen, T. Strand and T. Volden, 1998a. *Environmental Geochemical Atlas of the Central Barents Region*. Special Publication, Geological Survey of Norway, Trondheim, 745p.
- Reimann, C., P. de Caritat, T.E. Finne, G. Kashulina, H. Niskavaara and V. Pavlov, 1998b. Comparison of elements in O- and C-horizon soils from the surroundings of Nikel, Kola Peninsula, using different grain size fractions and extractions. *Geoderma* 84:65-87.
- Reimann, C., D. Banks and P. de Caritat, 2000a. Impacts of airborne contamination on regional soil and water quality: The Kola Peninsula, Russia. *Environmental Science and Technology*, 34:2727-2732.
- Reimann, C., D. Banks and G. Kashulina, 2000b. Processes influencing the chemical composition of the O-horizon of podzols along a 500-km north-south profile from the coast of the Barents Sea to the Arctic Circle. *Geoderma*, 95:113-139.
- Reimann, C., G. Kashulina, P. de Caritat and H. Niskavaara, 2001a. Multi-element, multi-medium regional geochemistry in the European Arctic: element concentrations, variation and correlation. *Applied Geochemistry*, 16:759-780.
- Reimann, C., F. Koller, G. Kashulina, H. Niskavaara and P. Englmaier, 2001b. Influence of extreme pollution on the inorganic chemical composition of some plants. *Environmental Pollution*, 115:239-252.
- Reimann, C., F. Koller, B. Frengstad, G. Kashulina, H. Niskavaara and P. Englmaier, 2001c. Comparison of the element composition in several plant species and their substrate from a 1 500 000 km<sup>2</sup> area in Northern Europe. *Science of the Total Environment*, 278:87-112.
- Reimann, C., H. Niskavaara, G. Kashulina, P. Filzmoser, R. Boyd, T. Volden, O. Tomilina and I. Bogatyrev, 2001d. Critical remarks on the use of terrestrial moss (*Hylocomium splendens* and *Pleurozium schreberi*) for monitoring airborne pollution. *Environmental Pollution*, 113:41-57.
- Reimann, C., F. Koller, B. Frengstad, G. Kashulina, H. Niskavaara and P. Englmaier, 2003. Total sulphur in leaves of several plant species from nine catchments within a 1 500 000 km<sup>2</sup> area in northern Europe: local vs. regional variability. *Geochemistry: Exploration, Environment, Analysis*, 3:205-215.
- Reuss, J.O. and D.W. Johnson, 1986. *Acid Deposition and the Acidification of Soils and Waters*. Ecological Studies. Springer Verlag, 119p.
- Reviews of the state of environment in Russian Federation in 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002. Moscow, Hydromeoizda. (In Russian)
- Ricard, V., J.-L. Jaffrezo, V.-M. Kerminen, R.E. Hillamo, M. Sillanpää, S. Ruellan, C. Lioussse and H. Cachier, 2002. Two years of continuous aerosol measurements in northern Finland. *Journal of Geophysical Research*, 107(D11), 10.1029/2001JD000952.
- Richter, V.A. and E.L. Zvereva, 1996. The tachinid species *Cleonece nitidiuscula* Zetterstedt new for fauna of Murmansk Province (Diptera: Tachinidae). *Zoosystematica Rossica*, 6:202.
- Riemer, J. and J.B. Whittaker, 1989. Air pollution and insect herbivores: observed interactions and possible mechanisms. In: E.A. Bernays (ed.). *Insect-Plant Interactions*, pp. 73-105. vol. 1. CRC Press.
- Rigina, O. and M. Kozlov, 1999. Pollution impact on sub-Arctic northern taiga forests in the Kola Peninsula, Russia. In: J.L. Innes and J. Oleksyn (eds.). *Forest Dynamics in Heavily Polluted Regions*, pp. 37-65. IUFRO Research Series, No. 1. CAB International.
- Ritter, C., J. Notholt, J. Fischer and C. Rathke, 2005. Direct thermal radiative forcing of tropospheric aerosol in the Arctic measured by ground based infrared spectrometry. *Geophysical Research Letters*, 32:L23816, doi:10.1029/2005GL024331.
- Rose, N.L., C.L. Rose, J.F. Boyle and P.G. Appleby, 2004. Lake-sediment evidence for local and remote sources of atmospherically deposited pollutants on Svalbard. *Journal of Paleolimnology*, 31:499-513.
- Rouse, W.R., M.S.W. Douglas, R.E. Hecky, A.E. Hershey, G.W. Kling, L. Lesack, P. Marsh, M. McDonald, B.J. Nicholson, N.T. Roulet and J.P. Smol, 1997. Effects of climate change on the freshwaters of arctic and subarctic North America. *Hydrological Processes*, 11:873-902.
- Rühland, K.M. and J.P. Smol, 1998. Limnological characteristics of 70 lakes spanning Arctic treeline from Coronation Gulf to Great Slave Lake in the Central Northwest Territories, Canada. *International Review of Hydrobiology*, 83:183-203.
- Rühland, K.M., J.P. Smol, X. Wang and D.C.G. Muir, 2003. Limnological characteristics of 56 lakes in the central Canadian Arctic tree-line region. *Journal of Limnology*, 62:9-27.
- Ruoho-Airola, T. and T. Salmi, 2001. Episodicity of sulphate deposition in Finland. *Water, Air and Soil Pollution*, 130:529-534.
- Ruoho-Airola, T., P. Anttila and T. Salmi, 2004. Airborne sulphur and nitrogen in Finland – trends and exposure in relation to air transport sector. *Journal of Environmental Monitoring*, 6:1-11.
- Ruohomäki, K., P. Kaitaniemi, M.V. Kozlov, T. Tammaru and E. Haukioja, 1996. Density and performance of *Epirrita autumnata* (Lep., Geometridae) along three air pollution gradients in northern Europe. *Journal of Applied Ecology*, 33:773-785.
- Ruotsalainen, A.L., A.M. Markkola and M.V. Kozlov, 2006. Root fungal colonization in *Deschampsia flexuosa*: effects of pollution and neighbouring trees. *Environmental Pollution*. In press.
- Sala, O.E., F.S. Chapin III, J.J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L.F. Huenneke, R.B. Jackson, A. Kinzig, R. Leemans, D.M. Lodge, H.A. Mooney, M. Oesterheld, N.L. Poff, M.T. Sykes, B.H. Walker, M. Walker and D.H. Wall, 2000. Global Biodiversity scenarios for the year 2100. *Science*, 287:1770-1774.
- Salmi, T., A. Määttä, P. Anttila, T. Ruoho-Airola and T. Amnell, 2002. Detecting trends of annual values of atmospheric pollutants by the Mann-Kendall test and Sen's slope estimates – the Excel template application MAKESSENS, Publications on Air Quality, no. 31, FMI-AQ-31.
- Salminen, R. and V. Gregorauskiene, 2002. Geochemistry of four soil profiles from Norilsk area, Taimyr region, Russia. Geological Survey of Finland, Report No. S/41/0000/4/2002. 17p.
- Salminen, R., V. Chekushin, M. Tenhola, I. Bogatyrev, S.P. Glavatskikh, E. Fedotova, V. Gregorauskiene, G. Kashulina, H. Niskavaara, A. Polischuk, K. Rissanen, L. Selenok, O. Tomilina and L. Zhdanova, 2004. Geochemical atlas of Eastern Barents region. *Journal of Geochemical Exploration*, 83:1-3. Special issue, 530p.
- Sandgren, C.D., 1988. The ecology of chrysophyte flagellates: their growth and perennation strategies as freshwater phytoplankton. In: C.D. Sandgren (ed.). *Growth and reproductive strategies of freshwater phytoplankton*, pp. 9-104. Cambridge University Press.
- Sarmaja-Korjonen, K., M. Nyman, S. Kultti and M. Välranta, 2006. Palaeolimnological development of Lake Njargajavri, northern Finnish Lapland, in a changing Holocene climate and environment. *Journal of Paleolimnology*, 35:65-81.
- Scheuer, E., R.W. Talbot, J.E. Dibb, G.K. Seid, L. DeBel and B. Lefer, 2003. Seasonal distributions of fine aerosol sulfate in the North American Arctic basin during TOPSE. *Journal of Geophysical Research*, 108(D4):8370, doi:10.1029/2001JD001364.
- Schiff, S., M. English, M. Ecclestone, R. Elgood, M. Hinton and L. Pezzuto, 1991. Constraints on the origin of acidity in Colour Lake, Axel Heiberg Island (79°25'N). In: T.D. Prowse, C.S.L. Ommaney (eds.). *NHRI Symposium No. 6*, 532 pp. Environment Canada.
- Schindler, D.W., 1997. Widespread effects of climatic warming on freshwater ecosystems. *Hydrological Processes*, 11:1043-1067.
- Schindler, D.W., 1999. From acid rain to toxic snow. *Ambio*, 28:350-355.
- Schindler, D.W., 2001. The cumulative effects of climate warming and other human stresses on Canadian freshwaters in the new millennium. *Canadian Journal of Fisheries and Aquatic Sciences*, 58:18-29.
- Schindler, D.W. and P.J. Curtis, 1997. The role of DOC in protecting freshwaters subjected to climatic warming and acidification from UV exposure. *Biogeochemistry*, 36:1-8.
- Schindler, D.W., K.A. Kidd, D.C.G. Muir and W.L. Lockhart, 1995. The effects of ecosystem characteristics on contaminant distribution in northern freshwater lakes. *Science of the Total Environment*, 161:1-17.
- Schindler, D.W., P.J. Curtis, B.R. Parker and M.P. Stainton, 1996. Consequences of climate warming and lake acidification for UV-B penetration in North American boreal lakes. *Nature*, 379:705-708.
- Schwarz, S.E., 1979. Residence times in reservoirs under non-steady-



- state conditions: application to atmospheric SO<sub>2</sub> and aerosol sulphate. *Tellus*, 31:520-547.
- Sehmel, G.A., 1980. Particle and gas dry deposition: A review. *Atmospheric Environment*, 14:983-1011.
- Seinfeld, J.H. and S.N. Pandis, 1998. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. John Wiley & Sons, 1326p.
- Seip, H.M. and K.A., 2002. Toward an integrated approach to environmental problems. *Cicerone*, 2002 (5).
- Sellers, P., C.A. Kelly, J.W.M. Rudd and A.R. MacHutchon, 1996. Photodegradation of methylmercury in lakes. *Nature*, 380:694-697.
- Sen, P.K., 1968. Estimates of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association*, 63:1379-1389.
- Seppä, H. and J. Weckström, 1999. Holocene vegetational and limnological changes in the Fennoscandian tree-line area as documented by pollen and diatom records from Lake Tsuolbmajavri, Finland. *Ecoscience*, 6:621-635.
- Seppä, H., M. Nyman, A. Korhola and J. Weckström, 2002. Changes of treelines and alpine vegetation in relation to post-glacial climate dynamics in northern Fennoscandia based on pollen and chironomid records. *Journal of Quaternary Science*, 17:287-301.
- SFT, 2002. *Air Pollution Effects in the Norwegian – Russian Border Area. A Status Report*. Norwegian Pollution Control Authority, Oslo. 34p.
- SFT, 2004. Norwegian monitoring programme for long-range transported air pollutants. Annual report – Effects 2003. Report 913/2004. Norwegian Pollution Control Authority (SFT), Oslo. 172p.
- SFT, 2005. Monitoring of long-range transported air pollution. Annual report – Effects 2004. Report 941/2005. Norwegian Pollution Control Authority (SFT), Oslo. 149p.
- Sharma, S., D. Lavoue, H. Cachier, L.A. Barrie and S.L. Gong, 2004. Long-term trends of the black carbon concentrations in the Canadian Arctic. *Journal of Geophysical Research*, 109, doi:10.1029/2003JD004331.
- Sharma, S., E. Andrews, L.A. Barrie, J.A. Ogren, and D. Lavoues, 2006. Variations and sources of the equivalent black carbon in the High Arctic revealed by long term observations at Alert and Barrow: 1989 – 2003, *Journal of Geophysical Research*, in press.
- Sharp, M., M. Skidmore and P. Nienow, 2002. Seasonal and spatial variations in the chemistry of a High Arctic supraglacial snow cover. *Journal of Glaciology*, 48:149-158.
- Shaw, G.E., 1975. The vertical distribution of atmospheric aerosols at Barrow, Alaska. *Tellus*, 27:39-50.
- Shaw, G.E., 1981. Eddy diffusion transport of Arctic pollution from the mid-latitudes: a preliminary model. *Atmospheric Environment*, 15:1483-1490.
- Shaw, G.E., 1983. Evidence for a central Eurasian source area of Arctic Haze in Alaska. *Nature*, 299:815-818.
- Shaw, G.E., 1984. Microparticle size spectrum of Arctic haze. *Geophysical Research Letters*, 11:409-412.
- Shaw, G.E., 1987. Aerosols as climate regulators: A climate biosphere linkage? *Atmospheric Environment*, 21:985-986.
- Shaw, G.E., 1995. The Arctic Haze phenomenon. *Bulletin of the American Meteorological Society*, 76:2403-2412.
- Shaw, G.E. and K. Stamnes, 1980. Arctic haze: perturbation of the Polar radiation budget. *Annals of the New York Academy of Sciences*, 338:533-539.
- Shaw, G.E., and G. Wendler, 1972. Atmospheric turbidity at McCall Glacier in northern Alaska. *Proceedings of the Atmospheric Radiation Conference*, Fort Collins, Colorado, August 7-9, 1972.
- Shaw, G.E., K. Stamnes and Y.X. Hu, 1993. Arctic haze: Perturbation to the radiation field. *Meteorology and Atmospheric Physics*, 51:227-235.
- Shipham, M.C., A.S. Bachmeier, D.R. Cahoon Jr. and E.V. Browell, 1992. Meteorological overview of the Arctic Boundary Layer Expedition (ABLE 3A) flight series. *Journal of Geophysical Research*, 97:16395-16419.
- Simonich, S.L. and R.A. Hites, 1995. Global distribution of persistent organochlorine compounds. *Science*, 269:1851-1854.
- Simpson, R., G. Williams, A. Petroeschovsky *et al.* 2001. The assessment of the impact of air pollution on daily mortality and morbidity in Australian cities: description of a study. *Environmental Health*, 1:13-17.
- Singh, H.B., D. O'Hara, D. Herlth, J.D. Bradshaw, S.T. Sandholm, G.L. Gregory, G.W. Sachse, D.R. Blake, P.J. Crutzen and M.A. Kanakidou, 1992. Atmospheric measurements of peroxyacetyl nitrate and other organic nitrates at high latitudes: possible sources and sinks. *Journal of Geophysical Research*, 97(D15):16511-16522.
- Sirois, A. and L.A. Barrie, 1999. Arctic lower tropospheric aerosol trends and composition at Alert, Canada: 1980–1995. *Journal of Geophysical Research*, 104:11599-11618.
- Sjörs, H. and U. Gunnarsson, 2002. Calcium and pH in north and central Swedish mire waters. *Journal of Ecology*, 90:650-657.
- Skjelkvåle, B.L., J.L. Stoddard and T. Andersen, 2001a. Trends in surface waters acidification in Europe and North America (1989–1998). *Water, Air and Soil Pollution*, 130:787-792.
- Skjelkvåle, B.L., T. Andersen, E. Fjeld, J. Mannio and A. Wilander, 2001b. Recovery from acidification of lakes in Finland, Norway and Sweden 1990–1999. *Hydrology and Earth System Sciences*, 5:327-338.
- Skjelkvåle, B.L., A. Henriksen, G.S. Jönsson, J. Mannio, A. Wilander, J.P. Jensen, E. Fjeld and L. Lien, 2001c. Chemistry of lakes in the Nordic region – Denmark, Finland with Åland, Iceland, Norway with Svalbard and Bear Island, and Sweden. *Acid Rain Research Report 52/2001 SNO 4391-2001*, Norwegian Institute for Water Research (NIVA) Oslo. 39p.
- Skjelkvåle, B.L., C. Evans, T. Larssen, A. Hindar and G. Raddum, 2003. Recovery from acidification in European surface waters: a view to the future. *Ambio*, 32:170-175.
- Skjelkvåle B.L., J.L. Stoddard, D.S. Jeffries, K. Tørseth, T. Høgåsen, J. Bowman, J. Mannio, D.T. Monteith, R. Mosello, M. Rogora, D. Rzychon, J. Veselý, J. Wieting, A. Wilander and A. Worsztynowicz, 2005. Regional scale evidence for improvements in surface water chemistry 1990-2001. *Environmental Pollution*, 137:165-176.
- Sliggers, J. and W. Kakebeeke (eds.), 2004. *Clearing the Air: 25 years of the Convention on Long-range Transboundary Air Pollution*. Geneva, United Nations Economic Commission for Europe.
- Smith, F.B. and R.D. Hunt, 1978. Meteorological aspects of the transport of pollution over long distances. *Atmospheric Environment*, 12:461-477.
- Smith, S.E. and D.J. Read, 1997. *Mycorrhizal symbiosis*. 2nd ed. Academic Press. 605p.
- Smith-Sivertsen, T., V. Tchachtchine, E. Lund, V. Bykov, Y. Thomassen and T. Norseth, 1998. Urinary nickel excretion in populations living in the proximity of two Russian nickel refineries: A Norwegian-Russian Population-based Study. *Environmental Health Perspectives*, 106:503-511.
- Smith-Sivertsen, T., V. Bykov, H. Melbye, V. Tchachtchine, A. Selnes and E. Lund, 2001. Sulphur dioxide exposure and lung function in a Norwegian and Russian population living close to a nickel smelter. *International Journal of Circumpolar Health*, 60:342-359.
- Smith-Sivertsen, T., V. Tchachtchine and E. Lund, 2003. Atopy in Norwegian and Russian adults: a population-based study from the common border area. *Allergy*, 58:357.
- Smol, J.P., 1992. Paleolimnology: an important tool for effective ecosystem management. *Journal of Aquatic Ecosystem Health*, 1:49-58.
- Smol, J.P., A.P. Wolfe, H.J.B. Birks, M.S.V. Douglas, V.J. Jones, A. Korhola, R. Pienitz, K. Rühland, S. Sorvari, D. Antoniades, S.J. Brooks, M.-A. Fallu, M. Hughes, B.E. Keatley, T.E. Laing, N. Michelutti, L. Nazarova, M. Nyman, A.M. Paterson, B. Perren, R. Quinlan, M. Rautio, E. Saulnier-Talbot, S. Siitonen, N. Solovieva and J. Weckström, 2005. Climate-Driven Regime Shifts in Arctic Lake Ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, 102:4397-4402.
- Solberg, S., C. Dye and N. Schmidbauer, 1996. Carbonyls and nonmethane hydrocarbons at rural European sites from the Mediterranean to the Arctic. *Journal of Atmospheric Chemistry*, 25:33-66.
- Solberg, S., B.L. Skjelkvåle, P.A. Aarrestad, O. Reitan and S.E. Walker, 1999. Regional konsekvensutredning for oljevirksomheten i Nordjøn. Report 5: Regulære utslipp til luft – konsekvenser. (In Norwegian)
- Solberg, S., M. Lazaridis, S.-E. Walker, S. Knudsen and A. Semb, 2003. The contribution to nitrogen deposition and ozone formation in south Norway from atmospheric emissions related to the petroleum activity in the North Sea. *Water, Air and Soil Pollution*, 148:289-321.
- Solovieva, N., 2000. A palaeolimnological study of Holocene environmental change in a small upland lake from the Kola Peninsula, Russia. Ph.D. Thesis, University College London. 413p.
- Solovieva, N. and V.J. Jones, 2002. A multiproxy record of Holocene environmental changes in the Kola Peninsula, northwest Russia. *Journal of Quaternary Science*, 17:303-318.
- Solovieva, N., V.J. Jones, P.G. Appleby and B.M. Kondratenok, 2002. Extent, environmental impact and long-term trends in atmospheric contamination in the Usa Basin of East-European Russian Arctic. *Water, Air and Soil Pollution*, 139:237-260.
- Solovieva, N., V.J. Jones, L. Nazarova, S.J. Brooks, H.J.B. Birks, J.-A. Grytnes, P.G. Appleby, T. Kauppila, B. Kondratenok, I. Renberg and V. Ponomarev, 2005. Palaeolimnological evidence for recent climatic change in lakes from the northern Urals, Arctic Russia. *Journal of Paleolimnology*, 33:463-482.
- Sorvari, S., A. Korhola and R. Thompson, 2002. Lake diatom response to recent Arctic warming in Finnish Lapland. *Global Change Biology*, 8:171-181.
- Sprenger, M. and H. Wernli, 2003. A northern hemispheric climatology of cross tropopause exchange for the ERA15 time period (1979–1993). *Journal of Geophysical Research*, 108:8521, doi:10.1029/2002JD002636.
- Steinnes, E., 1995. A critical evaluation of the use of naturally growing

- moss to monitor the deposition of atmospheric metals. *Science of the Total Environment*, 160/161:243-249.
- Steinnes, E., N. Lukina, V. Nikonov, D. Aamlid and O. Røyset, 2000. A gradient study of 34 elements in the vicinity of a copper-nickel smelter in the Kola Peninsula. *Environmental Monitoring and Assessment*, 60:71-88.
- Stocks, B.J., M.A. Fosberg, T.J. Lynham, L. Mearns, B.M. Wotton, Q. Yang, J.-Z. Jin, K. Lawrence, G.R. Hartley, J.A. Mason and D.W. McKennedy, 1998. Climate change and forest fire potential in Russian and Canadian boreal forests. *Climatic Change*, 38:1-13.
- Stoddard, J.L., D.S. Jeffries, A. Lükewille, T.A. Clair, P.J. Dillon, C.T. Driscoll, M. Forsius, M. Johannessen, J.S. Kahl, J.H. Kellogg, A. Kemp, J. Mannio, D. Monteith, P.S. Murdoch, S. Patrick, A. Rebsdorf, B.L. Skjelkvåle, M.P. Stainton, T.S. Traaen, H. van Dam, K.E. Webster, J. Wieting and A. Wilander, 1999. Regional trends in aquatic recovery from acidification in North America and Europe 1980-95. *Nature*, 401:575-578.
- Stoddard, J.L., J.S. Kahl, F.A. Deviney, D.R. DeWalle, C.T. Driscoll, A. T. Herlihy, J.H. Kellogg, P.S. Murdoch, J.R. Webb and K.E. Webster, 2003. Response of Surface Water Chemistry to the Clean Air Act Amendments of 1990. EPA/620/R-03/001, U.S. Environmental Protection Agency. 92p.
- Stohl, A., 2001. A one-year Lagrangian "climatology" of airstreams in the northern hemisphere troposphere and lowermost stratosphere. *Journal of Geophysical Research*, 106:7263-7279.
- Stohl, A., 2006. Characteristics of atmospheric transport into the Arctic troposphere. *Journal of Geophysical Research*, 111. No. D11, D11306 10.1029/2005JD006888.
- Stohl, A., S. Eckhardt, C. Forster, P. James and N. Spichtinger, 2002. On the pathways and timescales of intercontinental air pollution transport. *Journal of Geophysical Research*, 107:4684, doi: 10.1029/2001JD001396.
- Sturges, W.T. and L.A. Barrie, 1988. Chlorine, bromine, and iodine in Arctic aerosols. *Atmospheric Environment*, 22:1179-1194.
- Sturges, W.T. and G.E. Shaw, 1983. Halogens in aerosols in central Alaska. *Atmospheric Environment*, 27A:2969-2977.
- Suomela, J. and P. Palokangas, 1993. Effects of aerial pollutants on plant quality for small northern mammalian herbivores. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 377-378. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Swart, R., M. Amann, F. Raes and W. Tuinstra, 2004. A good climate for clean air: Linkages between climate change and air pollution. *Climatic Change*, 66:263-269.
- Tammi, J., M. Appelberg, U. Beier, T. Hesthagen, A. Lappalainen and M. Rask, 2003a. Fish status of Nordic lakes: effects of acidification, eutrophication and stocking activity on present fish species composition. *Ambio*, 32:98-105.
- Tammi, J., A. Lappalainen and T. Bergman, 2003b. Water quality and fish populations of acid sensitive waters in the Vätsäri area, north-eastern Finland: responses to reduced sulphur emissions from the Kola Peninsula, Russia, in the 1990s. *Boreal Environment Research*, 8:1-7.
- Targulian, V.O., 1971. Soil formation and weathering in cold humid regions. Moscow: "Nauka". 268p. (In Russian)
- Tarhanen, S., S. Metsärinne, T. Holopainen and J. Oksanen, 1999. Membrane permeability response of lichen *Bryoria fuscescens* to wet deposited heavy metals and acid rain. *Environmental Pollution*, 104:101-129.
- Tenow, O. and H. Bylund, 2000. Recovery of a *Betula pubescent* forest in northern Sweden after severe defoliation by *Epirrita autumnata*. *Journal of Vegetation Science*, 11:855-862.
- The Royal Society, 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Policy document 12/05.
- Thurston, D., 2003. Offshore Oil and Gas Activities in the Arctic. PAME Background paper.
- Thurston, G.D. and K. Ito, 2001. Epidemiological studies of acute ozone exposures and mortality. *Journal of Exposure Analysis and Environmental Epidemiology*, 11:286-294.
- Tikkanen, E. and I. Niemelä, 1995. Kola Peninsula pollutants and forest ecosystems in Lapland. Final Report of the Lapland Forest Damage Project. Finland's Ministry of Agriculture and Forestry and The Finnish Forest Research Institute, Gummerus, Jyväskylä, 82p.
- Tømmervik, H., B.E. Johansen and J.P. Pedersen, 1995. Monitoring the effects of air pollution on terrestrial ecosystems in Varanger (Norway) and Nikel-Pechenga (Russia) using remote sensing. *Science of the Total Environment*, 160/161:753-767.
- Tømmervik, H., M.E. Johansen, J.P. Pedersen and T. Guneriusson, 1998. Integration of remote sensed and in-situ data in an analysis of the air pollution effects on terrestrial ecosystems in the border areas between Norway and Russia. *Environmental Monitoring and Assessment*, 49:51-85.
- Tømmervik, H., K.A. Högda and I. Solheim, 2003. Monitoring vegetation changes in Pasvik (Norway) and Pechenga in Kola Peninsula (Russia) using multitemporal Landsat MSS/TM data. *Remote Sensing of Environment*, 85:370-388.
- Tømmervik, H., B. Johansen, I. Tombre, D. Thannheiser, K.A. Högda, E. Gaare and F.E. Wielgolaski, 2004. Vegetation changes in the Nordic mountain birch forest: the influence of grazing and climate change. *Arctic, Antarctic, and Alpine Research*, 36:323-332.
- Traaen, T.S., 1987. Forsuring av innsjøer i Finnmark. SFT-Rapport 299/87, SFT, Oslo.
- Traaen, T.S., 1991. Forsuring og tungmetallforurensning i Sør-Varanger. Framdriftsrapport for 1990. SFT-Rapport 481/92, SFT, Oslo.
- Tranvik, L.J. and M. Jansson, 2002. Climate change – Terrestrial export of organic carbon. *Nature*, 415:861-862.
- Treffeisen, R., A. Herber, J. Ström, M. Shiobara, T. Yamanouchi, S. Yamagata, K. Holmén, M. Kriews and O. Schrems, 2004. Interpretation of Arctic aerosol properties using cluster analysis applied to observations in the Svalbard area. *Tellus*, 56B:457-476.
- Trivett, N.B.A., L.A. Barrie, J.P. Blanchet, R.M. Bottenheim, R.M. Hoff and R.E. Mickle, 1989. An experimental investigation of Arctic haze at Alert, NWT, March 1985. *Atmosphere-Ocean*, 26:341-376.
- Tsvetkov, V.F., 1993. Threshold levels of air pollution and forest decline in surroundings of "Severonikel" smelter complex. In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.). *Aerial pollution on Kola Peninsula*, pp. 397-401. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Tuovinen, J.-P., T. Laurila, H. Lättilä, A. Ryaboshapko, P. Brukhanov and S. Korolev, 1993. Impact of the sulphur dioxide sources in the Kola Peninsula on air quality in northernmost Europe. *Atmospheric Environment*, 27A:1379-1395.
- Tutubalina, O.V. and W.G. Rees, 2001. Vegetation degradation in a permafrost region as seen from space: Noril'sk (1961-1999). *Cold Regions Science and Technology*, 32:191-203.
- Twomey, S., 1977. The influence of pollution on the shortwave albedo of clouds. *Journal of the Atmospheric Sciences*, 34:1149-1152.
- Twomey, S., 1991. Aerosols, clouds and radiation. *Atmospheric Environment*, 25A:2435-2442.
- UBA, 2004. Manual on methodologies and criteria for modelling and mapping critical loads & levels and air pollution effects, risks and trends. Texte 52/04, Umweltbundesamt, www.icpmapping.org
- UNECE, 1993. The 1993 report of the task force on mapping. EB.AIR/WG.1/R.85
- UNECE, 1994. Protocol to the 1979 Convention on Long-range Transboundary Air Pollution on Further Reduction of Sulphur Emissions. Document ECE/EB.AIR/40, United Nations Economic Commission for Europe, 106p.
- UNECE, 1999. The 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone. Document ECE/EB.AIR, United Nations Economic Commission for Europe.
- UNECE, 2004a. Mapping Manual 2004. UNECE Convention on Long-range Transboundary Air Pollution. Manual on methodologies and criteria for modelling and mapping critical loads and levels and air pollution effects, risks and trends.
- UNECE, 2004b. Review and assessment of air pollution effects and their recorded trends. Report of the Working Group on Effects of the Convention on Long-range Transboundary Air Pollution. United Nations Economic Commission for Europe.
- Valero, F.P.J., T.P. Ackerman and W.J.R. Gore, 1989. The effects of the arctic haze as determined from airborne radiometric measurements during AGASP II. *Journal of Atmospheric Chemistry*, 9:225-244.
- Valkama, J. and M.V. Kozlov, 2001. Impact of climatic factors on the developmental stability of mountain birch growing in a contaminated area. *Journal of Applied Ecology*, 38:665-673.
- van Rompaey, R.S.A.R., 1995. Acidification interacting with global changes: research to manage drifting systems. In: G.J. Heij and J.W. Erisman (eds.). *Acid rain research: do we have enough answers?* pp. 381-384. Elsevier Science.
- Vassilieva, N., M. Gytarsky and R. Karaban, 1995. Changes of the dominant species of the epiphytic lichens and ground vegetation cover in the forests subjected to "Pechenganicel" smelter impact. In: E. Løbersli and K. Venn (eds.). *Effect of air pollution on terrestrial ecosystems in the border area between Norway and Russia*, pp. 103-111. Proceedings from the second symposium, Svanvik, Norway, 3-5 October 1994. Directorate for Nature Management, Trondheim.
- Vestreng, V., 2003. Review and revision of emission data reported to CLRTAP. EMEP/ MSC-W Note1. Norwegian Meteorological Institute.
- Vinebrooke, R.D., K.L. Cottingham, J. Norberg, M. Scheffer, S.I. Dodson, S.C. Maberly and U. Sommer, 2004. Impacts of multiple stressors on biodiversity and ecosystem functioning: the role of species co-tolerance. *OIKOS*, 104:451-457.
- Virtanen, T., K. Mikkola, E. Patova and A. Nikula, 2002. Satellite image analysis of human caused changes in the tundra vegetation around the city of Vorkuta, North-European Russia. *Environment*

## References

- tal Pollution, 120:647-658.
- Vollenweider, R.A. and J. Kerekes, 1982. Eutrophication of waters, monitoring, assessment and control. Organization for Economic Cooperation and Development, Paris, 154p.
- VTI/IVL, 2004. Data on SO<sub>x</sub> and NO<sub>x</sub> emissions from Russian Coal Fired Power Plants for the year 2000; compiled by the VTI All Russia Thermal Engineering Institute, for the project 'Mercury Co-Benefit Activities with SO<sub>2</sub> Control Technology Development in Russia' led by the Swedish Environmental Research Institute (IVL) and funded by the US EPA Office of International Affairs International Cooperation.
- Vuorenmaa, J., M. Forsius and J. Mannio, 2006. Increasing trends of total organic carbon concentrations in small forest lakes in Finland from 1987 to 2003. *Science of the Total Environment*, 365:47-65.
- Waggoner, A.P. and R.E. Weiss, 1980. Comparison of the fine particle mass concentration and light scattering extinction in ambient aerosol. *Atmospheric Environment*, 14:623-626.
- Wählin, P., F. Palmgren and R.V. Dingenen, 2002. Experimental studies of ultrafine particles in streets and the relationship to traffic. *Atmospheric Environment*, 35(Suppl. 1):S63-S69.
- Walker, T.R., P.D. Crittenden and S.D. Young, 2003a. Regional variation in the chemical composition of winter snow pack and terricolous lichens in relation to sources of acid emissions in the Usa river basin, northeast European Russia. *Environmental Pollution*, 125:401-412.
- Walker, T.R., S.D. Young, P.D. Crittenden and H. Zhang, 2003b. Anthropogenic metal enrichment of snow and soil in North-Eastern European Russia. *Environmental Pollution*, 121:11-21.
- Wania, F., 2003. Assessing the potential of persistent organic chemicals for long-range transport and accumulation in polar regions. *Environmental Science and Technology*, 37:1344-1351.
- Warren, S.G. and W.J. Wiscombe, 1980. A model for the spectral albedo of snow. II. Snow containing atmospheric aerosols. *Aerosol Science and Technology*, 4:31-43.
- Watanabe, O., H. Motoyama, M. Igarashi, K. Kamiyama, S. Matoba, K. Goto-Azuma, H. Narita, H. Kameda, 2001. Studies on climatic and environmental changes during the last few hundred years using ice cores from various sites in Nordaustlandet, Svalbard. *Mem. Natl. Inst. Polar Research*, 54: 227-242.
- Weatherhead, B., A. Tanskanen and A. Stevermer, 2005. Chapter 5. Ozone and ultraviolet radiation. *In: Arctic Climate Impact Assessment*, pp. 151-182. Cambridge University Press.
- Weckström, J., A. Korhola and T. Blom, 1997. Diatoms as quantitative indicators of pH and water temperature in subarctic Fennoscandian lakes. *Hydrobiologia*, 347:171-184.
- Weckström, J., A. Korhola, J.A. Snyder, T.E. Laing and G.M. MacDonald, 2003. Diatom inferred acidity history of 32 lakes on the Kola Peninsula, Russia. *Water, Air and Soil Pollution*, 149:339-361.
- Wesely, M.L. and B.B. Hicks, 2000. A review of the current status of knowledge on dry deposition. *Atmospheric Environment*, 34:2261-2282.
- WHO, 1996. Diesel fuel and exhausts emissions. *Environmental health criteria No 171*. World Health Organization.
- WHO, 1997. Health and environment in sustainable development. (WHO/EHG/97.8). World Health Organization.
- WHO, 2001. Air quality guidelines for Europe. Updated version. World Health Organization.
- Wigington, P.J., T.O. Davices, M. Tranter and K.N. Eshleman, 1992. Comparison of episodic acidification in Canada, Europe and the United States. *Environmental Pollution*, 78:29-35.
- Wilander, A., 1994. Estimation of ground sulphate in natural surface waters in Sweden. *Water Air Soil Pollution*, 75:371-387.
- Wilander, A., 1998. Water chemistry in reference lakes during 12 years; status and trends. Report 4652, Naturvårdsverket Förlag, Stockholm, Sweden. 79p. (In Swedish)
- Willén, E., 2003. Dominance patterns of planktonic algae in Swedish forest lakes. *Hydrobiologia*, 502:315-324.
- Wolfe, A., 2002. Climate modulates the acidity of Arctic lakes on millennial time scales. *Geology*, 3:215-218.
- Wotawa, G., P.C. Novelli, M. Trainer and C. Granier, 2001. Interannual variability of summertime CO concentrations in the Northern Hemisphere explained by boreal forest fires in North America and Russia. *Geophysical Research Letters*, 28:4575-4578.
- Wright, R.F. and A. Jenkins, 2001. Climate change as a confounding factor in reversibility of acidification: RAIN and CLIMEX projects. *Hydrology and Earth System Sciences*, 5:477-486.
- Wright, R.F., L. Camarero, B.J. Cosby, R.C. Ferrier, M. Forsius, R. Hellwell, A. Jenkins, J. Kopáček, T. Larssen, V. Majer, F. Moldan, M. Posch, M. Rogora and W. Schöpp, 2005. Recovery of acidified European surface waters. *Environmental Science and Technology*, 39:64A-72A.
- Wrona, F.J., T.D. Prowse and J.D. Reist, 2005. Chapter 8. Freshwater ecosystems and fisheries. *In: Arctic Climate Impact Assessment*, pp. 353-452. Cambridge University Press.
- WWF, 2005. Areas vulnerable to acute oil pollution in the Norwegian Barents Sea. Report: 2005-0456. World Wide Fund for Nature.
- Yakovlev, V.A., 1992. Benthic invertebrates, zooplankton and phytoplankton communities in relation to pollution of waters in the Kola Peninsula. *In: K. Kinnunen and M. Varmola (eds.)*. Effects of air pollutants and acidification in combination with climatic factors on forests, soils and waters in Northern Fennoscandia, pp. 109-114. Nordic Council of Ministers, Copenhagen.
- Yakovlev, V.A., 1999. Acidity of small lakes in Finnish Lapland – based on aquatic macroinvertebrate studies in 1993-1995. Publication/brochure 1/18/1999, Lapland Regional Environmental Centre, Rovaniemi.
- Yakovlev, V.A., 2000. Trophic structure of zoobenthos as an ecological indicator for aquatic ecosystems and a water quality index. *Vodnye Resursy (Water Resources)*, 27:237-244.
- Zagurskaya, L.M. and S.S. Ziabchenko, 1994. Impact of industrial pollution on microbial processes in boreal forests of Kostomuksha region. *Pochvovedenie (Eurasian Soil Science)*, 5:105-110. (In Russian)
- Zenkova, I.V., 1999. The state of soil fauna in the impact zone of the 'Severonikel' smelter. *In: Y.A. Izrael, G.V. Kalabin and V.V. Nikonov (eds.)*. Anthropogenic impact on northern nature and its ecological consequences, pp. 293-310. Kola Science Centre, Apatity. (In Russian)
- Zenkova, I.V. and E.M. Zainagutdinova, 2002. Industrial pollution of soils and individual parameters of soil inhabited group of beetles. *In: N.B. Hitrov (ed.)*. Steadiness of soils to natural and human-induced impacts, pp. 257-258. Abstracts to presentations of All-Russian conference, 24-25 April 2002, Moscow. (In Russian)
- Zhang, T., K. Stamnes and S.A. Bowling, 1996. Impact of clouds on surface radiative fluxes and snowmelt in the Arctic and subarctic. *Journal of Climate*, 9:2110-2123.
- Zvereva, E.L., 1993a. Population densities of some flies in an air pollution gradient near Monchegorsk. *In: M.V. Kozlov, E. Haukioja and V.T. Yarmishko (eds.)*. Aerial pollution on Kola Peninsula, pp. 371-373. Proceedings of the International Workshop, April 14-16 1992, St. Petersburg. Kola Science Centre, Apatity.
- Zvereva, E.L., 1993b. Effect of industrial pollution on fly communities (Diptera, Brachycera). *Entomol. Obozr.*, 72:558-569. (In Russian; English translation: *Ent. Rev.* 1994, 73(2):45-57).
- Zvereva, E.L., 1999. Can global warming promote insect outbreaks by changing host plant quality? *In: M.A. Lange, B. Bartling and K. Grosfeld (eds.)*. Global Changes and the Barents Sea Region. Proceedings of the First International BASIS Research Conference, pp 393-394. St.Petersburg, February 22-25 1998.
- Zvereva, E.L. and M.V. Kozlov, 2000. Natural enemies of the leaf beetle, *Melasoma lapponica*, in the impact zone of a nickel-copper smelter: density dependence and effects of air pollution. *Journal of Applied Ecology*, 37:298-308.
- Zvereva, E.L. and M.V. Kozlov, 2001. Effects of pollution-induced habitat disturbance on the response of willows to simulated herbivory. *Journal of Ecology*, 89:21-30.
- Zvereva, E.L. and M.V. Kozlov, 2005. Growth and reproduction of dwarf shrubs, *Vaccinium myrtillus* and *V. vitis-idaea*, in a severely polluted area. *Basic and Applied Ecology*, 6:261-274.
- Zvereva, E., M.V. Kozlov and S. Neuvonen, 1995a. Population density and performance of *Melasoma lapponica* (Coleoptera: Chrysomelidae) in surroundings of smelter complex. *Environmental Entomology*, 24:707-715.
- Zvereva, E.L., M.V. Kozlov and S. Neuvonen, 1995b. Decrease in feeding niche breadth of *Melasoma lapponica* (Coleoptera: Chrysomelidae) with increase in pollution. *Oecologia*, 104:323-329.
- Zvereva, E.L., M.V. Kozlov, P. Niemelä and E. Haukioja, 1997a. Delayed induced resistance and increase in leaf fluctuating asymmetry as responses of *Salix borealis* to insect herbivory. *Oecologia*, 102:368-373.
- Zvereva, E.L., M.V. Kozlov and E. Haukioja, 1997b. Population dynamics of a herbivore in an industrially modified landscape: case study with *Melasoma lapponica* (Coleoptera: Chrysomelidae). *Acta Phytoipatologica et Entomologica Hungarica*, 32:251-258.
- Zvereva, E.L., M.V. Kozlov and O. Yu. Kruglova, 2002. Color polymorphism in relation to population dynamics of the leaf beetle *Chrysomela lapponica*. *Evolutionary Ecology*, 16:523-539.
- Zvereva, E.L., V. Serebrov, V. Glupov and I. Dubovsky, 2003. Activity and heavy metal resistance of non-specific esterases in leaf beetle *Chrysomela lapponica* from polluted and unpolluted habitats. *Comparative Biochemistry and Physiology, Part C* 135:383-391.






## Abbreviations

*	of non-marine origin
1-D	one-dimensional
3-D	three-dimensional
Al	aluminum
AMAP	Arctic Monitoring and Assessment Programme
ANC	acid neutralizing capacity
AO	Arctic Oscillation
AOD	aerosol optical depth
BC	black carbon / base cation
C	carbon
Ca	calcium
C-horizon	parent material of soil
CH <sub>4</sub>	methane
Cl	chloride
CL <sub>Ac</sub>	critical load of acidity
CLE scenario	emissions scenario based on current legislation
CO	carbon monoxide
CO <sub>2</sub>	carbon dioxide
Cu	copper
DEHM	Danish Eulerian Hemispheric Model
DIC	dissolved inorganic carbon
DI-pH	diatom-inferred pH
DOC	dissolved organic carbon
EANET	Acid Deposition Monitoring Network in East Asia
EMEP	Co-operative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe
K	potassium

LRTAP Convention	Convention on Long-Range Transboundary Air Pollution
MFR scenario	emissions scenario based on maximum technically feasible reductions
Mg	magnesium
Mn	manganese
N	nitrogen
Na	sodium
NAO	North Atlantic Oscillation
N <sub>2</sub> O	nitrous oxide
NH <sub>3</sub>	ammonia
NH <sub>4</sub>	ammonium
NH <sub>4</sub> -N	ammonium nitrogen
Ni	nickel
NILU	Norwegian Institute for Air Research
nmVOC	non-methane volatile organic compounds
NO <sub>2</sub>	nitrogen dioxide
NO <sub>3</sub>	nitrate
NO <sub>3</sub> -N	nitrate nitrogen
NO <sub>x</sub>	nitrogen oxides
NO <sub>x</sub> -N	nitrogen oxides nitrogen
NSR	Northern Sea Route
O-horizon	organic horizon of soil
S	sulfur
SO <sub>2</sub>	sulfur dioxide
SO <sub>4</sub>	sulfate
SO <sub>4</sub> -S	sulfate sulfur
SO <sub>x</sub>	sulfur oxides
UN ECE	UN Economic Commission for Europe
UV	ultraviolet
Zn	zinc

The image is a cover page for the Arctic Monitoring and Assessment Programme (AMAP). The left side is white, and the right side is a dark, blurred photograph of a forest floor with green moss and brown tree trunks. A light blue curved line is at the bottom left. The AMAP logo is in the bottom left corner.

**AMAP**  
Arctic Monitoring and  
Assessment Programme

ISBN 82-7971-046-9